



HNF1 α and CDX2 Transcriptional Factors bind to Cadherin-17 (CDH17) gene promoter and modulate its expression in Hepatocellular Carcinoma

Journal:	<i>Journal of Cellular Biochemistry</i>
Manuscript ID:	JCB-09-0616
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	09-Nov-2009
Complete List of Authors:	Zhu, Rui; The University of Hong Kong, Surgery Wong, Kwong-Fai; The University of Hong Kong, Surgery Lee, Nikki; The University of Hong Kong, Surgery Lee, Kai-Fai; The University of Hong Kong, Obstetrics and Gynaecology Luk, John; University of Hong Kong, Department of Surgery
Keywords:	Cadherin-17, Hepatocellular carcinomas, Hepatic nuclear factor 1 α , Caudal-related homeobox 2, Promoter



1
2
3
4 **HNF1 α and CDX2 Transcriptional Factors bind to Cadherin-17 (CDH17)**
5
6 **gene promoter and modulate its expression in Hepatocellular Carcinoma**
7
8

9
10
11 **Rui Zhu¹, Kwong-Fai Wong¹, Nikki P.Y. Lee¹, Kai-Fai Lee^{2*} and John M.C. Luk^{1*}**
12
13

14
15
16 ¹*Department of Surgery and Center for Cancer Research, The University of Hong Kong,*
17 *Pokfulam, Hong Kong, China*

18
19 ²*Department of Obstetrics and Gynaecology, The University of Hong Kong, Pokfulam,*
20 *Hong Kong, China*
21
22
23

24
25
26 **Correspondence to:**

27
28 Dr JMC Luk, Department of Surgery, The University of Hong Kong, Pokfulam, Hong
29 Kong. Tel: +852-2819-9628; Fax: +852-2819-9636; E-mail: jmluk@hku.hk or

30
31 Dr KF Lee, Department of Obstetrics and Gynaecology, The University of Hong Kong,
32 Pokfulam, Hong Kong. Tel: +852-2819-9369; fax: +852-2816-1947; E-mail:
33 ckflee@hkucc.hku.hk
34
35
36
37
38
39

40 **Running head: HNF1 α and CDX2 regulate CDH17**
41
42

43 **Keywords: Cadherin-17; hepatocellular carcinomas; hepatic nuclear factor 1 α ;**
44 **caudal-related homeobox 2; promoter**
45
46
47
48
49

50 **Number of Figures: 5**
51

52
53 **Number of Table: 1**
54
55
56
57
58
59
60

ABSTRACT

Cadherin-17 (CDH17) is a novel adhesion molecule that expresses abundantly in fetal liver but is silenced in healthy adult. Recently, we have identified CDH17 is a cancer biomarker which is over-expressed in ~80% of hepatocellular carcinomas that associated with poor clinical outcomes. The present study aims to identify the transcription factor(s) and elucidate the regulatory mechanisms that drive CDH17 gene over-expression in human HCC. A 1-kb upstream sequence of CDH17 gene was cloned and the promoter activity was studied by luciferase reporter assay. By bioinformatics analysis, deletion and mutation assays, and chromatin immunoprecipitation studies, we identified hepatic nuclear factor 1 α (HNF1 α) and caudal-related homeobox 2 (CDX2) binding sites at the proximal (-81/+55 region relative to the transcriptional start site) promoter region which modulate the CDH17 promoter activities in two HCC cell lines (Hep3B and MHCC97L). Interestingly, the expression of CDH17 was high in mouse fetal livers (E13.5> E16.5> newborn> 3 weeks> adult) as well as in human HCC cell lines with higher metastatic potential (MHCC97L> Hep3B> MIHA), which is in concomitant with the expression levels of HNF1 α and CDX2 mRNA and proteins. Furthermore, knockdown of HNF1 α and CDX2 by small interfere RNA (siRNA) significantly decrease CDH17 transcript level in the transfected HCC cells. In sum, we identified -81/+55 region of CDH17 as minimal promoter region that is regulated by HNF1 α and CDX2. The expression levels of these transcriptional factors as well as CDH17, may serve as biomarkers for HCC, and may be potential targets for cancer treatment.

INTRODUCTION

Hepatocellular carcinoma (HCC) ranks the fifth most common malignancy in the world [Llovet et al., 2003]. Hepatic resection and liver transplantation are the mainstays of surgical management for HCC, but the 5-year survival rate is still poor [Liu and Fan, 2006; Poon and Fan, 2004]. Early diagnosis and treatment of HCC remain challenging due to lack of highly specific and sensitive markers. The currently used serological tumor marker for HCC, α -fetoprotein (AFP), lacks specificity and sensitivity [Johnson, 2001].

Liver-Intestine cadherin (LI-cadherin), or CDH17 (cadherin-17) is a member of non-classical cadherin family [Jung et al., 2004] and located on chromosome 8q21.1 [Dantzig et al., 1994]. Beside the cell adhesion function, CDH17 was initially identified as a proton-dependent peptide transporter-1 in human colon adenocarcinoma Caco-2 cells [Dantzig et al., 1994]. CDH17 is over-expressed in gastric metaplasia and gastric carcinoma [Grotzinger et al., 2001; Kaposi-Novak et al., 2006; Ko et al., 2004; Ko et al., 2005], as well as in pancreatic carcinoma [Takamura et al., 2003]. Recently, our group has reported CDH17 is over-expressed in HCC and is associated with poor survival rate and a higher tumor recurrence [Wang et al., 2005; Wong et al., 2003]. In human, over-expression of CDH17 is frequently detected in ~80% HCC cases, but the corresponding regulatory mechanisms remain largely unclear.

Recently, CDX2 was found to play important role(s) in up-regulation of CDH17 expression in colorectal cancer cell lines and gastric adenocarcinoma [Hinoi et al., 2002; Ko et al., 2005], which was in turn was regulated by bone morphogenetic protein 2 (BMP2)

1
2
3
4 and BMP4 in gastric cancer cells [Barros et al., 2008].
5
6

7 HNF1 is one of the key transcriptional factors that regulate serum α -fetoprotein
8
9 levels in hepatocellular carcinoma [Nakabayashi et al., 2004]. It is predominantly
10
11 expressed in liver and kidney, and binds to the promoters of a variety of genes that are
12
13 expressed exclusively in the liver including fibrinogen- α and - β , albumin, α -1-antitrypsin,
14
15 liver-type pyruvate kinase, transthyretin, aldolase B, and hepatitis B virus large surface
16
17 protein [Courtois et al., 1987; Xu et al., 2001]. Over-expression of HNF1 α was found in
18
19 well-differentiated HCC than in the surrounding non-HCC tissues [Wang et al., 1998].
20
21
22
23
24

25 In this study, we aimed to dissect the regulation of CDH17 expression in HCC
26
27 and in mouse liver during development, and functionally characterized two transcription
28
29 factors (HNF1 α and CDX2) on CDH17 expression. The specific regulation of these
30
31 transcriptional factors was confirmed by RNAi approach in the transfected HCC cells.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

MATERIALS AND METHODS

Plasmids, bacteria and reagents

The Dual-Luciferase® reporter assay system and the pGL3-Basic, pGL3-Control and pRL-TK reporter vectors were obtained from Promega (Madison, WI). *E. coli* DH5 α competent cells were prepared by chemical method. Restriction endonucleases, T4 DNA ligase and T4 polynucleotide kinase were purchased from New England Biolabs (Ipswich, MA). Bradford reagent was purchased from Bio-Rad (Hercules, CA) and all other reagents were purchased from Sigma (St. Louis, MO).

Mouse liver and clinical HCC tissues

The use of animal in this study protocol was approved by the Using Live Animals for Teaching and Research (CULATR), The University of Hong Kong. Informed written consent was obtained from patients and the usage of clinical tissues for this study was vetted and approved by the Institutional Review Board of the University of Hong Kong/Hospital Authority Hong Kong West Cluster. Messenger RNA and protein lysates were extracted from BALB/c mouse livers at different stages: embryonic day 13.5 (E13.5), E16.5, new-borne (NB), postnatal (3 weeks), and adult (3 months).

Cloning of 5'-promoter region of the CDH17 gene

The promoter region of the human CDH17 gene (NM_004063.3) was predicted and analyzed using the NCBI Map Viewer (<http://www.ncbi.nlm.nih.gov/mapview>). The

1
2
3
4 promoter fragment starting at -909bp and ending at +55bp was amplified by PCR using
5
6 specific forward primer 5'-TAGAGAGTGGGCTGGGCTC-3' and reverse primer
7
8 5'-TGGTCGAGACTCTTGCTACG-3' and Pfu Turbo high-fidelity DNA polymerase
9
10 (Stratagene, La Jolla, CA). In brief, PCR was carried out at 94 °C for 5 min and then 30
11
12 cycles at 94 °C for 40 s, 60 °C for 1 min, and 72 °C for 3 min and a final extension step at
13
14 72 °C for 10 min in a final reaction volume of 25 μ l. The genomic DNA of HCC cell line
15
16 MHCC97L was used as template and the CDH17 gene promoter was cloned by directional
17
18 restriction cloning method [Lee et al., 2009; Wong et al., 2003].
19
20
21
22
23
24
25
26
27

28 **Constructs of CDH17 mutant promoters**

29
30 Another four deletion constructs (-686/+55, -332/+55, -81/+55 and -40/+55) were
31
32 generated by PCR cloning with the following forward primers
33
34 (5'-GAAGCCTTGACTTGAGAAAT-3', 5'-TGATCAAGTCTCCTGTGCT-3',
35
36 5'-AGTGGCTCTCGAAGAGCAAT-3', and 5'-TTTGACTGAAGCTGAAGGG-3'),
37
38 respectively. These five CDH17 promoter constructs were cloned to the pGL3-Basic vector
39
40 by XhoI/KpnI restriction digestions. For site-directed mutagenesis, mutated promoter
41
42 constructs were generated using the QuickChange mutagenesis kit (Stratagene). All
43
44 deletion and point-mutation constructs were sequenced with an ABI 3700 Genetic Analyzer
45
46 (Applied Biosystems Inc., Foster City, CA).
47
48
49
50
51
52
53
54
55
56

57 **Cell Cultures and transfection**

58
59 The human HCC cell lines MHCC97L and Hep3B and a normal immortalized human
60

1
2
3
4 hepatocyte cell line MIHA were cultured in Dulbecco's minimal essential medium (Life
5
6 Technologies, Carlsbad, CA), supplemented with 10% fetal bovine serum, 50 U/ml
7
8 penicillin, 50 U/ml streptomycin and 2 mM L-glutamine at 37 °C and 5% CO₂. For
9
10 transfection, cells were grown to about 90% confluence in a 24-well tissue culture plate and
11
12 then added with 500 µl of serum-free medium containing 2 µl lipofectamine™ 2000
13
14 reagent (Invitrogen, Carlsbad, CA) mixed with 1 µg of purified plasmid cDNA and 0.1 µg
15
16 of pRL-TK vector (the internal control reporter) per well. Followed by incubation at 37°C
17
18 for 48 h, the promoter activity of the CDH17 gene was assessed by measuring luciferase
19
20 activity using the Dual-Luciferase® reporter assay (Promega). The intensity of
21
22 chemiluminescence was measured in a luminometer (Thermo, Waltham, MA). All
23
24 experiments were performed in duplicate and repeated at least three times with positive
25
26 (pGL3-Control) and negative (pGL3-Basic) controls.
27
28
29
30
31
32
33
34
35
36
37
38

39 **Western blotting**

40
41 Antibodies against CDX2 and β-actin were purchased from Millipore (Billerica, MA) and
42
43 Sigma (St Louis, MO), respectively. Antibodies against CDH17, HNF1α and HNF1β
44
45 were from Santa Cruz Biotechnology (Santa Cruz, CA). Protein extracts from the liver of
46
47 BALB/c mice at different developmental stages and from different cell lines MHCC97L,
48
49 Hep3B and MIHA was prepared. In brief, cells were washed twice with ice-cold phosphate
50
51 buffered saline (PBS) and then incubated in the lysis buffer (50 mM Tris-HCl pH 8, 150
52
53 mM NaCl, 1% NP-40) on ice for 30 min. After centrifugation at 4 °C for 15 min at 14,000g,
54
55 the supernatant (30 µg per lane) was collected and resolved on a 10% SDS-PAGE. Proteins
56
57
58
59
60

1
2
3
4 on the gel were then transferred onto a polyvinylidene fluoride membrane (0.4- μm pore
5 size; Millipore). Non-specific protein binding onto the membrane was blocked with
6 Tris-buffered saline containing 0.1% Tween 20 (TBST) and 5% non-fat milk at 4°C
7 overnight [Lee et al., 2006]. After washing three times with TBST, the membrane was
8 incubated with the primary antibodies (1:500 in TBST containing 5% non-fat milk) at room
9 temperature for 1 h. After washing with TBST, bound antibodies were detected by
10 horseradish peroxidase (HRP)-conjugated secondary antibodies (1:20,000 dilution) and
11 ECL reagent (Amersham Biosciences). The immunoreactive signals were developed on
12 Kodak® Biomax™ MS film.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

31 **Quantitative PCR**

32
33 Total RNA was isolated from cell cultures or frozen tissues using Trizol reagent
34 (Invitrogen). Two micrograms of total RNA were used to synthesize first-strand cDNA
35 using random hexamer primers and reverse transcriptase (Promega). A 5 μl aliquot of 1:10
36 diluted cDNA was mixed with 10 μl SYBR® GreenERTM qPCR superMix (Invitrogen)
37 and 0.5 μM of the designated primers (Table 1). The thermal cycling conditions were 50 °C
38 for 2 minutes, 95 °C for 10 minutes, followed by 40 cycles at 95 °C for 15 seconds and
39 60 °C for 1 minute. All reactions were done at least twice and the products were analyzed
40 using the ABI Prism 7700 detection system (Applied Biosystems Inc.). Ribosomal 18S
41 RNA was included in all reactions for internal house-keeping control and normalization.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Electromobility shift assay (EMSA)

1
2
3
4 Nuclear protein extracts from MHCC97L and Hep3B cells were prepared as described
5
6 previously [Luk et al., 2004], and the amount of protein extract was quantified using the
7
8 Bradford protein assay kit (Bio-Rad). Double-strand oligonucleotide probes were
9
10 synthesized with annealing buffer (200 mM Tris-HCl;pH 9.0, 40mM MgCl₂, 1 M NaCl, 20
11
12 mM EDTA) and labeled with [γ -³²P] ATP (PerkinElmer, Waltham, MA) using the
13
14 Ready-To-Go T4 Polynucleotide Kinase kit (Amersham BioSciences) with T4
15
16 polynucleotide kinase. Unincorporated nucleotides were removed by spin column in a
17
18 Tris-EDTA buffer, pH 8.0. An aliquot (10 μ g) of nuclear extract was incubated with or
19
20 without 1 μ l unlabeled oligonucleotide probes or the previously described antibodies in 1x
21
22 binding buffer (4% glycerol, 1 μ g poly(dI-dC).(dI-dC), 10 mM Tris-HCl (pH 7.5), 50 mM
23
24 NaCl, 1 mM MgCl₂, 0.5 mM EDTA and 0.5 mM DTT) in a 10 μ l reaction volume for 20
25
26 min at room temperature. Radiolabeled probe was added to each reaction tube and the tubes
27
28 were incubated for a further 20 min at room temperature. The reaction products were then
29
30 resolved on a 5% non-denaturing polyacrylamide gel. The gels were dried under vacuum
31
32 and autoradiographed on Kodak® Biomax™ MS films overnight at -70°C.
33
34
35
36
37
38
39
40
41
42
43
44
45
46

47 **Chromatin Immunoprecipitation (ChIP) assay**

48
49 Standard ChIP assays were performed with Millipore ChIP Assay kit according to the
50
51 manufacturer's instructions. Briefly, approximately 1 x 10⁶ cells were cross-linked using
52
53 1% formaldehyde at 37°C for 10 min, washed twice with PBS, and then resuspended in
54
55 200 μ l SDS lysis buffer containing 1x protease inhibitor cocktail (Roche Applied Science,
56
57 Indianapolis, IN) and 10 mM of DTT. After incubation on ice for 10 min, the cells were
58
59
60

1
2
3
4 sonicated (Sonic Dismembrator Model 100, Fisher Scientific, Waltham, MA) four times at
5
6
7 20% maximum power for 10 s and the lysate was centrifuged at 4°C for 15 min at 14,000g.
8
9
10 The resultant supernatant was diluted 1:10 with a dilution buffer containing a protease
11
12 inhibitor. Twenty microliter of the diluted supernatant was collected as input control before
13
14 immunoprecipitation. After precleared with 75 μ L of salmon sperm DNA/protein A agarose
15
16 (50% slurry) for 1 hour at 4°C with agitation, the immune complexes were incubated at
17
18 4°C overnight with either 2 μ g of primary antibodies or IgG control, and then recovered
19
20 with 60 μ l of salmon sperm DNA/protein A agarose 50% slurry at 4°C for 1 hour.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Precipitates were washed once sequentially with washing solutions provided by the manufacturer (Millipore), and finally with two washes of TE buffer. Chromatin complexes were incubated at room temperature for 15 min with 250 μ l of elution buffer containing 1% SDS and 100mM sodium bicarbonate. The eluate was reverse cross-linked by heating at 65°C for 4 hours with 200 mM NaCl and 200 mg/ml proteinase K. The DNA was recovered by phenol/chloroform extraction and ethanol precipitation and then amplified by PCR with forward primer 5'-GTGCTAAGTGTTGGGGGTACAA-3' and reverse primer 5'-TTGACTGAAGCTGAAGGGAGAG-3'. PCR was carried out at 94 °C for 5 min and then 30 cycles at 94 °C for 40 s, 60 °C for 1 min, and 72 °C for 1 min and a final extension step at 72 °C for 10 min in a final volume of 25 μ l.

HNF1 α and CDX2 siRNA knockdown

Small interference RNA (siRNA) was used to knockdown the expression of HNF1 α and CDX2 in HCC cells (MHCC97L and Hep3B). RNA oligonucleotides were designed

1
2
3
4 according to the published sequences [Hinoi et al., 2005]. HCC cells were transfected with
5
6 the designated siRNA duplex or scramble control using Lipofectamine™ 2000 (Invitrogen)
7
8 as described above. After 48-hour incubation, both mRNA and protein extracts were
9
10 purified for qPCR and Western blot analyses, respectively.
11
12
13
14
15
16

17 **Statistical analysis**

18
19 All results are represented as mean \pm S.D. Statistical comparisons were performed by
20
21 One-way ANOVA or paired Student's t-test using the SPSS statistical software package
22
23 version 13.0 for Window (SPSS Inc., Chicago, IL) and $p < 0.05$ was considered to be
24
25 statistically significant.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RESULTS

Cloning and characterization of CDH17 gene promoter

CDH17 promoter region was amplified by PCR and cloned into pGL3-Basic reporter vector. The position and sequence of the primers used to generate the parental fragment and the four deletion constructs were shown in Figure 1A. Two bioinformatics programs, ALGGEN-PROMO (<http://algggen.lsi.upc.es>) and MatInspector (<http://www.genomatix.de>) were used to identify potential transcription factors at the -909/+55 promoter region and the results were compared to the transcription factor database TRANSFAC (version 8.3). A conserved CAAT box, but not the TATA-box, was found in the CDH17 gene promoter region. Interestingly, two putative transcription factors, HNF1 and CDX2 were identified. At this stage, the binding of HNF1 α from HNF1 β cannot be differentiated as they shared similar binding sequences. There are three putative binding sites for each HNF1 and CDX2 transcription factors in the CDH17 promoter, respectively. Figure 1B shows the luciferase reporter activities of different promoter deletion constructs when compared to the parental (-909/+55) control in two different HCC cell lines: MHCC97L (CDH17^{high}) and Hep3B (CDH17^{low}). The data indicated that CDX2 and HNF1 α are two critical factors for the transcription of CDH17 because the luciferase activities decreased significantly from the pGL-81/+55Luc to pGL3-40/+55Luc. Both cell lines exhibited similar activity patterns although the signals were generally stronger in the MHCC97L cell line.

To confirm the functional roles of these transcriptional factors (TF), point mutations were introduced at the TF core binding sites in the CDH17 promoter sequence.

1
2
3
4 Reporter activities were markedly reduced for HNF1 mutant (-47/-45, $p < 0.01$), followed by
5
6 the CDX2 mutant (-91/-89, $p < 0.05$) (Fig. 2). Mutations on other HNF1 and CDX2 binding
7
8 sites did not drastically reduce promoter activities. Double mutations on the HNF1 (-47/-45)
9
10 and CDX2 (-91/-89) binding sites reduced the CDH17 reporter activity to near basal level.
11
12
13
14
15
16

17 **Binding of CDX2 and HNF1 α to CDH17 promoter *in vitro* and *in vivo***

18
19
20 The binding of HNF1 and CDX2 to the CDH17 gene promoter in MHCC97L and Hep3B
21
22 cells were studied by EMSA and ChIP assays. As shown in Fig. 3A, both the HNF1 α and
23
24 CDX2 wild-type (WT), but not the mutant probes were able to complexes with the HCC
25
26 lysates. Further, addition of 10 and 50-fold excess of “cold” WT probes, but not the mutant
27
28 probes, could displace or compete with the binding. Moreover, CDX2, HNF1 α , but not
29
30 HNF1 β antibodies were able to retard the migration of the DNA-protein complex,
31
32 suggesting the binding of the nuclear extract is specific. Meanwhile, the ChIP assay was
33
34 performed to study TF-DNA interactions *in vivo*. After immunoprecipitation by designated
35
36 antibodies, the recovered DNA was amplified by PCR using CDH17 promoter-specific
37
38 primers spanning from -318 to -18 regions. Positive signals were detected by the HNF1 α
39
40 and CDX2, but not the HNF1 β and isotype control (Fig. 3B).
41
42
43
44
45
46
47
48
49
50
51

52 **Differential expression of CDH17, HNF1 and CDX in HCC and fetal mouse liver**

53
54
55 Luciferase reporter assays revealed differential promoter activities in Hep3B and
56
57 MHCC97L cell lines (Fig. 1B and Fig. 2). To study if the endogenous CDH17 transcript
58
59 level correlates with the HNF1 α and CDX2 expression levels, we determined the
60

1
2
3
4 expression of these transcripts in HCC cells and mouse livers at different developmental
5
6 stages. MHCC97L which has a high metastatic potential also expressed a high level of
7
8 CDH17 transcript as well as the CDX2 and HNF1 α transcripts; whereas the primary HCC
9
10 (Hep3B) and MIHA (immortalized normal human hepatocyte) expressed at a lower or
11
12 undetectable levels of the gene (Fig. 4A).
13
14
15
16

17
18 Since CDH17 plays important roles during embryonic liver development, the
19
20 expression of CDH17 and TFs levels in mouse liver at different development stages (E13.5,
21
22 E16.5, newborn, 3 weeks and adult) were measured. Bioinformatics analysis of these
23
24 transcriptional factor binding sites on CDH17 promoter between mouse, rat and human are
25
26 highly conserved using ClustalW2 program ([http://www.ebi.ac.uk/](http://www.ebi.ac.uk/Tools/clustalw2/index.html)
27
28 Tools/clustalw2/index.html), suggesting a conserved regulatory mechanism of CDH17
29
30 expression among different animal species (data not shown). It was found that CDH17
31
32 expressed abundantly in the early embryonic liver, but the expression of the gene becomes
33
34 undetectable in late fetal stages and adult livers (Fig. 4A). Similar to the human HCC
35
36 findings, the differential expression of CDH17 transcript in mouse liver largely correlated
37
38 with the patterns of HNF1 α and CDX2 levels. The result was supported by Western blot
39
40 analysis of the CDH17, HNF1 α and CDX2 protein levels (Fig. 4B). Interestingly, the
41
42 decreases of CDH17 and HNF1 α were more drastic in both human HCC cells and mouse
43
44 livers, than the CDX2.
45
46
47
48
49
50
51
52
53
54
55
56

57 **Knockdown of HNF1 α and CDX2 suppress CDH17 expression**

58
59 To demonstrate HNF1 α and CDX2 regulate CDH17 expression, we used siRNA approach
60

1
2
3
4 to knockdown HNF1 α and CDX2 expression and study the expression of CDH17 in
5
6 MHCC97L cells (Fig. 5). RNAi-mediated targeting HNF1 α or CDX2 lowered the mRNA
7
8 level of the respective transcriptional factor (HNF1 α or CDX2) by approximate 50%, when
9
10 compared with the vehicle or scramble control. These down-regulations were accompanied
11
12 with significant reduction of CDH17 transcript level in MHCC97L cells ($p < 0.05$). An
13
14 additive effect of combined HNF1 α or CDX2 RNAi treatment on CDH17 suppression was
15
16 also observed, indicating a coordinate regulation of HNF1 α and CDX2 in the cells.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

DISCUSSION

Liver-intestine cadherin or cadherin-17 (CDH17) is expressed in the rat liver and intestinal epithelial cells [Angres et al., 2001], as well as in the embryonic mouse liver [Takamura et al., 2004], but no CDH17 expression was detected in normal liver tissue of mature humans and mice [Takamura et al., 2003]. CDH17 is involved in the maintenance of tissue integrity, tissue growth and differentiation [Berndorff et al., 1994]. Aberrant expression of CDH17 mRNA and protein was found in upper gastrointestinal malignancies including liver cancer [Wang et al., 2006; Wang et al., 2005; Wong et al., 2003]. In this study, we demonstrated that the expression of CDH17 is regulated by HNF1 α and CDX2 transcriptional factors in HCC and mouse liver.

The human CDH17 gene promoter has a conserved CAAT box at the -65/-62 position, but lacks the TATA box, an initiator element in the predominant control region of typical eukaryotic genes [Suzuki et al., 2001]. In both primary and metastatic human HCC cell lines, the minimal CDH17 promoter (-81/-40) region contains one CDX2 and two HNF1 sites that mediated strong promoter activity in vitro. Further deletion of this minimal promoter region diminished the reporter activity to near basal levels. We also found that the transactivation activity of the pGL3-81/+55Luc construct was similar to longer constructs (up to -909 position). Although the deletion construct pGL3-332/+55Luc shown a decrease in promoter activity, the reason behind is still unknown. It is possible that an enhancer sequence is located at -686 and -332 region, and a repressor sequence is located at -32 to -81 region, thereby modulate the changes in promoter activity. As a result, the deletion construct pGL3-81/+55Luc shown a full-promoter activity. Yet, further

1
2
3
4 experiments are needed to demonstrate if these regulatory sequences play any significant
5
6
7 role in vivo.

8
9
10 Mutation analysis identified a HNF1 site at -47/-45 position is crucial for CDH17
11
12 expression. To a lesser extent, CDX2 (-91/-89) regulates the CDH17 promoter activity in
13
14 vitro. Furthermore, both EMSA and ChIP assays demonstrated binding of HNF1 and
15
16 CDX2 to the CDH17 promoter sequences. HNF1 α , but not the HNF1 β isoform bound to
17
18 the CDH17 promoter and the protein-DNA complex was recognized by HNF1 α specific
19
20 antibody. The temporal expression of HNF1 α and CDX2 transcript in related to CDH17
21
22 expression was studied in human liver cell lines and mouse liver tissues. It was found that
23
24 an elevated levels of HNF1 α and CDX2 expression are closely associated with a higher
25
26 expression of CDH17 in metastatic HCC cell line (MHCC97L >> Hep3B > MIHA) as well
27
28 as in early embryonic mouse liver tissues [E13.5 > E16.5 > new born (NB)]. The role of
29
30 HNF1 α , and CDX2 on co-regulating the CDH17 gene expression was further demonstrated
31
32 by iRNA suppression on CDX2 and HNF1 α expression in HCC cells.
33
34
35
36
37
38
39
40

41
42 HNF1 and CDX2 are homeodomain transcription factors that regulate a number
43
44 of intestine-specific genes including Lewis type 1 antigen synthase (β 3Gal-T5) and
45
46 UDP-glucuronosyltransferases (UGT) [Gregory et al., 2004; Isshiki et al., 2003]. CDX2
47
48 regulates CDH17 expression in colorectal cancer cell lines and gastric adenocarcinoma
49
50 [Barros et al., 2008; Hinoi et al., 2002; Ko et al., 2005]. In this study, 3 putative
51
52 CDX2-binding sites were found in the 1-kb CDH17 promoter region (Fig. 1A) and it is the
53
54 -91/-89 site that contributed to CDH17 promoter activity and binding in HCC. In addition,
55
56 we identified HNF1 homeobox A (HNF1 α) that may play an important role in regulating
57
58
59
60

1
2
3
4 CDH17 transcription in human HCC as well as in mouse fetal liver. Although there are
5
6 three putative HNF1-binding sites in the CDH17 promoter, deletion/mutation studies
7
8 identified -47/-45 position is crucial in regulating CDH17 expression (Fig. 1B and Fig. 2).
9
10

11 HNF1 α and HNF1 β are two related transcriptional factors that bind to DNA as
12
13 homo- or heterodimers. They were identified initially as liver-enriched transcription factors
14
15 that were involved in the expression of several plasma proteins including α 1-antitrypsin
16
17 [Mendel and Crabtree, 1991]. However, no detectable HNF1 β transcript and protein was
18
19 found in the two HCC cell lines studied (data not shown), and HNF1 β failed to bind to
20
21 CDH17 promoter DNA sequence in EMSA or ChIP assays. These findings suggest that
22
23 HNF1 α , but not HNF1 β is associated with the regulation of the CDH17 gene in HCC.
24
25 Interestingly, HNF1 β has recently been demonstrated to direct the expression of
26
27 kidney-specific cadherin (cadherin-16) [Bai et al., 2002].
28
29
30
31
32
33
34
35

36 In conclusion, the present study provides the first evidence that HNF1 α plays a
37
38 critical role, together with CDX2, on the regulation of CDH17 gene expression in HCC as
39
40 well as in early stages of fetal liver development in mice. Clinically, over-expression of
41
42 CDH17 in HCC is associated with poor clinical outcomes; HNF1 α and CDX2 in
43
44 accompany with CDH17 may be potential therapeutic targets for disease diagnosis and
45
46 treatment.
47
48
49
50
51
52
53
54

55 **ACKNOWLEDGMENTS**

56
57 The work was supported by a grant from the General Research Fund (HKU771607M) of
58
59 the Research Grants Council of Hong Kong. We thank Dr. Jana Wo for the technical
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

assistance on microscopic surgery of fetal mouse liver and Dr. Arieh Bomzon for critical review and professional editing of the manuscript.

For Peer Review

REFERENCES

Angres B, Kim L, Jung R, Gessner R, Tauber R. 2001. LI-cadherin gene expression during mouse intestinal development. *Dev Dyn* 221:182-193.

Bai Y, Pontoglio M, Hiesberger T, Sinclair AM, Igarashi P. 2002. Regulation of kidney-specific Ksp-cadherin gene promoter by hepatocyte nuclear factor-1beta. *Am J Physiol Renal Physiol* 283:F839-51.

Barros R, Pereira B, Duluc I, Azevedo M, Mendes N, Camilo V, Jacobs RJ, Paulo P, Santos-Silva F, van Seuningen I, van den Brink GR, David L, Freund JN, Almeida R. 2008. Key elements of the BMP/SMAD pathway co-localize with CDX2 in intestinal metaplasia and regulate CDX2 expression in human gastric cell lines. *J Pathol* 215:411-420.

Berndorff D, Gessner R, Kreft B, Schnoy N, Lajous-Petter AM, Loch N, Reutter W, Hortsch M, Tauber R. 1994. Liver-intestine cadherin: molecular cloning and characterization of a novel Ca(2+)-dependent cell adhesion molecule expressed in liver and intestine. *J Cell Biol* 125:1353-69.

Courtois G, Morgan JG, Campbell LA, Fourel G, Crabtree GR. 1987. Interaction of a liver-specific nuclear factor with the fibrinogen and alpha-1-antitrypsin promoters. *Science* 238:688-692.

Dantzig AH, Hoskins JA, Tabas LB, Bright S, Shepard RL, Jenkins IL, Duckworth DC, Sportsman JR, Mackensen D, Rosteck PR, Jr., et al. 1994. Association of intestinal peptide transport with a protein related to the cadherin superfamily. *Science* 264:430-3.

Gregory PA, Lewinsky RH, Gardner-Stephen DA, Mackenzie PI. 2004. Coordinate regulation of the human UDP-glucuronosyltransferase 1A8, 1A9, and 1A10 genes by hepatocyte nuclear factor 1alpha and the caudal-related homeodomain protein 2. *Mol Pharmacol* 65:953-63.

Grotzinger C, Kneifel J, Patschan D, Schnoy N, Anagnostopoulos I, Faiss S, Tauber R, Wiedenmann B, Gessner R. 2001. LI-cadherin: a marker of gastric metaplasia and neoplasia. *Gut* 49:73-81.

Hinoi T, Lucas PC, Kuick R, Hanash S, Cho KR, Fearon ER. 2002. CDX2 regulates liver intestine-cadherin expression in normal and malignant colon epithelium and intestinal metaplasia. *Gastroenterology* 123:1565-1577.

Isshiki S, Kudo T, Nishihara S, Ikehara Y, Togayachi A, Furuya A, Shitara K, Kubota T, Watanabe M, Kitajima M, Narimatsu H. 2003. Lewis type 1 antigen synthase (beta3Gal-T5) is transcriptionally regulated by homeoproteins. *J Biol Chem* 278:36611-36620.

Johnson PJ. 2001. The role of serum alpha-fetoprotein estimation in the diagnosis and management of hepatocellular carcinoma. *Clin Liver Dis* 5:145-159.

Jung R, Wendeler MW, Danevad M, Himmelbauer H, Gessner R. 2004. Phylogenetic origin of LI-cadherin revealed by protein and gene structure analysis. *Cell Mol Life Sci*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

61:1157-1166.

Kaposi-Novak P, Lee JS, Gomez-Quiroz L, Coulouarn C, Factor VM, Thorgeirsson SS. 2006. Met-regulated expression signature defines a subset of human hepatocellular carcinomas with poor prognosis and aggressive phenotype. *J Clin Invest* 116:1582-1595.

Ko S, Chu KM, Luk JM, Wong BW, Yuen ST, Leung SY, Wong J. 2004. Overexpression of LI-cadherin in gastric cancer is associated with lymph node metastasis. *Biochem Biophys Res Commun* 319:562-8.

Ko S, Chu KM, Luk JM, Wong BW, Yuen ST, Leung SY, Wong J. 2005. CDX2 co-localizes with liver-intestine cadherin in intestinal metaplasia and adenocarcinoma of the stomach. *J Pathol* 205:615-622.

Lee KF, Xu JS, Lee YL, Yeung WSB. 2006. Demilune cell and parotid protein (Dcpp) from murine oviductal epithelium stimulates pre-implantation embryo development. *Endocrinology* 147:79-87.

Lee NP, Chen L, Lin MC, Tsang FH, Yeung C, Poon RT, Peng J, Leng X, Beretta L, Sun S, Day PJ, Luk JM. 2009. Proteomic expression signature distinguishes cancerous and nonmalignant tissues in hepatocellular carcinoma. *J Proteome Res* 8:1293-303.

Liu CL, Fan ST. 2006. Adult-to-adult live-donor liver transplantation: the current status. *J Hepatobiliary Pancreat Surg* 13:110-6.

Llovet JM, Burroughs A, Bruix J. 2003. Hepatocellular carcinoma. *Lancet* 362:1907-17.

Luk JM, Tong MK, Mok BW, Tam PC, Yeung WSB, Lee KF. 2004. Sp1 site is crucial for the mouse claudin-19 gene expression in the kidney cells. *FEBS Letters* 578:251-256.

Mendel DB, Crabtree GR. 1991. HNF-1, a member of a novel class of dimerizing homeodomain proteins. *J Biol Chem* 266:677-80.

Nakabayashi H, Koyama Y, Suzuki H, Li HM, Sakai M, Miura Y, Wong NC, Nishi S. 2004. Functional mapping of tissue-specific elements of the human alpha-fetoprotein gene enhancer. *Biochem Biophys Res Commun*. 318:773-85.

Poon RT, Fan ST. 2004. Hepatectomy for hepatocellular carcinoma: patient selection and postoperative outcome. *Liver Transpl* 10:S39-45.

Suzuki Y, Tsunoda T, Sese J, Taira H, Mizushima-Sugano J, Hata H, Ota T, Isogai T, Tanaka T, Nakamura Y, Suyama A, Sakaki Y, Morishita S, Okubo K, Sugano S. 2001. Identification and characterization of the potential promoter regions of 1031 kinds of human genes. *Genome Res* 11:677-84.

Takamura M, Ichida T, Matsuda Y, Kobayashi M, Yamagiwa S, Genda T, Shioji K, Hashimoto S, Nomoto M, Hatakeyama K, Ajioka Y, Sakamoto M, Hirohashi S, Aoyagi Y. 2004. Reduced expression of liver-intestine cadherin is associated with progression and lymph node metastasis of human colorectal carcinoma. *Cancer Lett* 212:253-9.

Takamura M, Sakamoto M, Ino Y, Shimamura T, Ichida T, Asakura H, Hirohashi S. 2003.

1
2
3
4 Expression of liver-intestine cadherin and its possible interaction with galectin-3 in ductal
5 adenocarcinoma of the pancreas. *Cancer Sci* 94:425-30.

6
7 Wang W, Hayashi Y, Ninomiya T, Ohta K, Nakabayashi H, Tamaoki T, Itoh H. 1998.
8 Expression of HNF-1 alpha and HNF-1 beta in various histological differentiations of
9 hepatocellular carcinoma. *J Pathol.* 184:272-278.

10
11 Wang XQ, Luk JM, Garcia-Barcelo M, Miao X, Leung PP, Ho DW, Cheung ST, Lam BY,
12 Cheung CK, Wong AS, Lau SS, So MT, Yu WC, Cai Q, Liu KS, Hui CK, Lau GK, Poon RT,
13 Wong J, Fan ST. 2006. Liver intestine-cadherin (CDH17) haplotype is associated with
14 increased risk of hepatocellular carcinoma. *Clin Cancer Res* 12:5248-52.

15
16 Wang XQ, Luk JM, Leung PP, Wong BW, Stanbridge EJ, Fan ST. 2005. Alternative mRNA
17 splicing of liver intestine-cadherin in hepatocellular carcinoma. *Clin Cancer Res*
18 11:483-489.

19
20 Wong BW, Luk JM, Ng IO, Hu MY, Liu KD, Fan ST. 2003. Identification of liver-intestine
21 cadherin in hepatocellular carcinoma--a potential disease marker. *Biochem Biophys Res*
22 *Commun* 311:618-624.

23
24 Xu L, Hui L, Wang S, Gong J, Jin Y, Wang Y, Ji Y, Wu X, Han Z, Hu G. 2001. Expression
25 profiling suggested a regulatory role of liver-enriched transcription factors in human
26 hepatocellular carcinoma. *Cancer Res* 61:3176-3181.

FIGURE LEGENDS

Figure 1 Cloning and characterization of CDH17 gene promoter. (A) The nucleotide sequences of the 5'-flanking regions of human CDH17 gene and the putative transcriptional regulatory elements from -955 were depicted. The G at position +1 indicates the transcriptional initiation site based on the GenBank database. The translation start site (ATG) is underlined and the amino acid sequences are indicated by bold and capital letters above nucleotide sequences. The primer sequences used for making the parental fragment and the four deletion constructs were each marked with arrows (forward and reverse primers). (B) The promoter activity of CDH17 deletion constructs in Hep3B and MHCC97L cells was studied. The pGL-332/+55Luc and pGL-40/+55Luc constructs exhibited drastic reduction in luciferase activity ($p < 0.05$ and $p < 0.01$, respectively) when compared with the control fragment (pGL-909/+55Luc). The luciferase activity of the empty expression vector pGL3-Basic was assigned a value of 1 and the relative activities of each construct were normalized with the pRL-TK expression levels. Values are represented as mean \pm S.D. from at least three independent experiments.

Figure 2 Mutation of Transcriptional factor binding sites on CDH17 promoter activity. The relative luciferase activities of mutated HNF1 and CDX2 sites in CDH17 promoter constructs () were studied in Hep3B and MHCC97L cells. The mutated HNF1 (filled oval) and/or CDX2 (filled rectangle) binding sites on CDH17 promoters were transfected into the HCC cells. Mutation at the CDX2 (-91 position) and/or HNF1 (-47 position) binding site(s) significantly reduced ($p < 0.05$ and $p < 0.01$, respectively) the

1
2
3
4 luciferase activity. Combined mutation of these two sites (-91 and -47) also resulted in a
5
6
7 significant reduction in luciferase activity ($p < 0.01$). Yet, no significant different in
8
9
10 luciferase activity was found between single (HNF1) and double (HNF1 plus CDX2) sites
11
12 mutation. Values are expressed as mean \pm S.D. of at least three independent experiments.
13
14
15
16

17 **Figure 3 Binding of HNF1 and CDX2 on CDH17 promoter sequence in vitro and**
18 **in vivo. (A)** Radiolabeled CDX2 and HNF1 probes (WT and mutant) were incubated with
19
20 Hep3B and MHCC97L nuclear extracts and then resolved on a 5% polyacrylamide gel. The
21
22 presence of CDH17-TF (either CDX2 or HNF1 α) complexes suggests binding of TF and
23
24 DNA in vitro. Unbound probes migrated to the bottom of gels. Addition of excessive (10X
25
26 and 50X) specific WT (wt), but not mutant (mut) cold probes reduced the signal
27
28 specifically. While the presence of specific antibodies binds to the TF-DNA complexes
29
30 further retarded the migration of the complex in the gel (supershift). The sequences of the
31
32 probes used were shown at the bottom. **(B)** ChIP assay was used to detect the presence of
33
34 DNA-TF(CDX2 and HNF1 α) complexes in MHCC97L and Hep3B cells in vivo.
35
36 Antibodies against the CDX2, HNF1 α , HNF1 β , and IgG control, were used to
37
38 immunoprecipitate the TF-DNA complexes. After recovery, the purified DNA was
39
40 amplified by CDH17-specific PCR reaction. Input controls were genomic DNA extracted
41
42 from MHCC97L and Hep3B cells without cross-linking and immunoprecipitation steps.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57


58 **Figure 4 Expression of CDH17, HNF1 α and CDX transcripts in human HCC cells**
59 **and mouse liver. (A)** Quantitative PCR was used to quantify the expression of CDH17,
60

1
2
3
4 HNF1 α and CDX2 transcripts. The expression of CDH17, HNF1 α and CDX2 transcripts
5
6 were highest in human MHCC97L and lower in Hep3B and MIHA cells; while gradually
7
8 decreases from E13.5, E16.5, new born (NB), 3 weeks old (3 wk) adult mouse liver. (B)
9
10 Western blot analysis confirmed a high expression level of CDH17, HNF1 α and CDX2
11
12 proteins in MHCC97L cells. A lower expression level was found in Hep3B and MIHA
13
14 cells. The protein expression levels of CDH17 and HNF1 α decrease drastically in mouse
15
16 fetal liver from E135 until born. Yet, the expression of CDX2 protein remains constant
17
18 through development. The expression of β -actin protein was used as loading control.
19
20
21
22
23
24
25
26
27

28 **Figure 5 siRNA-mediated knockdown of HNF1 α and CDX2 in HCC cells. (A)**


29
30 Targeting siRNA oligonucleotides against the HNF1 α and CDX2 genes were transfected
31
32 into the CDH17 expressing MHCC97L cells. Quantity of the CDH17, HNF1 α and CDX2
33
34 transcripts was measured by real-time qPCR. Vector and scramble siRNA were included as
35
36 controls in each experiments. Data were shown as mean \pm S.D. from three independent
37
38 experiments. * denotes statistically significantly different at $p < 0.05$. (B) Western blotting
39
40 demonstrated that CDH17 protein expression was down-regulated after siRNA against
41
42 HNF1 α and CDX2.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



253x190mm (600 x 600 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




253x190mm (600 x 600 DPI)


Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

A




B



253x190mm (600 x 600 DPI)

Review



253x190mm (600 x 600 DPI)

Table 1 Primers used for real-time PCR for detection of transcripts in human liver cells and mouse liver tissues

Primer Name	Sequence	Product Size (bp)
Human		
CDH17 Forward	5'-GGGGGAGATACTCCAGTCGT-3'	167
CDH17 Reverse	5'-TCCAGTTGCCAAATAAAGCA-3'	
CDX2 Forward	5'-GAACCTGTGCGAGTGGATG-3'	158
CDX2 Reverse	5'-TCCTCCGGATGGTGATGTAG-3'	
HNF1 α Forward	5'-CCATCCTCAAAGAGCTGGAG-3'	173
HNF1 α Reverse	5'-CTGGTTGAGGCCAGTGGTAT-3'	
Mouse		
CDH17 Forward	5'-GGCCAAGAACCGAGTCAAGTC-3'	190
CDH17 Reverse	5'-CTCCATGAGAATCCAAGGCTG-3'	
CDX2 Forward	5'-CAAGGACGTGAGCATGTATCC-3'	106
CDX2 Reverse	5'-GTAACCACCGTAGTCCGGGTA-3'	
HNF1 α Forward	5'-GACCTGACCGAGTTGCCTAAT-3'	103
HNF1 α Reverse	5'-CCGGCTCTTTCAGAATGGGT-3'	