

BMJ Open Semi-individualised Chinese medicine treatment as an adjuvant management for diabetic nephropathy: a pilot add-on, randomised, controlled, multicentre, open-label pragmatic clinical trial

Kam Wa Chan,¹ Tai Pang Ip,² Alfred Siu Kei Kwong,³ Sing Leung Lui,² Gary Chi Wang Chan,⁴ Benjamin John Cowling,⁵ Wai Han Yiu,¹ Dickson Wai Leong Wong,¹ Yang Liu,¹ Yibin Feng,⁶ Kathryn Choon Beng Tan,⁷ Loretta Yuk Yee Chan,¹ Joseph Chi Kam Leung,¹ Kar Neng Lai,¹ Sydney Chi Wai Tang¹

To cite: Chan KW, Ip TP, Kwong ASK, *et al*. Semi-individualised Chinese medicine treatment as an adjuvant management for diabetic nephropathy: a pilot add-on, randomised, controlled, multicentre, open-label pragmatic clinical trial. *BMJ Open* 2016;**6**:e010741. doi:10.1136/bmjopen-2015-010741

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2015-010741>).

Received 3 December 2015
Revised 8 July 2016
Accepted 14 July 2016



CrossMark

For numbered affiliations see end of article.

Correspondence to
Professor Sydney Chi Wai Tang; scwtang@hku.hk

ABSTRACT

Introduction: Diabetes mellitus and diabetic nephropathy (DN) are prevalent and costly to manage. DN is the leading cause of end-stage kidney disease. Conventional therapy blocking the renin–angiotensin system has only achieved limited effect in preserving renal function. Recent observational data show that the use of Chinese medicine (CM), a major form of traditional medicine used extensively in Asia, could reduce the risk of end-stage kidney disease. However, existing clinical practice guidelines are weakly evidence-based and the effect of CM remains unclear. This trial explores the effect of an existing integrative Chinese–Western medicine protocol for the management of DN.

Objective: To optimise parameters and assess the feasibility for a subsequent phase III randomised controlled trial through preliminary evaluation on the effect of an adjuvant semi-individualised CM treatment protocol on patients with type 2 diabetes with stages 2–3 chronic kidney disease and macroalbuminuria.

Methods and analysis: This is an assessor-blind, add-on, randomised, controlled, parallel, multicentre, open-label pilot pragmatic clinical trial. 148 patients diagnosed with DN will be recruited and randomised 1:1 to a 48-week additional semi-individualised CM treatment programme or standard medical care. Primary end points are the changes in estimated glomerular filtration rate and spot urine albumin-to-creatinine ratio between baseline and treatment end point. Secondary end points include fasting blood glucose, glycated haemoglobin, brain natriuretic peptide, fasting insulin, C peptide, fibroblast growth factor 23, urinary monocyte chemoattractant protein-1, cystatin C, nephrin, transforming growth factor- β 1 and vascular endothelial growth factor. Adverse events are monitored through self-completed

Strengths and limitations of this study

- This study is the first to assess the potential efficacy of an adjuvant Chinese medicine programme in diabetic kidney disease.
- Chinese medicine is being extensively used in Asia and getting popular worldwide. Previous epidemiological data suggest that Chinese medicine may retard the deterioration of renal function among patients with diabetic kidney disease. Hence, it is timely to perform a clinical trial.
- This study is a pilot study and the sample size is not powered to achieve statistical significance.
- Since the trial is open-label, subjective outcomes including quality of life could not be assessed.

questionnaire and clinical visits. Outcomes will be analysed by regression models. Enrolment started in July 2015.

Ethics and registration: This protocol is approved by the Institutional Review Board of the University of Hong Kong/Hospital Authority Hong Kong West Cluster (reference number UW 14-301).

Trial registration number: NCT02488252.

INTRODUCTION

The epidemic of diabetes mellitus (DM) is predicted to rise steadily in incidence and prevalence all over the world, driven by the effects of an ageing population and rising levels of obesity.^{1 2} It is estimated that 387 million people were having DM in 2014,³ and by 2030, the DM population is projected to reach 439 million.⁴ Diabetic nephropathy

(DN) is the leading cause of end-stage kidney disease (ESKD)⁵ and is developed among one-third of all patients with diabetes.^{5 6} In Hong Kong, the incidence of ESKD primarily caused by DN requiring renal replacement therapy in the form of chronic dialysis or kidney transplantation doubled from 1996 (26.2%) to 2013 (49.6%).⁷ Patients with DN have the lowest adjusted 5-year survival probability of 50.6% when compared with patients with incident renal replacement therapy with other causes including hypertension (61.1%) and glomerulonephritis (71.1%).⁸ DN imposes a major burden on patients and healthcare systems in many developed regions.⁵

The pathogenesis of DN is diverse^{9–13} and yet to be fully understood. It involves glycototoxicity, advanced glycation end products and reactive oxygen species which induce haemodynamic changes and inflammation. Biopsy from patients with DN shows that podocyte loss, basement membrane thickening, interstitial fibrosis and tubular atrophy and DN are characterised by proteinuria clinically. Conventional therapy from the western literature pinpointing a few selected pathways, such as the widely advocated blockade of the renin–angiotensin system, has only achieved modest efficacy in treating DN.⁶ Combined therapy with several conventional regimes, for instance, ACE inhibitor and angiotensin receptor blocker may not be as efficacious as believed.^{6 14} No specific therapies for DN or renal fibrosis are available currently. Chinese medicine (CM) is a popular complementary remedy in addition to standard medical care among the Chinese in Hong Kong.^{15 16} A recent observational study in Taiwan shows that the use of CM is associated with an average risk reduction of 59% of ESKD when compared with non-users in a 6-year period. However, the risk reduction varies significantly across different subgroups of patients stratified clinically based on CM theory.¹⁷ Despite its popularity, there is no evidence-based clinical guideline for CM as an adjuvant therapy to standard medical care in managing DN.

An integrative Chinese–Western medicine clinical management protocol for DN was developed by reviews of existing clinical management guidelines and an expert consensus from CM nephrologists in mainland China in 2012.¹⁸ The adjuvant CM treatment consists of five different CM formulas corresponding to five subgroups of patients stratified by clinical manifestations. A preliminary evaluation on renal clinical outcomes among patients with DN attending an ongoing local CM service programme based on the captioned protocol shows stabilised glomerular filtration rate (GFR), reduced albuminuria and reduced glycated haemoglobin (HbA1c) level. Recent evidence from *in vitro* and *in vivo* studies suggested that the effect of these herbal medicines in reducing proteinuria and alleviating decline of renal function could be associated with attenuated podocyte apoptosis and detachment,^{19–21} reduced oxidative stress^{21 22} and suppressed transforming growth factor- β (TGF- β) production,^{23 24} through some of the important pathways including TGF- β /Smad,^{23 24} mitogen-activated protein kinase/extracellular signal-regulated kinase (MAPK/ERK)²⁵ and nuclear factor kappa B (NF- κ B)²⁶ signalling pathway. In this pilot study, we aim to explore the effect of the captioned protocol on renal outcomes when used as an add-on therapy for patients with DN in the existing healthcare setting.

OBJECTIVE


To optimise parameters and assess the feasibility for a subsequent phase III randomised controlled trial (RCT) through preliminary evaluation on the effect of the adjuvant semi-individualised CM treatment protocol on patients with type 2 diabetes with stages 2–3 chronic kidney disease (CKD) and macroalbuminuria.

METHOD/DESIGN

Study design

This is an assessor-blind, add-on, randomised, controlled, parallel, multicentre, open-label pragmatic clinical trial. The flow of study is summarised in [figure 1](#).

Figure 1 Flow of study.



Inclusion and exclusion criteria

Patients: (1) diagnosed with type 2 diabetes for at least 5 years; (2) with an estimated GFR $\geq 30 < 90$ mL/min/ 1.73 m^2 confirmed with repeat testing over 3 or more months calculated by the abbreviated Modification of Diet in Renal Disease (MDRD) study equation;²⁷ (3) persistent macroalbuminuria with spot urine albumin-to-creatinine ratio (UACR) ≥ 300 mg/g confirmed by at least two consecutive first morning void urine samples; (4) aged between 35 and 80 years; (5) on a stable dose of antidiabetic drug including insulin for 12 weeks and (6) on a stable dose of ACE inhibitor or angiotensin receptor blocker for 12 weeks will be recruited.

Patients: (1) with a known history of glomerulonephritis, polycystic kidney disease, systemic lupus erythematosus or any suggestive evidence of non-diabetic glomerulopathy; (2) with a known history of kidney transplant; (3) with concurrent severe disorders of the heart, brain, liver and haematopoietic system, tumour, mental disorder; (4) deranged liver function; (5) with poorly controlled blood pressure; (6) with a known history of intolerance or malabsorption of oral medications; (7) uncontrollable urinary infection; (8) experiencing pregnancy and (9) participating in other clinical trials within 30 days will be excluded.

Sample size calculation

Currently, there is no consensus on the sample size calculation for pilot studies. Previous studies suggested a sample size of 12–30 per arm.^{28–31} Since the primary objective of this trial is to explore the feasibility of a subsequent main study and have a preliminary evaluation on the effect of CM on patients with DN, the sample size calculation is based on control of the inflation factor (IF) to the estimation of sample size for the subsequent main study.³¹

The standard deviation used for sample size calculation for the main study could be underestimated by a pilot study and should be adjusted by an IF.³¹ IF is calculated based on the size of the pilot study and the confidence level of achieving at least the desired power in the main study. The actual achieved power of the main study depends on the nominal power set for the main study and the IF.

In order to be 95% confident (two-sided) that the subsequent main study actually achieves a power of 80% with nominal power set at 90% (ie, a 10% power forfeit), the IF should be < 1.15 . At $IF = 1.15$, a sample size of 80 is therefore needed to have 95% one-sided confidence that the main study will achieve at least the nominal power to test the hypothesis that an add-on of the whole CM treatment programme could be more effective in stabilising the GFR among patients with DN when compared with having standard care alone. For subgroup analysis, a sample size of 25 patients per subgroup could achieve 80% one-sided confidence that the effect of stabilising GFR is different within subgroups of similar CM clinical pattern.

With five subgroups, a sample size of 148 patients is needed in this pilot trial to allow a 15% attrition rate.

Recruitment and setting

Patients are recruited from Queen Mary Hospital Specialist Renal Outpatients Clinic, Sai Ying Pun Family Medicine Specialty Clinic, Sai Ying Pun Jockey Club General Out-patient Clinic, Kennedy Town Jockey Club General Out-patient Clinic, Aberdeen Family Medicine Specialty Clinic, Aberdeen Jockey Club General Out-patient Clinic; and Diabetes Complication Screening Clinic and Renal Clinic of Tung Wah Hospital. Patients diagnosed with diabetes and CKD stage 2–3 with macroalbuminuria are referred to the principal investigator (PI), co-investigator (Co-I), or Chinese medicine practitioner (CMP) for further assessment. The aim, procedures, nature of study, possible side effects of the CM treatment programme and right of unconditional withdrawal are explained by PI or Co-I before written consent is obtained from each patient who agrees to take part in the study.

All patients will undergo a 2-week run-in period, during which they are asked to keep their medication dosage fixed. A blood sample will be taken and tested to check liver function (γ -glutamyltransferase, alanine transaminase (ALT), alkaline phosphatase, total protein, aspartate transaminase (AST), bilirubin, albumin) and renal function (sodium, chloride, urea, potassium, creatinine) in the Department of Pathology & Clinical Biochemistry of Queen Mary Hospital or local clinical laboratories of other hospitals. Patients are considered eligible for the study if their liver functions are normal.

A random sequence is generated by and encrypted with computer by an independent research assistant (RA). Sealed opaque envelopes containing an allocation sequence are sent to the outpatient clinics. The password of the sequence is kept in a sealed, duly-signed opaque envelope locked by RA and Co-I. The allocation sequence is concealed from PI and Co-Is responsible for patient screening, recruitment and assignment of CM subgroup stratification. Eligible candidates are randomised to either the active treatment arm or control arm. The allocation will be explained by the CMP I or II to the patient. The allocation will be masked from the outcome assessor. The participant could not be masked due to the nature of treatment. Since the clinical outcomes under investigation are objective, placebo effect should be minimised. The distribution of known and unknown confounding factors among the intervention group and control group is balanced by the randomisation.

Intervention and control

The intervention under investigation is a semi-individualised CM treatment protocol used on top of standard medical care. The programme consists of five different CM herbal formulations corresponding to five

different subgroups of patients stratified by clinical manifestations according to classical CM theory:

A, for the subgroup of 'spleen and kidney Qi deficiency':

Radix Aucklandiae (Mu Xiang), Rhizoma Pinelliae Praeparatum (Fa Ban Xia), Rhizoma Atractylodis Macrocephalae (Bai Zhu), Poria (Fu Ling), Radix et Rhizoma Glycyrrhizae Praeparata cum Melle (Zhi Gan Cao), Fructus Amomi (Sha Ren), Pericarpium citri reticulatae (Chen Pi), Radix et Rhizoma Ginseng (Ren Shen), Radix Rehmanniae Praeparata (Shu Di Huang), Fructus Corni (Shan Zhu Yu), Rhizoma Dioscoreae (Shan Yao), Cortex Moutan (Mu Dan Pi), Rhizoma Alismatis (Ze Xie).

B, for the subgroup of 'spleen and kidney Yang deficiency':

Rhizoma Atractylodis Macrocephalae (Bai Zhu), Fructus Chaenomelis (Mu Gua), Radix Aucklandiae (Mu Xiang), Fructus Tsaoko (Cao Guo), Semen Arecae (Bing Lang), Radix Aconiti Lateralis Praeparata (Hei Shun Pian), Poria (Fu Ling), Rhizoma Zingiberis (Gan Jiang), Radix et Rhizoma Glycyrrhizae Praeparata cum Melle (Zhi Gan Cao), Rhizoma Zingiberis Recens (Sheng Jiang), Fructus Jujubae (Da Zao), Radix Rehmanniae Praeparata (Shu Di Huang), Fructus Corni (Shan Zhu Yu), Rhizoma Dioscoreae (Shan Yao), Cortex Moutan (Mu Dan Pi), Rhizoma Alismatis (Ze Xie), Ramulus Cinnamomi (Gui Zhi).

C, for the subgroup of 'spleen and kidney Qi and Ying deficiency':

Radix et Rhizoma Ginseng (Ren Shen), Radix Astragali (Huang Qi), Radix Rehmanniae Praeparata (Shu Di Huang), Fructus Corni (Shan Zhu Yu), Rhizoma Dioscoreae (Shan Yao), Poria (Fu Ling), Cortex Moutan (Mu Dan Pi), Rhizoma Zingiberis Recens (Sheng Jiang), Fructus Jujubae (Da Zao).

D, for the subgroup of 'liver and kidney Ying deficiency':

Radix Rehmanniae Praeparata (Shu Di Huang), Fructus Corni (Shan Zhu Yu), Rhizoma Dioscoreae (Shan Yao), Poria (Fu Ling), Cortex Moutan (Mu Dan Pi), Rhizoma Alismatis (Ze Xie), Fructus Ligustri Lucidi (Nv Zhen Zi), Herba Ecliptae (Mo Han Lian).

E, for the subgroup of 'Ying and Yang deficiency':

Radix Rehmanniae Praeparata (Shu Di Huang), Fructus Corni (Shan Zhu Yu), Rhizoma Dioscoreae (Shan Yao), Poria (Fu Ling), Cortex Moutan (Mu Dan Pi), Rhizoma Alismatis (Ze Xie), Ramulus Cinnamomi (Gui Zhi), Radix Aconiti Lateralis Praeparata (Hei Shun Pian), Fructus Ligustri Lucidi (Nv Zhen Zi), Herba Ecliptae (Mo Han Lian).

Patients with diabetic kidney disease provided consent are further assessed by the CMP and categorised into one of the five different CM subgroups based on the Guiding Principle of Clinical Research on New Drugs of Chinese Medicine.³² The CM diagnoses are based on clinical manifestation as follows:

A, for the subgroup of 'spleen and kidney Qi deficiency':

Primary manifestation: fatigue, shortness of breath, torpid intake, lumbago.

B, for the subgroup of 'spleen and kidney Yang deficiency':

Primary manifestation: fear of cold, fatigue, shortness of breath, torpid intake, lumbago.

C, for the subgroup of 'spleen and kidney Qi and Ying deficiency':

Primary manifestation: fatigue, lumbago, dry mouth, heat vexation.

D, for the subgroup of 'liver and kidney Ying deficiency':

Primary manifestation: dizziness or headache, lumbago, dry mouth, heat vexation.

E, for the subgroup of 'Ying and Yang deficiency':

Primary manifestation: fear of cold, lumbago, dry mouth, heat vexation.

Patients with at least one episode of all the primary manifestations of each CM subgroup listed above during the 2 weeks before consultation are considered eligible for the corresponding group. To enhance the specificity of categorisation, patients having multiple manifestations that fit more than one subgroup are excluded.

The treatment programme is originated from an existing clinical practice guideline being used in mainland China derived from expert consensus.¹⁸ Patients randomised to the intervention arm will receive the corresponding formulations on top of standard medication, as stated above. Minor adjustment of the herbs is allowed based on an individual's condition and determined by the CMP to reflect actual clinical practice. All patients continue their standard medications and follow-ups. Patients randomised to the control group will receive consultation from the CMP at baseline, study midpoint (week 24) and study end point (week 48) in addition to standard care. Patients will have 5 days of medicine per week and are advised to take the medication twice per day dissolved in boiling water in the first 5 days of the week.

Herbal safety

Herbal medicines with known toxicity that are listed in Schedule 1, Chapter 549, Chinese Herbal Medicines, Chinese Medicine Ordinance of Hong Kong are excluded. Soluble herbal granules prepared by PuraPharm are used in this study. The production process is in strict compliance with standards of Good Manufacturing Practice (GMP). The CM is prescribed and dispensed through the clinics of the School of Chinese Medicine, Li Ka Shing Faculty of Medicine or Clinical Trial Pharmacy of Queen Mary Hospital complying with the standard of practice of pharmacy. A fully registered CMP is responsible for clinical diagnosis and prescription. At week 6, patients will undergo liver function tests and brain natriuretic peptide (BNP) test for monitoring of liver toxicity and cardiotoxicity. To further ensure patient safety, we only recruit patients with stages 2–3 CKD who are more clinically stable to participate in the study.

Outcome measurement

The primary outcome measures are the changes in the estimated GFR and spot UACR between the baseline (week 0) and treatment end point (week 48). Secondary measures include changes from baseline (week 0) to end of treatment (week 48) in fasting blood glucose (FBG), HbA1c, serum BNP, fasting serum insulin, serum C peptide, serum fibroblast growth factor 23 (FGF23), urinary monocyte chemotactic protein-1 (MCP-1), urinary cystatin C, urinary nephrin, urinary TGF- β 1 and urinary vascular endothelial growth factor (VEGF). A self-complete questionnaire is distributed to the participants to monitor occurrence of adverse events, and they are advised to inform the PI or RA immediately.

Estimated GFR, UACR, FBG, HbA1c and liver function test are ordered by the Renal Clinic and assessed by the Division of Clinical Biochemistry, Queen Mary Hospital or local clinical laboratories of other hospitals which are accredited by the College of American Pathologists. BNP, insulin, C peptide, FGF23, MCP-1, cystatin C, nephrine, TGF- β 1 and VEGF assessment will be performed at PI's research laboratory by an independent RA. Estimated GFR is calculated using the abbreviated 4-variable MDRD study equation with serum creatinine, age, ethnicity and gender. Clinical presentations and CM syndromic diagnosis will be assessed in a structured consultation developed for this purpose. To ensure consistency and reliability of assessment and minimise bias from investigators across the study, patients will be assessed by the same CMP during each visit based on standard CM practice. Follow-up consultations are held for all patients weekly in the first month and monthly subsequently. Evaluation of outcomes will be performed at weeks 24 (on treatment) and 48 (end of treatment). The follow-up schedule is summarised below in [table 1](#).

The baseline demographics including age, sex, body mass index, duration of diabetes, duration of diagnosed diabetic kidney disease and concurrent medications used are collected through the clinical management

system of the Specialist Renal Outpatient Clinic. Patient flow measured as enrolled case per month and follow-up rate are recorded for the consideration of subsequent RCT design.

Handling of withdrawal and dropout

In order to maximise participants' compliance, first, we have a thorough consent process for all participants with details of the study schedule, potential side effects of treatment, and the responsibilities of the participants explained. Second, we scrutinise potential participants carefully during a 2-week run-in period in order to exclude ineligible and low compliance participants before randomisation. Third, an independent email account and a direct telephone line associated with this clinical trial have been set up to enable active communication with patients. Moreover, extra visits will be arranged for patients if necessary. If any patient plans to withdraw or drop out, the study team will communicate with the patient to determine and resolve the problem in order to retain the participant.

Termination criteria

The trial will be terminated for a specific participant if she or he has: (1) presence of serious adverse effect(s) in the interventional group; (2) hypersensitivity towards Chinese herbal medicine and (3) participation in another clinical trial. The whole study will be terminated under the following circumstances: (1) presence of serious adverse effect(s) related to Chinese herbal medicine with supportive evidence and (2) completion of all follow-up assessments.

Serious adverse events include adverse events that result in death, require either hospitalisation or the prolongation of hospitalisation, are life-threatening, result in a persistent or significant disability/incapacity or result in a congenital anomaly/birth defect. Other important medical events, based on appropriate medical judgement, may also be considered serious adverse events if a patient's health is at risk and intervention is required to prevent an outcome mentioned.

Table 1 Follow-up schedule

	Enrolment Before treatment (up to 30 days prior to baseline)	Allocation Week 0, day 1	Postallocation—treatment period		
			Weeks 1–4 (\pm 3 days)	After 6 weeks (\pm 7 days)	After 24 and 48 weeks (\pm 7 days)
Screening of eligibility	X				
Informed consent	X				
Medical history	X				
Renal and liver function test	X			X	X
Randomised allocation		X			
Intervention (for interventional group)		X	X	X	X
Standard care (for all patients)		X	X	X	X
Outcome assessment (Spot urine+blood)		X			X
Adverse events		X	X	X	X



Data analysis

Missing values, if any, will be imputed with regression. The analysis will follow the intention-to-treat approach. Patients without a postrandomisation assessment for a particular efficacy end point will be excluded from the analysis of that end point. SAS and SPSS will be used for the data analysis.

The demographics of each group will be presented as means. Sex will be presented as the male:female ratio. Regression analyses will be used to compare the adjusted mean of estimated GFR, UACR, HbA1c, FBG, BNP, insulin, C peptide, FGF23, MCP-1, cystatin C, nephrin, TGF- β 1 and VEGF at week 48 between (1) the combination of all intervention groups and combination of all control groups, (2) individual treatment subgroup and its matching control group and (3) different control groups with the corresponding baseline values as covariates. Data will be presented as the difference in adjusted means between the intervention arm and control arm with a 95% CI and the corresponding p value. A change score analysis will be performed as supplementary analysis.

The adverse events are recorded based on the Common Terminology Criteria for Adverse Events V4.03 and will be analysed in a narrative manner. The adverse events are categorised into five grades: grade (1) mild; asymptomatic or mild symptoms; clinical or diagnostic observations only; no intervention indicated; grade (2) moderate; minimal, local or non-invasive intervention indicated; limiting age-appropriate instrumental activities of daily living; grade (3) severe or medically significant but not immediately life-threatening; hospitalisation or prolongation of hospitalisation indicated; disabling; limiting self-care activities of daily living; grade (4) life-threatening consequences; urgent intervention indicated; and grade (5) death related to adverse events. The percentage of all adverse events and the rate of attrition due to adverse events will be compared between intervention groups and control groups.

To minimise type I error inflation, the analysis will follow a hierarchical approach in the order of (1) individual treatment subgroup versus its control subgroup and (2) combination of all intervention groups versus the combination of all control groups as to avoid type I error inflation. The primary outcomes are the change of GFR and UACR.

Subgroup analysis will be performed for CKD stages 2 and 3 separately. Sensitivity analyses will be performed for (1) missing data imputed with regression, (2) missing data imputed with last-observation-carried-forward (LOCF) and (3) per protocol dropout of patients. Adverse events will be analysed case-by-case and presented descriptively.

Data management

A trial management committee is formed by the study team members. Co-Is and RA collect, clean and send the study data to the committee on a weekly basis. All data are double-entered to computer and cleaned

before analysis to prevent data entry errors. All data transfer is encrypted to protect patients' confidentiality. The committee holds meetings monthly with experts to discuss the progress of the trial.

ETHICS CONSIDERATION

This trial is ethically justified as there are uncertainties on the therapeutic effect and adverse effect of the treatment programme which is generated by expert consensus but with no further validation of clinical evidence. Since the treatment is being used in mainland China and has the potential to be broadly used in Hong Kong and benefit local citizens, we believe this trial is worthwhile. All protocol amendments will be approved by the Institutional Review Board prior to execution.

Author affiliations

¹Division of Nephrology, Department of Medicine, The University of Hong Kong, Hong Kong, Hong Kong

²Department of Medicine, Tung Wah Hospital, Hong Kong, Hong Kong

³Department of Family Medicine and Primary Healthcare, Queen Mary Hospital, Hong Kong, Hong Kong

⁴Division of Nephrology, Department of Medicine, Queen Mary Hospital, Hong Kong, Hong Kong

⁵Division of Epidemiology and Biostatistics, School of Public Health, The University of Hong Kong, Hong Kong, Hong Kong

⁶School of Chinese Medicine, The University of Hong Kong, Hong Kong, Hong Kong

⁷Division of Endocrinology, Department of Medicine, The University of Hong Kong, Hong Kong, Hong Kong

Contributors KWC and SCWT conceived and designed the study. TPI, ASKK, SLL, GCWC, KCBT and YF contributed to the trial design and study operation. BJC contributed to the statistical design and analysis. WHY, DWLW, YL, JCKL, LYC and KNL contributed to the design and operation of biochemical sampling. All the authors made significant contributions to the development and conceptualisation of the protocol. All the authors reviewed the draft versions of this paper and have read and approved the final manuscript.

Funding This study is conducted as an investigator-initiated study with financial support from the Health and Medical Research Fund (project number 12133341) administered by the Food and Health Bureau, the Government of the Hong Kong Special Administrative Region.

Disclaimer The funding agency has no role in the design of this study and will not have any role during its execution, analyses, interpretation of the data or decision to submit results.

Competing interests None declared.

Patient consent Obtained.

Ethics approval This protocol is approved by the Institutional Review Board of the University of Hong Kong/Hospital Authority Hong Kong West Cluster (HKU/HA HKW IRB; reference number UW 14-301). A WHO Trial Registration Data Set is presented below.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

1. Tang SC. Diabetic nephropathy: a global and growing threat. *Hong Kong Med J* 2010;16:244–5.

2. Thomas MC, Cooper ME, Zimmet P. Changing epidemiology of type 2 diabetes mellitus and associated chronic kidney disease. *Nat Rev Nephrol* 2016;12:73–81.
3. International Diabetes Federation. *IDF Diabetes Atlas Sixth Edition: International Diabetes Federation*. 2014.
4. Shaw JE, Sicree RA, Zimmet PZ. Global estimates of the prevalence of diabetes for 2010 and 2030. *Diabetes Res Clin Pract* 2010;87:4–14.
5. Tuttle KR, Bakris GL, Bilous RW, *et al*. Diabetic kidney disease: a report from an ADA Consensus Conference. *Am J Kidney Dis* 2014;64:510–33.
6. Palmer SC, Mavridis D, Navarese E, *et al*. Comparative efficacy and safety of blood pressure-lowering agents in adults with diabetes and kidney disease: a network meta-analysis. *Lancet* 2015;385:2047–56.
7. Leung CB, Cheung WL, Li PK. Renal registry in Hong Kong—the first 20 years. *Kidney Int Suppl* 2015;5:33–8.
8. ERA-EDTA Registry. *ERA-EDTA Registry Annual Report 2013*. Amsterdam: ERA-EDTA Registry, 2015:1–148.
9. Karhaloo A. Anti-fibrosis therapy and diabetic nephropathy. *Curr Diab Rep* 2012;12:414–22.
10. Ban CR, Twigg SM. Fibrosis in diabetes complications: pathogenic mechanisms and circulating and urinary markers. *Vasc Health Risk Manag* 2008;4:575–96.
11. Liu Y. Renal fibrosis: new insights into the pathogenesis and therapeutics. *Kidney Int* 2006;69:213–17.
12. Lin M, Yiu WH, Wu HJ, *et al*. Toll-like receptor 4 promotes tubular inflammation in diabetic nephropathy. *J Am Soc Nephrol* 2012;23:86–102.
13. Tang SC, Lai KN. The pathogenic role of the renal proximal tubular cell in diabetic nephropathy. *Nephrol Dial Transplant* 2012;27:3049–56.
14. Chan GC, Tang SC. Diabetic nephropathy: landmark clinical trials and tribulations. *Nephrol Dial Transplant* 2016;31:359–68.
15. Census and Statistics Department. *Thematic Household Survey Report No. 45*. In: Department CaS, ed. Hong Kong 2010.
16. Chung VC, Ma PH, Lau CH, *et al*. Views on traditional Chinese medicine amongst Chinese population: a systematic review of qualitative and quantitative studies. *Health Expect* 2014;17:622–36.
17. Lin MY, Chiu YW, Chang JS, *et al*. Association of prescribed Chinese herbal medicine use with risk of end-stage renal disease in patients with chronic kidney disease. *Kidney Int* 2015;88:1365–73.
18. Zhang L. *The optimization study of clinical pathway formulation based on chronic renal failure diagnosis scheme*. Guangzhou University of Chinese Medicine, 2010.
19. Chen J, Chen Y, Luo Y, *et al*. Astragaloside IV ameliorates diabetic nephropathy involving protection of podocytes in streptozotocin induced diabetic rats. *Eur J Pharmacol* 2014;736:86–94.
20. Chen J, Gui D, Chen Y, *et al*. Astragaloside IV improves high glucose-induced podocyte adhesion dysfunction via alpha3beta1 integrin upregulation and integrin-linked kinase inhibition. *Biochem Pharmacol* 2008;76:796–804.
21. Gui D, Guo Y, Wang F, *et al*. Astragaloside IV, a novel antioxidant, prevents glucose-induced podocyte apoptosis in vitro and in vivo. *PLoS ONE* 2012;7:e39824.
22. Liu HR, Tang XY, Dai DZ, *et al*. Ethanol extracts of Rehmannia complex (Di Huang) containing no Corni fructus improve early diabetic nephropathy by combining suppression on the ET-ROS axis with modulate hypoglycemic effect in rats. *J Ethnopharmacol* 2008;118:466–72.
23. Wang Y, Lin C, Ren Q, *et al*. Astragaloside effect on TGF- β 1, SMAD2/3, and α -SMA expression in the kidney tissues of diabetic KKAY mice. *Int J Clin Exp Pathol* 2015;8:6828–34.
24. Nie Y, Li S, Yi Y, *et al*. Effects of astragalus injection on the TGF β /Smad pathway in the kidney in type 2 diabetic mice. *BMC Complement Altern Med* 2014;14:1472–6882.
25. Zheng R, Deng Y, Chen Y, *et al*. Astragaloside IV attenuates complement membranous attack complex induced podocyte injury through the MAPK pathway. *Phytother Res* 2012;26:892–8.
26. Gui D, Huang J, Guo Y, *et al*. Astragaloside IV ameliorates renal injury in streptozotocin-induced diabetic rats through inhibiting NF- κ B-mediated inflammatory genes expression. *Cytokine* 2013;61:970–7.
27. Levey AS, Bosch JP, Lewis JB, *et al*. A more accurate method to estimate glomerular filtration rate from serum creatinine: a new prediction equation. Modification of Diet in Renal Disease Study Group. *Ann Intern Med* 1999;130:461–70.
28. Hertzog MA. Considerations in determining sample size for pilot studies. *Res Nurs Health* 2008;31:180–91.
29. Julious SA. Sample size of 12 per group rule of thumb for a pilot study. *Pharm Stat* 2005;4:287–91.
30. Lancaster GA, Dodd S, Williamson PR. Design and analysis of pilot studies: recommendations for good practice. *J Eval Clin Pract* 2004;10:307–12.
31. Sim J, Lewis M. The size of a pilot study for a clinical trial should be calculated in relation to considerations of precision and efficiency. *J Clin Epidemiol* 2012;65:301–8.
32. Zheng XY. *Guiding Principle of Clinical Research on New Drugs of Chinese Medicine (Trial)*. Beijing: China Medical Science Press, 2002.

World Health Organisation Trial Registration Data Set

Data category	Information
Primary Registry and Trial Identifying Number	ClinicalTrials.gov NCT02488252
Date of registration in primary registry	25 June 2015
Secondary Identifying Numbers	HMRF-12133341
Source(s) of monetary or material support	Health and Medical Research Fund
Primary sponsor	The University of Hong Kong
Secondary sponsor(s)	Not applicable
Contact for public queries	Professor Sydney CW Tang, MD, PhD Tel: +852 2255 3879 Email: scwtang@hku.hk
Public title	Semi-individualised Chinese medicine treatment as an adjuvant management for diabetic nephropathy
Scientific title	Semi-individualised Chinese medicine treatment as an adjuvant management for diabetic nephropathy: a pilot add-on, randomised, controlled, multicentre, open-label pragmatic clinical trial
Countries of recruitment	Hong Kong Diabetic nephropathy

Continued



Continued

Data category	Information
Health condition(s) or problem (s) studied	
Intervention(s)	Active comparator: Standard medical care with ACE inhibitor or angiotensin receptor blocker and oral hypoglycaemic agents and/or insulin at stable dose Experimental arm: Semi-individualised Chinese medicine treatment programme on top of standard medical care
Key inclusion and exclusion criteria	Ages eligible for study: between 35 and 80 years Gender eligible for study: both Healthy volunteers: not accepted Inclusion criteria: <ul style="list-style-type: none"> ▶ Diagnosed with type 2 diabetes for at least 5 years; ▶ With an estimated glomerular filtration rate (GFR) $\geq 30 < 90$ mL/min/1.73 m² confirmed with repeat testing over 3 or more months calculated by the abbreviated MDRD study equation; ▶ Persistent macroalbuminuria with spot urine albumin-to-creatinine ratio (UACR) ≥ 300 mg/g confirmed by at least 2 of the 3 consecutive first morning void urine samples; ▶ On a stable dose of antidiabetic drug including insulin for 12 weeks; ▶ On a stable dose of ACE inhibitor or angiotensin receptor blocker for 12 weeks; ▶ Willing and able to give written informed consent. Exclusion criteria: <ul style="list-style-type: none"> ▶ With a known history of glomerulonephritis, polycystic kidney disease, systemic lupus erythematosus, any suggestive evidence of non-diabetic glomerulopathy; ▶ With a known history of kidney transplant; ▶ With concurrent severe disorders of the heart, brain, liver, and haematopoietic system, tumour and mental disorder; ▶ With deranged liver function; ▶ With poorly controlled blood pressure; ▶ With a known history of intolerance or malabsorption of oral medications; ▶ With an uncontrollable urinary infection; ▶ Experiencing pregnancy; ▶ Participating in other clinical trials within 30 days.
Study type	Interventional Allocation: randomised Intervention model: parallel assignment (2 arms) Blinding: open-label (assessor of primary outcome measures blinded) Primary purpose: treatment Phase: II/III Allocation concealment: sealed opaque envelope prepared by an independent research assistant Sequence generation: computer-generated random sequence
Date of first enrolment	July 2015
Target sample size	148
Recruitment status	Recruiting
Primary outcome(s)	Changes in estimated GFR and spot UACR (time frame: 48 weeks)
Key secondary outcome(s)	Changes in fasting blood glucose, glycated haemoglobin, serum, brain natriuretic peptide, fasting serum insulin, serum C peptide, serum fibroblast growth factor 23, urinary monocyte chemoattractant protein-1, urinary cystatin C, urinary nephrin, urinary transforming growth factor- β 1 and urinary vascular endothelial growth factor

BMJ Open

Semi-individualised Chinese medicine treatment as an adjuvant management for diabetic nephropathy: a pilot add-on, randomised, controlled, multicentre, open-label pragmatic clinical trial

Kam Wa Chan, Tai Pang Ip, Alfred Siu Kei Kwong, Sing Leung Lui, Gary Chi Wang Chan, Benjamin John Cowling, Wai Han Yiu, Dickson Wai Leong Wong, Yang Liu, Yibin Feng, Kathryn Choon Beng Tan, Loretta Yuk Yee Chan, Joseph Chi Kam Leung, Kar Neng Lai and Sydney Chi Wai Tang

BMJ Open 2016 6:
doi: [10.1136/bmjopen-2015-010741](https://doi.org/10.1136/bmjopen-2015-010741)

Updated information and services can be found at:
<http://bmjopen.bmj.com/content/6/8/e010741>

These include:

References

This article cites 27 articles, 3 of which you can access for free at:
<http://bmjopen.bmj.com/content/6/8/e010741#BIBL>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Evidence based practice](#) (581)
[Health services research](#) (1157)
[Renal medicine](#) (107)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>