

On Box-Perfect Graphs

Guoli Ding*

Department of Mathematics, Louisiana State University, Baton Rouge, USA

Wenan Zang[†] and Qiulan Zhao[‡]

Department of Mathematics, The University of Hong Kong, Hong Kong, China

September 18, 2018

Abstract

Let $G = (V, E)$ be a graph and let A_G be the clique-vertex incidence matrix of G . It is well known that G is perfect iff the system $A_G \mathbf{x} \leq \mathbf{1}$, $\mathbf{x} \geq \mathbf{0}$ is totally dual integral (TDI). In 1982, Cameron and Edmonds proposed to call G box-perfect if the system $A_G \mathbf{x} \leq \mathbf{1}$, $\mathbf{x} \geq \mathbf{0}$ is box-totally dual integral (box-TDI), and posed the problem of characterizing such graphs. In this paper we prove the Cameron-Edmonds conjecture on box-perfectness of parity graphs, and identify several other classes of box-perfect graphs. We also develop a general and powerful method for establishing box-perfectness.

1 Introduction

A rational system $A\mathbf{x} \leq \mathbf{b}$ is called *totally dual integral* (TDI) if the minimum in the LP-duality equation

$$\max\{\mathbf{w}^T \mathbf{x} : A\mathbf{x} \leq \mathbf{b}\} = \min\{\mathbf{y}^T \mathbf{b} : \mathbf{y}^T A = \mathbf{w}^T; \mathbf{y} \geq \mathbf{0}\} \quad (1.1)$$

has an integral optimal solution, for every integral vector \mathbf{w} for which the minimum is finite. Edmonds and Giles [14] proved that total dual integrality implies primal integrality: if $A\mathbf{x} \leq \mathbf{b}$ is TDI and \mathbf{b} is integral, then both programs in (1.1) have integral optimal solutions whenever they have finite optimum. So the model of TDI systems serves as a general framework for establishing min-max results in combinatorial optimization (see Schrijver [22] for an comprehensive and in-depth account). As summarized by Schrijver [20], the importance of a min-max relation is twofold: first, it serves as an optimality criterion and as a good characterization for the corresponding optimization problem; second, a min-max relation frequently yields an elegant combinatorial theorem, and allows a geometrical representation of the corresponding problem in terms of a polyhedron. Many well-known results and difficult conjectures in combinatorial optimization can be rephrased as saying that a certain linear system is TDI; in particular, by Lovász' Replication Lemma [16], a graph G is

*Supported in part by NSF grant DMS-1500699.

[†]Supported in part by the Research Grants Council of Hong Kong.

[‡]Corresponding author. E-mail: qiulanzhao@163.com.

perfect if and only if the system $A_G \mathbf{x} \leq \mathbf{1}$, $\mathbf{x} \geq \mathbf{0}$ is TDI, where A_G is the clique-vertex incidence matrix of G . The reader is referred to Chudnovsky *et al.* [10, 12] for the proof of the Strong Perfect Graph Theorem and to Chudnovsky *et al.* [8] for recognition of perfect graphs.

A rational system $A\mathbf{x} \leq \mathbf{b}$ is called *box-totally dual integral* (box-TDI) if $A\mathbf{x} \leq \mathbf{b}$, $\mathbf{l} \leq \mathbf{x} \leq \mathbf{u}$ is TDI for all vectors \mathbf{l} and \mathbf{u} , where each coordinate of \mathbf{l} and \mathbf{u} is either a rational number or $\pm\infty$. By taking $\mathbf{l} = -\infty$ and $\mathbf{u} = \infty$ it follows that every box-TDI system must be TDI. Cameron and Edmonds [3, 5] proposed to call a graph G *box-perfect* if the system $A_G \mathbf{x} \leq \mathbf{1}$, $\mathbf{x} \geq \mathbf{0}$ is box-TDI; they also posed the problem of characterizing such graphs.

We make some preparations before presenting an equivalent definition of box-perfect graphs. Let $G = (V, E)$ be a graph (all graphs considered in this paper are simple unless otherwise stated). For any $X \subseteq V$, let $G[X]$ denote the subgraph of G induced by X . For any $v \in V$, let $N_G(v)$ denote the set of vertices incident with v . Members of $N_G(v)$ are called *neighbors* of v . By *duplicating* a vertex v of G we obtain a new graph G' constructed as follows: we first add a new vertex v' to G , which may or may not be adjacent to v , and then we join v' to all vertices in $N_G(v)$.

As usual, let $\alpha(G)$ and $\chi(G)$ denote respectively the stable number and chromatic number of G . Let $\bar{\chi}(G) = \chi(\bar{G})$, which is the clique cover number of G . For any integer $q \geq 1$, let

$$\alpha_q(G) = \max\{|X| : X \subseteq V(G) \text{ with } \chi(G[X]) \leq q\}, \text{ and}$$

$$\bar{\chi}_q(G) = \min\{q\bar{\chi}(G - X) + |X| : X \subseteq V(G)\}.$$

Notice that $\alpha_1 = \alpha$ and $\bar{\chi}_1 = \bar{\chi}$. A graph G is called *q-perfect* if $\alpha_q(G[X]) = \bar{\chi}_q(G[X])$ holds for all $X \subseteq V(G)$. This concept was introduced by Lovász [17] as an extension of perfect graphs, since 1-perfect graphs are precisely perfect graphs. Let us call a graph *totally perfect* if it is q -perfect for all integers $q \geq 1$. Lovász pointed out that comparability graphs, incomparability graphs, and line graphs of bipartite graphs are totally perfect. However, S_3 is not 2-perfect, showing that a perfect graph does not have to be q -perfect when $q > 1$.




Figure 1.1: Graph S_3 and its complement \bar{S}_3

Theorem 1.1 (Cameron [4]). *A graph is box-perfect if and only if every graph obtained from this graph by repeatedly duplicating vertices is totally perfect.*

This theorem implies the following immediately.

Corollary 1.2 (Cameron [4]). (1) *Induced subgraphs of a box-perfect graph are box-perfect.*
 (2) *Duplicating vertices in a box-perfect graph results in a box-perfect graph.*
 (3) *Comparability and incomparability graphs are box-perfect.*

The next proposition contains a few other important observations made by Cameron [4]. A matrix A is *totally unimodular* if the determinant of every square submatrix of A is 0 or ± 1 . A

$\{0, 1\}$ -matrix A is *balanced* if none of its submatrices is the vertex-edge incidence matrix of an odd cycle. For each graph G , let B_G be the submatrix of A_G obtained by keeping only rows that correspond to maximal cliques of G . Let us call G *totally unimodular* or *balanced* if B_G is totally unimodular or balanced. It is worth pointing out that bipartite graphs and their line graphs are totally unimodular, and every totally unimodular graph is balanced. In addition, as shown by Berge [1], all balanced graphs are totally perfect. Let \bar{S}_3^+ be obtained from the complement \bar{S}_3 of S_3 by adding a new vertex v and joining v to all six vertices of \bar{S}_3 .

Proposition 1.3 (Cameron [4]). (1) \bar{S}_3^+ is not box-perfect.

(2) Totally unimodular graphs are box-perfect.

(3) Balanced graphs do not have to be box-perfect, shown by \bar{S}_3^+ .

(4) The complement of a box-perfect graph does not have to be box-perfect, shown by \bar{S}_3 .

(5) Box-perfectness is not preserved under taking clique sums, shown by S_3 .

As we have seen, many nice properties of perfect graphs are not satisfied by box-perfect graphs. Another property of this kind is substitution: substituting a vertex of a box-perfect graph by a box-perfect graph does not have to yield a box-perfect graph, as shown by \bar{S}_3^+ (which is obtained by substituting a vertex of K_2 with \bar{S}_3). To our knowledge, almost none of the known summing operations that preserve perfectness can carry over to box-perfectness – this makes it extremely hard to obtain a structural characterization of box-perfect graphs!

At this point, the only known box-perfect graphs are totally unimodular graphs, comparability graphs, incomparability graphs, and p -comparability graphs (where $p \geq 1$ and 1-comparability graphs are precisely comparability graphs) [3, 5]. Cameron and Edmonds [3] conjectured that every parity graph is box-perfect. In this paper we confirm this conjecture and identify several other classes of box-perfect graphs, including claw-free box-perfect graphs. In the next section we construct a class \mathcal{R} of non-box-perfect graphs, from which we characterize box-perfect split graphs. It turns out that every minimal non-box-perfect graph that we know of is contained in a graph from \mathcal{R} . This observation raises the question: is it true that a graph G is box-perfect if and only if G does not contain any graph in \mathcal{R} as an induced subgraph?

In addition to structural description, the other difficulty with the study of box-perfect graphs lies in the lack of a proper tool for establishing box-perfectness. In section 3 we introduce a so-called ESP property, which is sufficient for a graph to be box-perfect. Although recognizing box-perfectness is an optimization problem, our approach based on the ESP property is of transparent combinatorial nature and hence is fairly easy to work with. For convenience, we call a graph ESP if it has the aforementioned ESP property. In the remainder of this paper, we shall establish several classes of box-perfect graphs by showing that they are actually ESP, including all classes obtained by Cameron [3, 4, 5]. We strongly believe that the ESP property is exactly the tool one needs for the study of box-perfect graphs.

Conjecture 1.4. *A perfect graph is box-perfect if and only if it is ESP if and only if it contains none of the members of \mathcal{R} as an induced subgraph.*

We close this section by mentioning a result on the complexity of recognizing box-perfect graphs.

Theorem 1.5 (Cook [13]). *The class of box-perfect graphs is in co-NP.*

2 A class of non-box-perfect graphs

Let S_n be the graph obtained from cycle $v_1v_2\dots v_{2n}v_1$ by adding edges v_iv_j for all distinct even i, j . It was proved in [4] that S_{2n+1} is not box-perfect for all $n \geq 1$. In this section we construct a class of non-box-perfect graphs, which include \bar{S}_3^+ and S_{2n+1} ($n \geq 1$). We will use this result to characterize box-perfect split graphs (a graph is *split* if its vertex set can be partitioned into a clique and a stable set).

Let $G = (U, V, E)$ be a bipartite graph, where $U = \{u_1, \dots, u_m\}$ and $V = \{v_1, \dots, v_n\}$. The *biadjacency matrix* of G is the $\{0, 1\}$ -matrix M of dimension $m \times n$ such that $M_{i,j} = 1$ if and only if $u_iv_j \in E$. Let \mathcal{Q} be the set of bipartite graphs G such that its biadjacency matrix M is not totally unimodular but all submatrices of M are. The following is a classical result of Camion.

Lemma 2.1 (Camion [6]). *Every graph $G = (U, V, E)$ in \mathcal{Q} is Eulerian. In addition, G satisfies $|U| = |V|$ and $|E| \equiv 2 \pmod{4}$.*

Let \mathcal{R} be the class of graphs constructed as follows. Take a bipartite graph $G' = (U, V, E') \in \mathcal{Q}$ and a graph $G'' = (V, E'')$ such that $N_{G''}(u)$ is a clique of G'' for all $u \in U$. Let $G = (U \cup V, E' \cup E'')$. If there exists $u \in U$ with $N_{G'}(u) = V$ then $G - u$ belongs to \mathcal{R} ; otherwise G belongs to \mathcal{R} .

Examples. For each odd $n \geq 3$, S_n belongs to \mathcal{R} since S_n can be constructed from a cycle $G' = C_{2n} \in \mathcal{Q}$ and a complete graph $G'' = K_n$, where no vertex is deleted in the construction. Graph \bar{S}_3^+ also belongs to \mathcal{R} . In this case a vertex is deleted in the construction, see Figure 2.1.




Figure 2.1: Graph \bar{S}_3^+ is constructed from a bipartite graph in \mathcal{Q} and K_4

Lemma 2.2. *No graph in \mathcal{R} is box-perfect.*

Proof. Let $G \in \mathcal{R}$ be constructed from $G' = (U, V, E') \in \mathcal{Q}$ and $G'' = (V, E'')$. Let A_G and B_G be the clique and maximal clique matrices of G . Then A_G can be expressed as $A_G = \begin{bmatrix} B_{G'} \\ C \end{bmatrix}$. Let M be the biadjacency matrix of G' and let $n := |U|$ ($= |V|$). Since every $u \in U$ belongs to exactly one maximal clique of G , the column of B_G that corresponds to u has precisely one nonzero entry. If no vertex was deleted in the construction of G then B_G can be expressed as $B_G = \begin{bmatrix} M & I_n \\ N & \mathbf{0} \end{bmatrix}$, where the first n columns are indexed by V and the last n columns are indexed by U . If a vertex $u_0 \in U$ was deleted in the construction of G , then G'' has to be a complete graph. In this case, since U does not have a second vertex adjacent to all vertices in V , B_G can be expressed as $[M, J]$, where $J_{n \times (n-1)} = \begin{bmatrix} I_{n-1} \\ \mathbf{0} \end{bmatrix}$ and the last row of M , which corresponds to u_0 , is a vector of all ones. By Lemma 2.1, all entries of $\mathbf{1}^T M$ and $M\mathbf{1}$ are even, and $\mathbf{1}^T M\mathbf{1} = 4m + 2$, for an integer $m > 0$. We consider the dual programs (with $A = A_G$)

$$\max\{\mathbf{w}^T \mathbf{x} : A\mathbf{x} \leq \mathbf{1}; \mathbf{x} \geq \mathbf{l}\} = \min\{\mathbf{y}^T \mathbf{1} - \mathbf{z}^T \mathbf{l} : \mathbf{y}^T A - \mathbf{z}^T = \mathbf{w}^T; \mathbf{y}, \mathbf{z} \geq \mathbf{0}\}. \quad (2.1)$$

Suppose no vertex was deleted in the construction of G . Let $p > 2m + 1$ be a prime and let

$$\mathbf{w} = \begin{bmatrix} \frac{1}{2}M^T\mathbf{1} \\ \mathbf{0} \end{bmatrix}, \quad \mathbf{l} = \begin{bmatrix} \mathbf{0} \\ \mathbf{1} - \frac{1}{2p}M\mathbf{1} \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} \frac{1}{2p}\mathbf{1} \\ \mathbf{1} - \frac{1}{2p}M\mathbf{1} \end{bmatrix}, \quad \mathbf{y} = \begin{bmatrix} \frac{1}{2}\mathbf{1} \\ \mathbf{0} \\ \mathbf{0} \end{bmatrix}, \quad \mathbf{z} = \begin{bmatrix} \mathbf{0} \\ \frac{1}{2}\mathbf{1} \end{bmatrix}.$$

Then it is routine to verify that \mathbf{w} is integral, $\mathbf{l} \geq \mathbf{0}$, and $\mathbf{x}, (\mathbf{y}, \mathbf{z})$ are feasible solutions to (2.1). Moreover $\mathbf{w}^T\mathbf{x} = \frac{2m+1}{2p} = \mathbf{y}^T\mathbf{1} - \mathbf{z}^T\mathbf{l}$, so $\mathbf{x}, (\mathbf{y}, \mathbf{z})$ are optimal solutions. Since the optimal value is not $\frac{1}{p}$ -integral, while \mathbf{l} is, it follows that the dual does not have an integral optimal solution and so G is not box-perfect. Next, suppose that a vertex was deleted in the construction of G . The proof for this case is almost identical to the proof for the last case. The only difference is that B_G has $2n - 1$ columns, instead of $2n$ columns. Thus we need to truncate the corresponding vectors. To be precise, let

$$\mathbf{w} = \begin{bmatrix} \frac{1}{2}M^T\mathbf{1} \\ \mathbf{0} \end{bmatrix}, \quad \mathbf{l} = \begin{bmatrix} \mathbf{0} \\ J^T(\mathbf{1} - \frac{1}{n}M\mathbf{1}) \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} \frac{1}{n}\mathbf{1} \\ J^T(\mathbf{1} - \frac{1}{n}M\mathbf{1}) \end{bmatrix}, \quad \mathbf{y} = \begin{bmatrix} \frac{1}{2}\mathbf{1} \\ \mathbf{0} \end{bmatrix}, \quad \mathbf{z} = \begin{bmatrix} \mathbf{0} \\ \frac{1}{2}\mathbf{1} \end{bmatrix}.$$

Using the fact that the last row of M is $\mathbf{1}^T$ we deduce that \mathbf{x} and (\mathbf{y}, \mathbf{z}) are feasible solutions, and $\mathbf{w}^T\mathbf{x} = \frac{2m+1}{n} = \mathbf{y}^T\mathbf{1} - \mathbf{z}^T\mathbf{l}$, which implies that both solutions are optimal. Furthermore, since $M\mathbf{1}$ is even and its last entry is n , we deduce that n is even and thus \mathbf{l} is $\frac{1}{n/2}$ -integral. However, the optimal value $\frac{2m+1}{n}$ is not $\frac{1}{n/2}$ -integral, so G is not box-perfect, which proves the theorem. \blacksquare

To identify all minimally non-box-perfect split graphs, we consider the following subsets of \mathcal{Q} . Let \mathcal{Q}_1 consist of all bipartite graphs $G = (U, V, E) \in \mathcal{Q}$ such that U has a vertex adjacent to all vertices of V . Let \mathcal{Q}_2 consist of all bipartite graphs $G = (U, V, E) \in \mathcal{Q} \setminus \mathcal{Q}_1$ such that the graph obtained from G by adding a vertex and making it adjacent to all vertices of V does not contain any graph in \mathcal{Q}_1 as an induced subgraph. Let \mathcal{S} consist of all graphs in \mathcal{R} that are constructed from a bipartite graph $G' \in \mathcal{Q}_1 \cup \mathcal{Q}_2$ and a complete graph G'' . It is clear that all members of \mathcal{S} are split graphs. Moreover, \bar{S}_3^+ and S_{2n+1} ($n \geq 1$) belong to \mathcal{S} .

Theorem 2.3. *The following are equivalent for any split graph G .*

- (1) G is box-perfect;
- (2) no graph in \mathcal{S} is an induced subgraph of G ;
- (3) G is totally unimodular.

Proof. Implication (3) \Rightarrow (1) follows from Proposition 1.3(2) and implication (1) \Rightarrow (2) follows from Lemma 2.2 and Corollary 1.2(1). To prove (2) \Rightarrow (3), let $G = (U, V, E)$ be a split graph, where U is a stable set and V is a clique. Let $G'' = G[V]$ and $G' = G \setminus E(G'')$. Let G''' be the bipartite graph obtained from G' by adding a vertex w adjacent to all vertices in V . Let M be the biadjacency matrix of G''' .

We first prove that M is totally unimodular. Suppose otherwise. Then G''' has an induced subgraph $H' \in \mathcal{Q}$. Let us choose H' so that H' contains the new vertex w whenever it is possible. Consequently, $H' \in \mathcal{Q}_1 \cup \mathcal{Q}_2$. Let H be constructed from H' and a complete graph H'' . Then $H \in \mathcal{S}$ and, by the construction of G , G contains H as an induced subgraph. This contradicts (2) and thus M has to be totally unimodular.

Let N be the biadjacency matrix of G' . Then $B_G = [N, I]$ or $\begin{bmatrix} N & I \\ \mathbf{1} & \mathbf{0} \end{bmatrix}$, depending on if V is a maximal clique of G . Notice that $M = \begin{bmatrix} N \\ \mathbf{1} \end{bmatrix}$. So B_G , and thus G , is totally unimodular. \blacksquare

This theorem shows that all minimally non-box-perfect split graphs are contained in \mathcal{S} . In fact, \mathcal{S} consists of precisely such graphs.

Theorem 2.4. *A split graph G belongs to \mathcal{S} if and only if G is not box-perfect but all its induced subgraphs are.*

Proof. The backward implication follows immediately from Theorem 2.3. To prove the forward implication, let $G \in \mathcal{S}$. By Lemma 2.2, we only need to show that $G - w$ is box-perfect for all $w \in V(G)$. Suppose G is constructed from a bipartite graph $G' = (U, V, E') \in \mathcal{Q}_1 \cup \mathcal{Q}_2$ and a complete graph $G'' = (V, E'')$. Let M be the biadjacency matrix of G' and let $n := |U| = |V|$. Observe that if $G' \in \mathcal{Q}_1$ then $B_G = [N, J]$, where $N = M$ and $J = \begin{bmatrix} I_{n-1} \\ \mathbf{0} \end{bmatrix}$; if $G' \in \mathcal{Q}_2$ then $B_G = [N, J]$, where $N = \begin{bmatrix} I \\ \mathbf{1} \end{bmatrix}$ and $J = \begin{bmatrix} I_n \\ \mathbf{0} \end{bmatrix}$.

Now it is straightforward to verify that, for each $u \in U$, $B_{G-u} = [N', J']$ is obtained from B_G by deleting the row and the column indexed by u ; for each $v \in V$, $B_{G-v} = [N', J']$ is obtained from B_G by deleting the column indexed by v and also possibly the last row. In both cases, N' is a proper submatrix of N . This implies that N' is totally unimodular and thus so is $[N', J']$. Consequently, $G - w$ is box-perfect (totally unimodular) for all $w \in V(G)$, which proves the theorem. ■

As we observed earlier that \bar{S}_3^+ and S_{2n+1} ($n \geq 1$) belong to \mathcal{S} . Thus these graphs are minimally non-box-perfect. We point out that, in addition to graphs in \mathcal{S} , other minimally non-box-perfect graphs can also be obtained using Lemma 2.2. For instance, the graph illustrated in Figure 2.2 is constructed from $G' = C_{10}$ and $G'' = C_5 + e$. By Lemma 2.2, this graph G is not box-perfect. However, G is not minimally non-box-perfect since $H = G - \{9, 0\}$ is not box-perfect, which is certified by vectors $\mathbf{w}^T = (1, 1, 1, 1, 1, 0, 0, 0)$, $\mathbf{l}^T = (0, 0, 0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$, $\mathbf{x}^T = (\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{3}{4}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$, $\mathbf{y}^T = (0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$, and $\mathbf{z}^T = (0, 0, 0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$, where the first row of B_H is the triangle 123. It can be shown that H is in fact minimally non-box-perfect because $H - x$ is totally unimodular for $x = 1, 2, 3, 4, 5, 6, 7$, and $H - 8$ has the ESP property defined in the next section which implies the box-perfectness.




Figure 2.2: A new non-box-perfect graph G

3 ESP graphs

In this section we introduce a so-called ESP property, which is sufficient for a graph to be box-perfect. We shall use this combinatorial property to identify several new classes of box-perfect graphs. We begin with a few lemmas.

Lemma 3.1 (Chen, Ding and Zang [7]). *Suppose \mathbf{a}_1 and \mathbf{a}_2 are rational vectors with $\mathbf{a}_1 \geq \mathbf{a}_2$, and b_1 and b_2 are rational numbers with $b_1 \leq b_2$. Then the system $\mathbf{A}\mathbf{x} \leq \mathbf{b}$, $\mathbf{a}_1^T \mathbf{x} \leq b_1$, $\mathbf{a}_2^T \mathbf{x} \leq b_2$,*

$\mathbf{x} \geq \mathbf{0}$ is box-TDI if and only if the system $A\mathbf{x} \leq \mathbf{b}$, $\mathbf{a}_1^T \mathbf{x} \leq b_1$, $\mathbf{x} \geq \mathbf{0}$ is box-TDI.

Lemma 3.2 (Cameron [4]). *The system $A\mathbf{x} \leq \mathbf{b}$ is box-TDI if and only if the system $A\mathbf{x} \leq \mathbf{b}$, $\mathbf{x} \leq \mathbf{u}$ is TDI, for all vectors \mathbf{u} , where each coordinate of \mathbf{u} is either a rational number or $+\infty$.*

The next two lemmas are reformulations of Theorem 22.7 and Theorem 22.13 of Schrijver [21].

Lemma 3.3 (Schrijver [21]). *Suppose the system $A\mathbf{x} \leq \mathbf{b}$, $x_1 \leq u$ is TDI for all rational numbers u , where x_1 is the first coordinate of \mathbf{x} . Then $A\mathbf{x} \leq \mathbf{b}$ is TDI.*

Lemma 3.4 (Schrijver [21]). *A rational system $A\mathbf{x} \leq \mathbf{b}$, $\mathbf{x} \geq \mathbf{0}$ is TDI if and only if $\min\{\mathbf{y}^T \mathbf{b} : \mathbf{y}^T A \geq \mathbf{w}^T, \mathbf{y} \geq \mathbf{0}\}$ is half-integral is finite and is attained by an integral \mathbf{y} , for each integral vector \mathbf{w} for which $\min\{\mathbf{y}^T \mathbf{b} : \mathbf{y}^T A \geq \mathbf{w}^T, \mathbf{y} \geq \mathbf{0}\}$ is finite.*

The next are two easy corollaries.

Lemma 3.5. *A graph G is box-perfect if and only if the system $B_G \mathbf{x} \leq \mathbf{1}$, $\mathbf{0} \leq \mathbf{x} \leq \mathbf{u}$ is TDI for all rational vectors $\mathbf{u} \geq \mathbf{0}$.*

Proof. The forward implication follows immediately from the definition of box-TDI and Lemma 3.1. Conversely, Lemma 3.2 and Lemma 3.3 imply that $B_G \mathbf{x} \leq \mathbf{1}$, $\mathbf{x} \geq \mathbf{0}$ is box-TDI. Then the result follows from Lemma 3.1. \blacksquare

Lemma 3.6. *A graph G is box-perfect if and only if for all rational $\mathbf{u} \geq \mathbf{0}$ and integral $\mathbf{w} \geq \mathbf{0}$,*

$$\begin{aligned} & \min\{\mathbf{y}^T \mathbf{1} + \mathbf{z}^T \mathbf{u} \mid \mathbf{y}^T B_G + \mathbf{z}^T \geq 2\mathbf{w}^T; \mathbf{y}, \mathbf{z} \geq \mathbf{0} \text{ integral}\} \\ & \geq 2 \min\{\mathbf{y}^T \mathbf{1} + \mathbf{z}^T \mathbf{u} \mid \mathbf{y}^T B_G + \mathbf{z}^T \geq \mathbf{w}^T; \mathbf{y}, \mathbf{z} \geq \mathbf{0} \text{ integral}\}. \end{aligned} \quad (3.1)$$

Proof. Observe that, for all vectors $\mathbf{u} \geq \mathbf{0}$ and \mathbf{w} , the three programs

$$\begin{aligned} & \min\{\mathbf{y}^T \mathbf{1} + \mathbf{z}^T \mathbf{u} \mid \mathbf{y}^T B_G + \mathbf{z}^T \geq \mathbf{w}^T; \mathbf{y}, \mathbf{z} \geq \mathbf{0}\} \\ & \min\{\mathbf{y}^T \mathbf{1} + \mathbf{z}^T \mathbf{u} \mid \mathbf{y}^T B_G + \mathbf{z}^T \geq \mathbf{w}^T; \mathbf{y}, \mathbf{z} \geq \mathbf{0} \text{ half-integral}\} \\ & \min\{\mathbf{y}^T \mathbf{1} + \mathbf{z}^T \mathbf{u} \mid \mathbf{y}^T B_G + \mathbf{z}^T \geq \mathbf{w}^T; \mathbf{y}, \mathbf{z} \geq \mathbf{0} \text{ integral}\} \end{aligned}$$

are finite. Moreover, replacing \mathbf{w} by \mathbf{w}_+ does not change the minimum values of these programs, where \mathbf{w}_+ is obtained from \mathbf{w} by turning its negative coordinates into zero. Therefore, the result follows immediately from Lemma 3.5 and Lemma 3.4. \blacksquare

Let $G = (V, E)$ be a graph. For any multiset Λ of cliques of G and any $v \in V$, let $d_\Lambda(v)$ denote the number of members of Λ that contain v . We call G *equitably subpartitionable (ESP)* if for every set Λ of maximal cliques of G there exist two multisets Λ_1 and Λ_2 of cliques of G (which are not necessarily members of Λ) such that

- (i) $|\Lambda_1| + |\Lambda_2| \leq |\Lambda|$;
- (ii) $d_{\Lambda_1}(v) + d_{\Lambda_2}(v) \geq d_\Lambda(v)$, for all $v \in V$; and
- (iii) $\min\{d_{\Lambda_1}(v), d_{\Lambda_2}(v)\} \geq \lfloor d_\Lambda(v)/2 \rfloor$, for all $v \in V$.

We call (Λ_1, Λ_2) an *equitable subpartition* of Λ , and refer to the above (i), (ii), and (iii) as *ESP property*. Note that (i) is equivalent to $|\Lambda_1| + |\Lambda_2| = |\Lambda|$ since we may include empty cliques in Λ_1 and Λ_2 . Similarly, (ii) is equivalent to $d_{\Lambda_1}(v) + d_{\Lambda_2}(v) = d_\Lambda(v)$ for all v , since cliques in Λ_1, Λ_2 can be replaced by smaller ones. Finally, it is also easy to see that in an ESP graph every multiset Λ of cliques admits an equitable subpartition. We will use these facts without further explanation.

Theorem 3.7. *Every ESP graph $G = (V, E)$ is box-perfect.*

Proof. By Lemma 3.6 we only need to show that inequality (3.1) holds for all rational $\mathbf{u} \geq \mathbf{0}$ and all integral $\mathbf{w} \geq \mathbf{0}$. Let $(\mathbf{y}^T, \mathbf{z}^T)$ be an optimal solution of the first minimum in (3.1). Let \mathcal{C} be the set of maximal cliques of G and let \mathcal{D} be the multiset of members of \mathcal{C} such that each $C \in \mathcal{C}$ appears in \mathcal{D} exactly y_C times. Let Λ be the set of $C \in \mathcal{C}$ such that y_C is odd. Since G is ESP, Λ admits a equitable subpartition (Λ_1, Λ_2) . Since every clique can be extended into a maximal clique, we may assume without loss of generality that members of Λ_1 and Λ_2 are all in \mathcal{C} . Let \mathcal{D}_0 be the multiset of members of \mathcal{C} such that each $C \in \mathcal{C}$ appears $\lfloor y_C/2 \rfloor$ times. It follows that $\mathcal{D} = \mathcal{D}_0 \uplus \mathcal{D}_0 \uplus \Lambda$, where \uplus stands for multiset sum. For $i = 1, 2$, let $\mathcal{D}_i = \mathcal{D}_0 \uplus \Lambda_i$. We deduce from (i) that

$$(1) \quad |\mathcal{D}_1| + |\mathcal{D}_2| \leq |\mathcal{D}|.$$

Let $\mathbf{p} = \mathbf{y}^T B_G + \mathbf{z}^T - 2\mathbf{w}^T$ and let $v \in V$. Without loss of generality we assume

$$(2) \quad d_{\mathcal{D}_1}(v) \geq d_{\mathcal{D}_2}(v) \quad \text{and} \quad \mathbf{p}_v \mathbf{z}_v = 0.$$

Since $d_{\mathcal{D}}(v) = 2d_{\mathcal{D}_0}(v) + d_{\Lambda}(v)$, we deduce from (ii-iii) that $d_{\mathcal{D}_1}(v) + d_{\mathcal{D}_2}(v) \geq d_{\mathcal{D}}(v)$ and $d_{\mathcal{D}_i}(v) = d_{\mathcal{D}_0}(v) + d_{\Lambda_i}(v) \geq \lfloor d_{\mathcal{D}}(v)/2 \rfloor$ ($i = 1, 2$). Thus we conclude from (2) that

$$(3) \quad d_{\mathcal{D}_1}(v) \geq \lceil d_{\mathcal{D}}(v)/2 \rceil \quad \text{and} \quad d_{\mathcal{D}_2}(v) \geq \lfloor d_{\mathcal{D}}(v)/2 \rfloor.$$

By the definition of \mathcal{D} we have $d_{\mathcal{D}}(v) = \mathbf{y}^T B_v$, where B_v is the column of B_G indexed by v . So

$$(4) \quad d_{\mathcal{D}}(v) + \mathbf{z}_v = \mathbf{p}_v + 2\mathbf{w}_v \geq 2\mathbf{w}_v.$$

Since \mathbf{w}_v is an integer, we deduce that

$$(5) \quad \mathbf{w}_v \leq \lfloor (d_{\mathcal{D}}(v) + \mathbf{z}_v)/2 \rfloor.$$

Setting $\mathbf{z}_{1v} = \lfloor \mathbf{z}_v/2 \rfloor$ and $\mathbf{z}_{2v} = \lceil \mathbf{z}_v/2 \rceil$, we have

$$(6) \quad \mathbf{z}_{1v} \mathbf{u}_v + \mathbf{z}_{2v} \mathbf{u}_v = \mathbf{z}_v \mathbf{u}_v.$$

We further claim that

$$(7) \quad d_{\mathcal{D}_i}(v) + \mathbf{z}_{iv} \geq \mathbf{w}_v, \quad \text{for } i = 1, 2.$$

To see (7), recall $\mathbf{p}_v \mathbf{z}_v = 0$ from (2). If $d_{\mathcal{D}}(v)$ is even, we deduce from (4) that \mathbf{z}_v is even, which implies, by (3-4), that $d_{\mathcal{D}_i}(v) + \mathbf{z}_{iv} \geq \frac{1}{2}(d_{\mathcal{D}}(v) + \mathbf{z}_v) \geq \mathbf{w}_v$. So we assume that $d_{\mathcal{D}}(v)$ is odd. If $\mathbf{z}_v = 0$ then, by (3) and (5), $d_{\mathcal{D}_i}(v) + \mathbf{z}_{iv} = d_{\mathcal{D}_i}(v) \geq \lfloor d_{\mathcal{D}}(v)/2 \rfloor \geq \mathbf{w}_v$. Else, by (2) and (4), \mathbf{z}_v is odd. Thus $d_{\mathcal{D}_i}(v) + \mathbf{z}_{iv} \geq \frac{1}{2}(d_{\mathcal{D}_i}(v) \pm 1) + \frac{1}{2}(\mathbf{z}_v \mp 1) = \frac{1}{2}(d_{\mathcal{D}_i}(v) + \mathbf{z}_v) \geq \mathbf{w}_v$, because of (3), (5), and the definition of \mathbf{z}_{iv} . So (7) holds.

For $i = 1, 2$, let $\mathbf{z}_i = (\mathbf{z}_{iv} : v \in V)$ and $\mathbf{y}_i \in \mathbb{Z}_+^{\mathcal{C}}$ be the multiplicity function of \mathcal{D}_i . It follows from (7) that $\mathbf{y}_i^T B_G + \mathbf{z}_i^T \geq \mathbf{w}^T$, which means that both $(\mathbf{y}_1, \mathbf{z}_1)$ and $(\mathbf{y}_2, \mathbf{z}_2)$ are feasible solutions of the second program in (3.1). From (1) and (6) we also conclude that $\mathbf{y}_i^T \mathbf{1} + \mathbf{z}_i^T \mathbf{u} \leq (\mathbf{y}^T \mathbf{1} + \mathbf{z}^T \mathbf{u})/2$ holds for at least one $i \in \{1, 2\}$. Hence inequality (3.1) holds, which proves the Theorem. \blacksquare

For a perfect graph G , being ESP can be characterized as follows. Let \mathbb{Z}_+ denote the set of nonnegative integers. For any $d \in \mathbb{Z}_+^{V(G)}$, let G^d denote the graph obtained from G by substituting each vertex v with a stable set of size $d(v)$. Note that v is deleted when $d(v) = 0$. Let $c_G = \mathbf{1}^T B_G$. In other words, for each $v \in V(G)$, $c_G(v)$ is the number of maximal cliques of G that contain v .

Theorem 3.8. *Let G be perfect. Then G is ESP if and only if for every $d \in \mathbb{Z}_+^{V(G)}$ with $d \leq c_G$ there exists $d' \in \mathbb{Z}_+^{V(G)}$ such that $\lfloor d/2 \rfloor \leq d' \leq \lceil d/2 \rceil$ and $\alpha(G^{d'}) + \alpha(G^{d-d'}) \leq \alpha(G^d)$.*

Proof. To prove the forward implication, let G be ESP and let $d \in \mathbb{Z}_+^{V(G)}$. Since G^d is perfect, its vertex set can be partitioned into $\alpha(G^d)$ cliques. These cliques naturally correspond to a multiset Λ of $\alpha(G^d)$ cliques of G . Note that $|\Lambda| = \alpha(G^d)$ and $d_\Lambda = d$. Since G is ESP, Λ admits an equitable subpartition (Λ_1, Λ_2) . By deleting vertices from cliques in Λ_1 and Λ_2 we can obtain multisets Λ_1^* and Λ_2^* of cliques of G such that $|\Lambda_1^*| + |\Lambda_2^*| \leq |\Lambda_1| + |\Lambda_2|$, $d_{\Lambda_1^*} + d_{\Lambda_2^*} = d$, and $\min\{d_{\Lambda_1^*}, d_{\Lambda_2^*}\} \geq \lfloor d/2 \rfloor$. Let $d' = d_{\Lambda_1^*}$. Then $\lfloor d/2 \rfloor \leq d' \leq \lceil d/2 \rceil$ and

$$\alpha(G^{d'}) + \alpha(G^{d-d'}) \leq \alpha(G^{d_{\Lambda_1}}) + \alpha(G^{d_{\Lambda_2}}) \leq |\Lambda_1| + |\Lambda_2| \leq |\Lambda| = \alpha(G^d),$$

which proves the forward implication.

To prove the backward implication, let Λ be a set of maximal cliques of G . Then $d := d_\Lambda \leq c_G$ and thus there exists d' as stated in the theorem. Let $d_1 = d'$ and $d_2 = d - d'$. For $i = 1, 2$, vertices of G^{d_i} can be partitioned into $\alpha(G^{d_i})$ cliques, and these cliques correspond to a multiset Λ_i of $\alpha(G^{d_i})$ cliques of G . Note that $d_{\Lambda_i} = d_i$. Thus (Λ_1, Λ_2) is an equitable subpartition of Λ , which proves the theorem. ■

We first remark that $\alpha(G^d)$ is exactly the maximum of $\sum_{v \in S} d(v)$ over all stable sets S of G . Sometimes this interpretation is more convenient. We also remark that we do not know a box-perfect graph that is not ESP. It seems reasonable to conjecture that no such a graph exists.

4 Known box-perfect graphs

Cameron [4] identified a few classes of box-perfect graphs. In this section we prove that they are in fact ESP graphs. Our results could be stronger than the results of Cameron if ESP and box-perfect are not equivalent. But the main reason for establishing our results is for future applications. We envision that more ESP graphs (possibly all box-perfect graphs) can be constructed from basic ESP graphs. Therefore, it is important to make sure that all known box-perfect graphs are ESP.

4.1 Totally unimodular graphs

It is well known (see Theorem 19.3 of [21]) that in a totally unimodular matrix, each set of rows can be partitioned so that the sum of one part minus the sum of the other part is a $\{0, \pm 1\}$ -vector. If G is totally unimodular then B_G has this partition property, which implies immediately that G satisfies the definition of ESP graphs. Thus we have the following.

Theorem 4.1. *Totally unimodular graphs are ESP.*

We point out that totally unimodular graphs include graphs like interval graphs, bipartite graphs, and block graphs (every block is a complete graph).

4.2 Incomparability graphs

Theorem 4.2. *Every incomparability graph G is ESP.*

Proof. Since G is perfect, we may apply Theorem 3.8. Let $d \in \mathbb{Z}_+^{V(G)}$. Note that G^d is again an incomparability graph. In fact, let P be a poset such that G is the incomparability of P and let P^d be obtained from P by replacing each element v with a chain of size $d(v)$. Then G^d is the incomparability graph of poset P^d . For each positive integer i , let A_i be the set of maximal elements of $P^d - (A_1 \cup \dots \cup A_{i-1})$. Then (A_1, \dots, A_n) is a partition of $V(G^d)$ into cliques, where $n = \alpha(G^d)$. Let V_1 be the union of A_i for all odd i and let V_2 be the union of A_i for all even i . Then $G^d[V_1]$ and $G^d[V_2]$ can be expressed as G^{d_1} and G^{d_2} , respectively, for some $d_1, d_2 \in \mathbb{Z}_+^{V(G)}$. It is easy to see that $d_1 + d_2 = d$ and $\lfloor d/2 \rfloor \leq d_j \leq \lceil d/2 \rceil$ ($j = 1, 2$). Moreover, each $\alpha(G^{d_j})$ is bounded by the number of A_i s contained in V_j . Therefore, $\alpha(G^{d_1}) + \alpha(G^{d_2}) \leq \alpha(G^d)$, which implies that $d' = d_1$ satisfies Theorem 3.8 and thus G is ESP. \blacksquare

4.3 p -Comparability graphs

p -Comparability graphs were introduced in [3] and were shown [3, 5] to be box-perfect. We show that they are ESP. Let D be a digraph with a special set T of vertices such that every arc is in a dicycle (directed cycle) and every dicycle meets T exactly once. In particular, D has no arc between any two vertices of T . If p is an integer with $|T| \leq p$, then a p -comparability graph G is defined from D by adding all chords of all dicycles, then deleting T , and finally ignoring all directions on edges. Note that 1-comparability graphs are precisely comparability graphs.

Theorem 4.3. *Every p -comparability graph G is ESP.*

To prove this theorem we will need the following Lemma. Let $D = (V, A)$ be a digraph. For each dicycle C of D , the *incidence vector* of C is the vector $\chi^C \in \{0, 1\}^A$ such that $\chi^C(a) = 1$ if and only if a is on C . A sum of incidence vectors of (not necessarily distinct) dicycles of D is called a *circulation* of D . The following is a special case of Corollary 11.2b of [22].

Lemma 4.4. *Every circulation f is the sum of two circulations f_1, f_2 such that $\lfloor f/2 \rfloor \leq f_i \leq \lceil f/2 \rceil$ holds for both $i = 1, 2$.*

Proof of Theorem 4.3. Let G be constructed from D and T . Let D^* be obtained from D by splitting each vertex v into v' and v'' such that arcs entering v are now entering v' , and arcs leaving v are now leaving v'' . We also add an arc from v' to v'' . Observe that for every dicycle C of D , D^* has a unique dicycle C^* such that $A(C^*) \cap A(D) = A(C)$. Moreover, every dicycle of D^* can be expressed as C^* for a dicycle C of D .

We will use a fact proved in [5] that for every clique K of G , there exists a dicycle C_K of D such that $K \subseteq V(C_K)$.

Let Λ be a set of maximal cliques of G . We prove the theorem by showing that Λ admits an equitable subpartition. Let f be the sum of incidence vectors of C_K^* over all $K \in \Lambda$. Since each C_K meets T exactly once, each C_K^* must meet $T^* = \{t't'' : t \in T\}$ exactly once. As a result, $|\Lambda|$ equals

the sum of $f(a)$ over all $a \in T^*$. In addition, since each $K \in \Lambda$ is a maximal clique, we must have $V(C_K) - T = K$. This implies that $d_\Lambda(v) = f(v'v'')$ holds for all $v \in V(D)$.

Let f_1 and f_2 be the two circulations of D^* determined by Lemma 4.4. For $i = 1, 2$, let \mathcal{C}_i^* be the multiset of dicycles of D^* such that f_i is the sum of χ^{C^*} over all $C^* \in \mathcal{C}_i^*$. Then let \mathcal{C}_i be the multiset $\{C : C^* \in \mathcal{C}_i^*\}$ and $\Lambda_i = \{V(C) - T : C \in \mathcal{C}_i\}$. By the construction of G , each member of Λ_i is a clique of G . Moreover, $d_{\Lambda_i}(v) = f_i(v'v')$ holds for all $v \in V(G)$, and $|\Lambda_i| = \sum_{a \in T^*} f_i(a)$. Therefore, (Λ_1, Λ_2) is an equitable subpartition of Λ , which proves that G is ESP. \blacksquare

Remark. Let us call a graph *strong ESP* if every set Λ of maximal cliques admits an equitable subpartition (Λ_1, Λ_2) with $\max\{|\Lambda_1|, |\Lambda_2|\} \leq \lceil |\Lambda|/2 \rceil$. This proof also proves that (1-)comparability graphs are in fact strong ESP.

5 Parity graphs

A graph is called a *parity graph* if any two induced paths between the same pair of vertices have the same parity. These are natural extensions of bipartite graphs and they are perfect [19]. Cameron and Edmonds [3] conjectured that every parity graph is box-perfect. The objective of this section is to present a proof of this conjecture.

To establish our result we need a structural characterization of parity graphs. Let H be a graph with a stable set S such that all vertices of S have the same set of neighbors. Let B be a bipartite graph and let T be a subset of a color class of B with $|T| = |S|$. Let G be obtained from the disjoint union of H and B by identifying S with T . We call G a *bipartite extension* of H by B , and we also call the construction of G from H *bipartite extension*.

Lemma 5.1 (Burlet and Uhry [2]). *Every connected parity graph can be constructed from a single vertex by repeatedly duplicating vertices and bipartite extensions.*

Lemma 5.2. *Duplicating a vertex in an ESP graph results in an ESP graph.*

Proof. Let ESP graph G have a vertex v . Let G' be obtained by duplicating v and let v' be the new vertex. For any set Λ' of maximal cliques of G' , we prove that Λ' has an equitable subpartition.

We define Λ as follows. If vv' is an edge then $\Lambda = \{K - v' : K \in \Lambda'\}$; if vv' is not an edge then $\Lambda = \{K : v' \notin K \in \Lambda'\} \uplus \{K - v' + v : v' \in K \in \Lambda'\}$. Note that Λ is a multiset of maximal cliques of G . Since G is ESP, Λ admits an equitable subpartition (Λ_1, Λ_2) . By deleting vertices from cliques in Λ_1 and Λ_2 we may assume that $d_{\Lambda_1} + d_{\Lambda_2} = d_\Lambda$ and $\lfloor d_\Lambda/2 \rfloor \leq d_{\Lambda_i} \leq \lceil d_\Lambda/2 \rceil$ ($i = 1, 2$).

If vv' is an edge, let $\Lambda'_i = \{K : v \notin K \in \Lambda_i\} \uplus \{K + v' : v \in K \in \Lambda_i\}$ ($i = 1, 2$). Then (Λ'_1, Λ'_2) is an equitable subpartition of Λ' because $d_X(v') = d_X(v)$ holds for $X \in \{\Lambda, \Lambda'_1, \Lambda'_2\}$.

Now suppose vv' is not an edge. Note that $d_\Lambda(v) = d_{\Lambda'}(v) + d_{\Lambda'}(v')$. Also we may assume that $d_{\Lambda_1}(v) = \lfloor d_\Lambda(v)/2 \rfloor$ and $d_{\Lambda_2}(v) = \lceil d_\Lambda(v)/2 \rceil$. Let

$$m_1 = \lfloor d_{\Lambda'}(v)/2 \rfloor, \quad m_2 = \lceil d_{\Lambda'}(v)/2 \rceil, \quad m'_1 = d_{\Lambda_1}(v) - m_1, \quad m'_2 = d_{\Lambda_2}(v) - m_2.$$

Then

$$m_1 + m_2 = d_{\Lambda'}(v), \quad m'_1 + m'_2 = d_{\Lambda'}(v'), \quad \min\{m'_1, m'_2\} \geq \lfloor d_{\Lambda'}(v')/2 \rfloor.$$

For $i = 1, 2$, let Λ'_i be obtained from Λ_i by turning m'_i cliques K that contain v into $K - v + v'$. Then the above equalities and inequalities imply that (Λ'_1, Λ'_2) is an equitable subpartition of Λ' . ■

Remark. Clearly, this proof also proves that duplicating a vertex in a strong ESP graph results in a strong ESP graph.

Theorem 5.3. *Parity graphs are ESP.*

Proof. By Lemma 5.2, we only need to show that if G is a bipartite extension of an ESP graph H by a bipartite graph $B = (X, Y, E)$, then G is ESP. Let $X_0 \subseteq X$ be the intersection of H and B . Let Λ be a set of maximal cliques of G . Naturally, Λ can be partitioned into Λ_H and Λ_B , which are maximal cliques of H and edges of B , respectively. Now we find an equitable subpartition $(\Lambda'_B, \Lambda''_B)$ of Λ_B and an equitable subpartition $(\Lambda'_H, \Lambda''_H)$ of Λ_H such that $(\Lambda'_B \cup \Lambda'_H, \Lambda''_B \cup \Lambda''_H)$ is an equitable subpartition of Λ . Let X_0 be partitioned into (X_1, X_2) such that X_1 consists of $x \in X_0$ with both $d_{\Lambda_B}(x)$ and $d_{\Lambda_H}(x)$ odd. Since $(\Lambda'_B, \Lambda''_B)$ and $(\Lambda'_H, \Lambda''_H)$ are always compatible on vertices in X_2 , we only need to focus on vertices in X_1 .

Without loss of generality, let $\Lambda_B = E$. Suppose B has $2t$ vertices of odd degree. Then E can be partitioned into cycles and t paths P_1, \dots, P_t . Let $(\Lambda'_B, \Lambda''_B)$ be defined by assigning edges to the two parts alternatively along the cycles and paths. Then $(\Lambda'_B, \Lambda''_B)$ is an equitable partition. Note that we have the following freedom in the assignment. Let $x \in X_1$ and let P_i be the path with x as an end. If the other end of P_i is not in X_1 , then we may choose $d_{\Lambda'_B}(x)$ to be $\lfloor d_{\Lambda_B}(x)/2 \rfloor$ or $\lceil d_{\Lambda_B}(x)/2 \rceil$, as we wish (without changing $d_{\Lambda'_B}(z)$ and $d_{\Lambda''_B}(z)$ for any other $z \in X_1$). If the other end of P_i is a vertex x' in X_1 , then we may assume that $d_{\Lambda'_B}(x) = \lfloor d_{\Lambda_B}(x)/2 \rfloor$ and $d_{\Lambda'_B}(x') = \lceil d_{\Lambda_B}(x')/2 \rceil$. Let $(x_1, x'_1), \dots, (x_k, x'_k)$ be these pairs in X_1 .

Let H_1 be obtained from H by deleting x'_1, \dots, x'_k and let Λ_1 be obtained from Λ_H by replacing each x'_i with x_i . Note that $d_{\Lambda_1}(x_i) = d_{\Lambda_H}(x_i) + d_{\Lambda_H}(x'_i)$ for all i , while $d_{\Lambda_1}(v) = d_{\Lambda_H}(v)$ for all other vertices v of H_1 . Since H is ESP, so is H_1 . Let $(\Lambda'_1, \Lambda''_1)$ be an equitable subpartition of Λ_1 . Without loss of generality, we assume $d_{\Lambda'_1}(x_i) = d_{\Lambda''_1}(x_i) = d_{\Lambda_1}(x_i)/2$ for all i . Let Λ'_H be obtained from Λ'_1 by turning $\lfloor d_{\Lambda_H}(x'_i)/2 \rfloor$ of its cliques K that contain x_i into $K - x_i + x'_i$ (for every i). Then $d_{\Lambda'_H}(x_i) = \lceil d_{\Lambda_H}(x_i)/2 \rceil$ and $d_{\Lambda'_H}(x'_i) = \lfloor d_{\Lambda_H}(x'_i)/2 \rfloor$. Let Λ''_H be obtained analogously. Now it is straightforward to verify that, the freedom on partition $(\Lambda'_B, \Lambda''_B)$ allows us to make adjustments so that $(\Lambda'_B \cup \Lambda'_H, \Lambda''_B \cup \Lambda''_H)$ is an equitable subpartition of Λ . ■

6 Complements of line graphs

In the rest of this paper we allow some graphs to have loops and parallel edges. We call these *multigraphs* and we reserve the word *graph* for simple graphs. If a multigraph H is obtained from a graph H_0 by adding loops and parallel edges, then H_0 is called a *simplification* of H and is denoted by $si(H)$.

Let $L(H)$ denote the line graph of a multigraph H . Under this circumstance, we always make the following implicit assumptions:

- (i) H has no isolated vertices (deleting an isolated vertex does not affect $L(H)$);
- (ii) H has no loops (replacing a loop with a pendent edge does not affect $L(H)$);

- (iii) H has no distinct vertices x, y, z such that z is the only neighbor of x and the only neighbor of y (replacing edges between y and z by edges between x and z does not affect $L(H)$).

The complement of $L(H)$ will be denoted by $\bar{L}(H)$. Our results in the next two sections imply a characterization of box-perfect line graphs. The goal of this section is to characterize box-perfect graphs that are complements of line graphs.

Theorem 6.1. *Let $G = \bar{L}(H)$ be perfect. Then G is box-perfect if and only if G is $\{S_3, \bar{S}_3^+\}$ -free.*

Our proof of this theorem is divided into a sequence of lemmas. We first determine the structure of $\{S_3, \bar{S}_3^+\}$ -free perfect graphs of the form $\bar{L}(H)$, and then we confirm that all such graphs are ESP. We will see that some of these graphs are in fact strong ESP.

We need a result of Gallai [15] which identifies eight classes and ten individual graphs such that a graph is a comparability graph if and only if it does not contain any of these identified graphs as an induced subgraph. We will use the following immediate consequence of Gallai's theorem. Let Γ be the graph obtained from a 6-cycle $v_1v_2v_3v_4v_5v_6v_1$ by adding two edges v_1v_3 and v_1v_5 .

Lemma 6.2. *Let G be claw-free and perfect. Then G is an incomparability graph if and only if G does not contain any of S_3, \bar{S}_3, Γ , and C_{2n} ($n \geq 3$) as an induced subgraph.*

Let K_4^+ denote the graph obtained from K_4 by adding two pendent edges to two of its distinct vertices. Let $K_{2,n}^+$ denote the graph obtained from $K_{2,n}$ ($n \geq 3$) by adding a pendent edge to a degree-2 vertex and an edge between the two degree- n vertices.

Lemma 6.3. *Let $\bar{L}(H)$ be $\{C_5, S_3, \bar{S}_3^+\}$ -free. If H contains \bar{S}_3 as a subgraph then $si(H)$ is either K_4^+ or a subgraph of $K_{2,n}^+$ for some $n \geq 3$.*

Proof. Since \bar{S}_3 is a subgraph of H , we assume $V(H) = \{x_1, x_2, x_3, y_1, y_2, y_3, z_1, \dots, z_m\}$ such that $x_1x_2x_3$ is a triangle and $x_iy_i \in E(H)$ ($i = 1, 2, 3$). If $m = 0$ then it is straightforward to verify the conclusion of the lemma, using the fact that H does not contain C_5 as a subgraph. So we assume $m > 0$. Let $K_{1,3}^*$ denote the graph obtained from $K_{1,3}$ by subdividing each edge exactly once. Note that $K_{1,3}^*$ is not a subgraph of H since $\bar{L}(K_{1,3}^*) = S_3$. As a result, each z_i is adjacent to none of y_1, y_2, y_3 , and at most two of x_1, x_2, x_3 . Furthermore, since $\bar{L}(H)$ is \bar{S}_3^+ -free, the entire neighborhood of each z_i must be a subset of $\{x_1, x_2, x_3\}$ of size one or two (here we also use assumption (i) above). By assumption (iii) above we may assume that each z_i is adjacent to exactly two of x_1, x_2, x_3 . Since C_5 is not a subgraph of H , all z_i 's must have the same set of neighborhood. Now, since $m > 0$, it is straightforward to verify that $si(H)$ is a subgraph of $K_{2,m+3}^+$. \blacksquare

Let C be an even cycle of length ≥ 4 . Let X be a stable set of C and let $Y = V(C) - X - N_C(X)$, where X is allowed to be empty. We construct a bipartite graph from C by adding a pendent edge to each vertex in Y and by repeatedly duplicating vertices in X . Let \mathcal{C} consist of all graphs that can be constructed in this way.

Lemma 6.4. *Let $L(H)$ be perfect and \bar{S}_3 -free. Suppose H is connected and H does not contain \bar{S}_3 as a subgraph. If $L(H)$ contains an induced Γ or C_{2n} ($n \geq 3$), then $si(H)$ is a subgraph of a graph in $\mathcal{C} \cup \{K_{3,3}\}$.*

Proof. Suppose Γ is an induced subgraph of $L(H)$. Then H has a subgraph with a 4-cycle $x_1x_2x_3x_4$ and two pendent edges x_1y_1, x_2y_2 . Note that x_1x_3 and x_2x_4 are not edges of H since \bar{S}_3 is not

a subgraph of H . Let z_1, \dots, z_m be the remaining vertices of H . If $m = 0$, then either $si(H)$ is a subgraph of $K_{3,3}$ or H contains a 5-cycle. So we assume $m > 0$. Like in the proof of the last lemma, since C_5 and $K_{1,3}^*$ are not subgraphs of H , for each i we must have $N_H(z_i) = \{x_1, x_3\}$ or $\{x_2, x_4\}$, or $\{x_j\}$ for some j . In addition, $N_H(y_i) \subseteq \{x_i, x_{i+2}\}$ ($i = 1, 2$) and $|N_H(y_1) \cup N_H(y_2)| \leq 3$. Now, since H does not contain $K_{1,3}^*$, it is routine to check that $si(H)$ is a subgraph of a graph in \mathcal{C} .

Next, suppose $L(H)$ is Γ -free. Then H contains a $2n$ -cycle $x_1x_2\dots x_{2n}$ ($n \geq 3$). Note that this cycle has no chord (otherwise $L(H)$ contains an induced Γ , \bar{S}_3 , or C_{2k+1} with $k \geq 2$). Let z_1, \dots, z_m be the remaining vertices of H . Using the same argument we used in the last paragraph it is straightforward to show that each $N_H(z_i)$ is $\{x_j\}$ or $\{x_j, x_{j+2}\}$ for some j (where x_{2n+t} is x_t). In addition, if $N_H(z_i) = \{x_j, x_{j+2}\}$ then $N_H(x_{j+1}) = N_H(z_i)$. Therefore, $si(H)$ is a subgraph of a graph in \mathcal{C} . ■

Lemma 6.5. *Suppose G has a vertex u such that $G - u$ is bipartite and $G - N(u)$ is edge-less. Then G is totally unimodular.*

Proof. By Theorem 19.3 of [21], we only need to show that each set Λ of maximal cliques admits an *equitable* partition (Λ_1, Λ_2) , meaning that $\min\{d_{\Lambda_1}(v), d_{\Lambda_2}(v)\} \geq \lfloor d_\Lambda(v) \rfloor$, for all $v \in V(G)$. Suppose to the contrary that some Λ does not admit such a partition. We choose Λ with $|\Lambda|$ as small as possible.

Let A, B, C, D be a partition of $V(G) - u$ such that $A \cup C, B \cup D$ are stable and $N(u) = B \cup C$. Let G' be the subgraph of G formed by edges in $K - u$, over all $K \in \Lambda$. We claim that G' is a forest. Suppose G' has a cycle $x_1x_2\dots x_n$. Note that for each i , exactly one of $x_i x_{i+1}$ and $u x_i x_{i+1}$ is a clique in Λ . Let Λ' be the rest cliques in Λ . By the minimality of $|\Lambda|$, Λ' admits an equitable partition (Λ'_1, Λ'_2) . Let us extend Λ'_j ($j = 1, 2$) to Λ_j by including $x_i x_{i+1}$ or $u x_i x_{i+1}$ (whichever belongs to Λ) for all i with $i - j$ even. Then it is easy to see that (Λ_1, Λ_2) is an equitable partition of Λ . This contradicts the choice of Λ and thus the claim is proved. The same argument also shows that G' has no maximal path with two ends both in $A \cup B$ or both in $C \cup D$. Thus all components of G' are paths with one end in $A \cup B$ and one end in $C \cup D$. If G' has only one path then the same argument still works. If G' has two or more paths then we can take any two of them and treat their union as a cycle and again apply the same argument. ■

Recall that a graph G is *strong ESP* if every set Λ of maximal cliques of G admits an equitable subpartition (Λ_1, Λ_2) with $\max\{|\Lambda_1|, |\Lambda_2|\} \leq \lceil |\Lambda|/2 \rceil$. The next lemma follows immediately from this definition.

Lemma 6.6. (1) *If G is strong ESP then so are all its induced subgraphs.*

(2) *Let G_1, G_2 be strong ESP and let G be obtained from the disjoint union of G_1, G_2 by adding all edges between them. Then G is also strong ESP.*

In a (loopless) multigraph G , the *degree* of a vertex v , denoted $d_G(v)$, is the number of edges incident with v . The next is the key step for proving Theorem 6.1.

Lemma 6.7. *For every $H \in \mathcal{C} \cup \{K_{3,3}\}$, $\bar{L}(H)$ is strong ESP.*

Proof. For each $\mu \in \mathbb{Z}_+^{E(H)}$, let μH denote the multigraph with vertex set $V(H)$ such that the number of edges between any two vertices x, y is zero (if $xy \notin E(H)$) or $\mu(xy)$ (if $xy \in E(H)$). Note that μH is bipartite since H is bipartite. Let $\Delta(\mu)$ denote the maximum degree of μH . By König's

edge-coloring theorem, $E(\mu H)$ is the union of k matchings if and only if $k \geq \Delta(\mu)$. Because of this theorem and the one-to-one correspondence between cliques of $\bar{L}(H)$ and matchings of H , to prove the lemma it is enough for us to show that

- (*) for any $\mu \in \mathbb{Z}_+^{E(H)}$ there exist $\mu_1, \mu_2 \in \mathbb{Z}_+^{E(H)}$ such that $\mu_1 + \mu_2 = \mu$, $\mu_i \geq \lfloor \mu/2 \rfloor$ ($i = 1, 2$), $\Delta(\mu_1) \leq \lceil \Delta(\mu)/2 \rceil$, and $\Delta(\mu_2) \leq \lfloor \Delta(\mu)/2 \rfloor$.

In the following we construct a partition (E_1, E_2) of $E(\mu H)$ such that the multiplicity functions μ_i of E_i ($i = 1, 2$) satisfies (*). This partition will be constructed in several steps. In the process we determine a partition (E_1, E_2, E_3) of $E(\mu H)$, where we begin with $(E_1, E_2, E_3) = (\emptyset, \emptyset, E(\mu(H)))$ and we keep moving edges from E_3 to E_1, E_2 until E_3 becomes empty. For $i = 1, 2, 3$, let H_i denote the subgraph of μH formed by edges in E_i .

First, for each edge $e = xy$ of H , among all $\mu(e)$ edges of E_3 that are between x and y , we move $\lfloor \mu(e)/2 \rfloor$ of them to E_1 and $\lfloor \mu(e)/2 \rfloor$ of them to E_2 . At the end of this process, H_3 becomes a simple graph. It follows that $\mu_i \geq \lfloor \mu/2 \rfloor$ ($i = 1, 2$) and this inequality will be satisfied no matter how edges of H_3 are moved to E_1 and E_2 in later steps.

If H_3 has a cycle C , since H is bipartite, $E(C)$ can be partitioned into two matchings M_1, M_2 . We move M_i from E_3 to E_i ($i = 1, 2$). We repeat this process until H_3 become a forest. At this point, H_1 and H_2 have the same degree on every vertex.

Let $S = \{v : d_{\mu H}(v) = \Delta(\mu)\}$. Suppose H_3 has a leaf v that is not in S . Let P be a maximal path of H_3 starting from v . Let $E(P)$ be partitioned into two matchings M_1, M_2 , where we assume the edge of P that is incident with the other end u of P belongs to M_1 . Then we move M_i from E_3 to E_i ($i = 1, 2$). After this change, $d_{H_1}(u) = \lceil d_{\mu H}(u)/2 \rceil \leq \lceil \Delta(\mu)/2 \rceil$, $d_{H_2}(u) = \lfloor d_{\mu H}(u)/2 \rfloor \leq \lfloor \Delta(\mu)/2 \rfloor$, and $d_{H_i}(v) \leq \lceil d_{\mu H}(v)/2 \rceil \leq \lfloor \Delta(\mu)/2 \rfloor$ ($i = 1, 2$). In addition, $d_{H_1}(w) = d_{H_2}(w)$ for all $w \neq u, v$, and $d_{H_i}(u), d_{H_i}(v)$ will remain unchanged in the remaining process. By repeating this process we may assume that all leaves of H_3 are in S . As a consequence, $\Delta(\mu)$ is odd. Note that the same argument works if H_3 has a maximal path with an odd number of edges. Thus we further assume that in every component of H_3 , all leaves are in the same color class (of any 2-coloring of H_3).

We first consider the case $H = K_{3,3}$. We claim that each component of H_3 is a path. Suppose a component H'_3 of H_3 is not a path. Then H'_3 has at least three leaves. Since all these leaves are in the same color class, H'_3 must have exactly three leaves z_1, z_2, z_3 and they form a color class of H . Consequently, $H'_3 = H_3 = K_{1,3}$. Moreover, in the previous steps of reducing H_3 , no path was ever deleted because otherwise H_3 would be a subgraph of $K_{2,2}$. It follows that $d_{\mu H}(v^*)$ is even, where $v^* \in V(H) - V(H_3)$. However, the fact $z_1, z_2, z_3 \in S$ implies that μH is $\Delta(\mu)$ -regular, and thus $d_{\mu H}(v^*) = \Delta(\mu)$ is odd. This contradiction proves our claim. Now, since each non-leaf v of H_3 has degree two, its degree in μH is even and thus $v \notin S$. It follows that moving all edges of E_3 to E_1 results in the required partition.

Next suppose $H \in \mathcal{C}$. Let H'_3 be a component of H_3 . Then H'_3 is a *caterpillar* since $K_{1,3}^*$ is not a subgraph of H . Therefore, H'_3 has a path $x_1x_2\dots x_{2k+1}$ such that every leaf of H'_3 is adjacent to some x_{2i+1} . We assume that H'_3 is not a path because otherwise we may move the entire path from E_3 to E_1 . We make two observations before we continue. First, $d_H(v) > 1$ holds for every leaf v of H'_3 , because otherwise the only edge of H that is incident with v would be the only edge of H'_3 (as $v \in S$). Second, if $u, v \in V(H'_3)$ are of degree-2 in H and are contained in a 4-cycle $uxvy$ of H , then at most one of u, v is in S . This is because otherwise $\mu(ux) = \mu(vy)$, $\mu(uy) = \mu(vx)$, and

both $x, y \in S$, which implies that H'_3 is a subgraph of the 4-cycle $uxvy$. It follows from these two observations and the construction of graphs in \mathcal{C} that each x_{2i+1} is adjacent to at most two leaves of H'_3 . For the same reasons, there must exist $i_0 \in \{0, 1, \dots, k\}$ such that $d_{H'_3}(x_{2i_0+1}) = 2$.

For $i = 1, 2, 3, 4$, let $V_i = \{v : d_{H'_3}(v) = i\}$. Note that $V_2 \cup V_3 \cup V_4 = \{x_1, \dots, x_{2k+1}\}$. Let M be the matching $\{x_{2i-1}x_{2i} : i = 1, \dots, i_0\} \cup \{x_{2i}x_{2i+1} : i_0 + 1, \dots, k\}$. From H'_3 we move M to E_2 and the rest of $E(H'_3)$ to E_1 . Now we verify that, after this change, $d_{H_1}(v) \leq \lceil \Delta(\mu)/2 \rceil$ and $d_{H_2}(v) \leq \lfloor \Delta(\mu)/2 \rfloor$ hold for all $v \in V_1 \cup V_2 \cup V_3 \cup V_4$. For each $v \in V_1$ it is easy to see that in fact $d_{H_1}(v) = \lceil \Delta(\mu)/2 \rceil$ and $d_{H_2}(v) = \lfloor \Delta(\mu)/2 \rfloor$. For each even i , we have $x_i \in V_2$ and $d_{H_1}(x_i) = d_{H_2}(x_i) = d_{\mu H}(x_i)/2 \leq \lfloor \Delta(\mu)/2 \rfloor$. For each odd i we consider two cases. If $x_i \in V_3$ then $d_{H_1}(x_i) = (d_{\mu H}(x_i) + 1)/2 \leq \lceil \Delta(\mu)/2 \rceil$ and $d_{H_2}(x_i) = (d_{\mu H}(x_i) - 1)/2 \leq \lfloor \Delta(\mu)/2 \rfloor$. If $x_i \in V_2 \cup V_4$ then $d_{H_1}(x_i) \leq (d_{\mu H}(x_i) + 2)/2 \leq \lceil \Delta(\mu)/2 \rceil$ and $d_{H_2}(x_i) \leq d_{\mu H}(x_i)/2 \leq \lfloor \Delta(\mu)/2 \rfloor$. Therefore, we may apply this split to all components of H_3 and create the required partition E_1, E_2 . ■

Proof of Theorem 6.1. The forward implication is obvious so we only show that $G = \bar{L}(H)$ is ESP when G is perfect and $\{S_3, \bar{S}_3^+\}$ -free.

Suppose $L(H)$ contains an induced S_3 . Then H contains \bar{S}_3 as a subgraph. By Lemma 6.3, $si(H)$ is either K_4^+ or a subgraph of $K_{2,n}^+$ for some $n \geq 3$. In both cases, it is straightforward to verify that $\bar{L}(si(H))$ satisfies the assumptions in Lemma 6.5. So $\bar{L}(si(H))$ is totally unimodular and thus is also ESP. By Lemma 5.2, $\bar{L}(H)$ is ESP.

Now suppose $L(H)$ is S_3 -free. We claim that $\bar{L}(H')$ is strong ESP for every component H' of H . If $\bar{L}(H')$ is a comparability graph, then the claim follows immediately from the Remark at the end of Section 4. So we assume that $\bar{L}(H')$ is not a comparability graph. By Lemma 6.2, $L(H)$ contains an induced Γ or C_{2n} ($n \geq 3$). This implies, by Lemma 6.4, that $si(H')$ is a subgraph of a graph in $\mathcal{C} \cup \{K_{3,3}\}$. Then the claim follows from Lemma 6.7, Lemma 6.6(1), and the Remark of Lemma 5.2. Finally, this claim and Lemma 6.6(2) imply that $\bar{L}(H)$ is ESP. ■

7 Trigraphs

Our next objective is to characterize claw-free box-perfect graphs. To accomplish this goal, we will need a result of Chudnovsky and Plumettaz [9] on the structure of claw-free perfect graphs. The purpose of this section is to explain their result, which requires many definitions.

A *trigraph* G consists of a finite set V of *vertices* and an *adjacency function* $\theta : \binom{V}{2} \rightarrow \{1, 0, -1\}$ such that $\{uv : \theta(uv) = 0\}$ is a matching. Two distinct vertices u and v of G are *strongly adjacent* if $\theta(uv) = 1$, *strongly antiadjacent* if $\theta(uv) = -1$, and *semiadjacent* if $\theta(uv) = 0$. We call u, v *adjacent* if $\theta(uv) \geq 0$, and *antiadjacent* if $\theta(uv) \leq 0$. Note that every graph can be considered as a trigraph with $\{uv : \theta(uv) = 0\} = \emptyset$. In other words, graphs are exactly trigraphs with no semiadjacent pairs. The result of Chudnovsky and Plumettaz is in fact about trigraphs.

For any trigraph $G = (V, \theta)$, let $G^{\geq 0}$ denote the graph $(V, \{uv : \theta(uv) \geq 0\})$. Conversely, for any graph $G = (V, E)$, let $tri(G)$ denote the set of all trigraphs (V, θ) such that for any distinct $u, v \in V$, $\theta(uv) \geq 0$ if $uv \in E$ and $\theta(uv) \leq 0$ if $uv \notin E$.

Let $G = (V, \theta)$ be a trigraph. We call G *connected* if $G^{\geq 0}$ is connected. For each $v \in V$, let

$N_G(v) = N_{G \geq 0}(v)$. We often write $N(v)$ for $N_G(v)$ if the dependency on G is clear. For any $X \subseteq V$, let $G|X$ be the trigraph such that its vertex set is X and its adjacency function is the restriction of θ to $\binom{X}{2}$. If a trigraph H is isomorphic to $G|X$ for some $X \subseteq V$, then we call H a *subtrigraph* of G and we say that G *contains* H .

A trigraph is a *hole* if it belongs to $\text{tri}(C_n)$ for some $n \geq 4$. A trigraph (V, θ) is an *antihole* if $(V, -\theta)$ is a hole. A hole or antihole is *odd* if its number of vertices is odd. A trigraph is *Berge* if it contains neither odd hole nor odd antihole. A trigraph is a *claw* if it belongs to $\text{tri}(K_{1,3})$. A trigraph is *claw-free* if it does not contain any claw. In general, if \mathcal{H} is a set of trigraphs, then a trigraph is *\mathcal{H} -free* if it does not contain any trigraph in \mathcal{H} . The result of Chudnovsky and Plumettaz characterizes {claw, holes, antiholes}-free trigraphs, that is, claw-free Berge trigraphs. To describe the resulting structure we need more definitions.

Let $G = (V, \theta)$ be a trigraph. For any two disjoint $X, Y \subseteq V$, we say that X is *complete* (resp. *strongly complete*, *anticomplete*, *strongly anticomplete*) to Y if every $x \in X$ and every $y \in Y$ are adjacent (resp. strongly adjacent, antiadjacent, strongly antiadjacent). A *clique* (resp. *strong clique*) of G is a set $C \subseteq V$ such that any two distinct vertices of C are adjacent (resp. strongly adjacent). A *stable set* (resp. *strong stable set*) of G is a set $S \subseteq V$ such that any two distinct vertices of S are antiadjacent (resp. strongly antiadjacent).

A trigraph H is a *thickening* of a trigraph G if $V(H)$ admits a partition $(X_v : v \in V(G))$ such that

- if $v \in V(G)$ then $X_v \neq \emptyset$ is a strong clique of H ;
- if $u, v \in V(G)$ are strongly adjacent in G then X_u is strongly complete to X_v in H ;
- if $u, v \in V(G)$ are strongly antiadjacent, then X_u is strongly anticomplete to X_v in H ;
- if $u, v \in V(G)$ are semiadjacent, then X_u is neither strongly complete nor strongly anticomplete to X_v in H .

Let \mathcal{C} be the class of all trigraphs illustrated in Figure 7.1, where

- $|B_i^j| \leq 1$ for all $i, j \in \{1, 2, 3\}$
- $|B_2^1 \cup B_3^1|, |B_1^2 \cup B_3^2|, |B_1^3 \cup B_2^3| \in \{0, 2\}$
- if $\theta(a_1 a_3) = 0$ then $B_2^1 \cup B_3^1 = \emptyset$
- there exists $x_i \in B_i^1 \cup B_i^2 \cup B_i^3$ for $i = 1, 2, 3$, such that $\{x_1, x_2, x_3\}$ is a clique.

It turns out that there are two kinds of claw-free Berge trigraphs. The first are thickenings of trigraphs in \mathcal{C} . The second are constructed (in a way like constructing line graphs) from certain basic trigraphs. In the following, we first define the building blocks and then describe the construction.

Let G have three vertices v, z_1, z_2 such that $\theta(vz_1) = \theta(vz_2) = 1$ and $\theta(z_1 z_2) = -1$. Then the pair $(G, \{z_1, z_2\})$ is a *spot*. Let G have four vertices v_1, v_2, z_1, z_2 such that $\theta(v_1 z_1) = \theta(v_2 z_2) = 1$, $\theta(v_1 v_2) = 0$, $\theta(z_1 z_2) = \theta(z_1 v_2) = \theta(z_2 v_1) = -1$. Then the pair $(G, \{z_1, z_2\})$ is a *spring*.

A trigraph is a *linear interval* if its vertices can be ordered as v_1, \dots, v_n such that if $i < j < k$ and $\theta(v_i v_k) \geq 0$ then $\theta(v_i v_j) = \theta(v_j v_k) = 1$. Let G be such a trigraph with $n \geq 4$. We call $(G, \{v_1, v_n\})$ a *linear interval stripe* if: v_1 and v_n are strongly antiadjacent, v_i and v_{i+1} are adjacent for every $i \in \{1, \dots, n-1\}$, no vertex is complete to $\{v_1, v_n\}$, and no vertex is semiadjacent to v_1 or v_n .

Let $(G, \{p, q\})$ be a spring or a linear interval strip. Let H be a thickening of G and let X_v ($v \in V(G)$) be the corresponding sets. If $|X_p| = |X_q| = 1$, then $(H, X_p \cup X_q)$ is called a *thickening*




Figure 7.1: Trigraphs in \mathcal{C}

of $(G, \{p, q\})$.

Let \mathcal{C}' be the class of all pairs $(H, \{z\})$ such that H is a thickening of a trigraph $G \in \mathcal{C}$ and $z \in X_{a_i}$ for some $i \in \{1, 2, 3\}$ for which $B_{i+1}^{i+2} \cup B_i^{i+2} = \emptyset$ and $N(z) \cap (X_{a_{i+1}} \cup X_{a_{i+2}}) = \emptyset$ (here we use the notation from the definitions of \mathcal{C} and thickening).

A *signed graph* (G, s) consists of a multigraph $G = (V, E)$ and a function $s : E \rightarrow \{0, 1\}$. If $\sum_{e \in E(C)} s(e)$ is even for all cycles C of G , then (G, s) is an *evenly signed graph*. In the following we define another three classes of signed graphs. For any $F \subseteq E$, let $G[F] = (V, F)$.

Let \mathcal{F}_1 be the class of loopless signed graphs (G, s) such that $si(G) = K_4$ and $s \equiv 1$. Let \mathcal{F}_2 be the class of loopless signed graphs (G, s) such that $si(G)$ is obtained from $K_{2,n}$ ($n \geq 1$) by adding an edge e^* between its two degree- n vertices, and edges in $\{e : s(e) = 0\}$ are all parallel to e^* (while $s(e^*) = 1$). We remark that our \mathcal{F}_1 is \mathcal{F}_2 of [9] and our \mathcal{F}_2 is $\mathcal{F}_1 \cup \mathcal{F}_3$ of [9]

In a connected multigraph G with $E(G) \neq \emptyset$, a subgraph B is a *block* of G if B is a loop or B is maximal with the property that B is loopless and $si(B)$ is a block of $si(G)$. A signed graph (G, s) is called an *even structure* if $E(G) \neq \emptyset$ and for all blocks B of G , $(B, s|_{E(B)})$ is a member of $\mathcal{F}_1 \cup \mathcal{F}_2$ or an evenly signed graph or a loop.

Now we describe how the pieces defined above can be put together. A trigraph $G = (V, \theta)$ is called an *evenly structured linear interval join* if it can be constructed in the following manner:

- Let (H, s) be an even structure.
- For each edge $e \in E(H)$, let $Z_e \subseteq V(H)$ be the set of ends of e (so $|Z_e| = 1$ or 2).
Let $S_e = (G_e, Z_e)$ such that G_e is a trigraph with $V(G_e) \cap V(H) = Z_e$ and
 - * if e is not on any cycle then S_e is a spot or a thickening of a linear interval stripe,
 - * if e is on a cycle of length > 1 and $s(e) = 0$ then S_e is a thickening of a spring,
 - * if e is on a cycle of length > 1 and $s(e) = 1$ then S_e is a spot,
 - * if e is a loop then $S_e \in \mathcal{C}'$.
- For all distinct $e, f \in E(H)$, $V(G_e) \cap V(G_f) \subseteq Z_e \cap Z_f$.
- Let $V = \cup_{e \in E(H)} V(G_e) \setminus Z_e$ and let θ be given by: for any $u, v \in V$
 - * if $u, v \in V(G_e) \setminus Z_e$ for some $e \in E(H)$ then $\theta(uv) = \theta_{G_e}(uv)$

- * if $u \in N_{G_e}(x)$ and $v \in N_{G_f}(x)$ for distinct $e, f \in E(H)$ with a common end x , then $\theta(uv) = 1$
- * in all other cases, $\theta(uv) = -1$.
- We will write $G = \Omega(H, s, \{S_e : e \in E(H)\})$.

Theorem 7.1 (Chudnovsky and Plumettaz [9]). *A connected trigraph is claw-free and Berge if and only if it is a thickening of a trigraph in \mathcal{C} or an evenly structured linear interval join.*

In the following we produce a different formulation of this result. A vertex x of a trigraph is *simplicial* if $N(x) \neq \emptyset$ and $\{x\} \cup N(x)$ is a strong clique. For $i = 1, 2$, let $G_i = (V_i, \theta_i)$ be a trigraph with a simplicial vertex x_i and with $|V_i| \geq 3$. The *simplicial sum* of G_1, G_2 (over x_1, x_2) is the trigraph $G = (V, \theta)$ such that $V = (V_1 - x_1) \cup (V_2 - x_2)$ and, for all distinct $v_1, v_2 \in V$,

- $\theta(v_1v_2) = \theta_i(v_1v_2)$ if $\{v_1, v_2\} \subseteq V_i$ for some $i = 1, 2$
- $\theta(v_1v_2) = 1$ if $v_i \in N_{G_i}(x_i)$ for both $i = 1, 2$
- $\theta(v_1v_2) = -1$ if otherwise.

We point out that both G_1 and G_2 are contained in G . Moreover, using the language of [9], G admits either a 1-join or a homogeneous set of size ≥ 2 .

Lemma 7.2. *Let G be a simplicial sum of G_1, G_2 . Then G is claw-free if and only if both G_1, G_2 are; and G is Berge if and only if both G_1, G_2 are.*

We omit the proof since it is straightforward. This lemma suggests that we can characterize claw-free Berge trigraphs by determining all such trigraphs that are not simplicial sums. In the following we describe these trigraphs.

Let \mathcal{I} be the class of linear interval trigraphs. Let \mathcal{L} be the class of trigraphs G such that $G^{\geq 0}$ is the line graph of a bipartite multigraph and every *triangle* (a clique of size 3) of G is a strong clique. Let J_1 be the first graph in Figure 7.2. We consider J_1 as a trigraph with no semiadjacent pairs. Let \mathcal{J}_1 consists of trigraphs obtained from J_1 by deleting k of its cubic vertices ($0 \leq k \leq 4$). Let $J_2(n)$ be the second trigraph in Figure 7.2, where Q_1, Q_2 , and all vertical triples are strong cliques, $\theta(uv)$ could be 0, 1, or -1 , and all other pairs are strongly antiadjacent. Note that $J_2(0) \in \mathcal{I}$. Let \mathcal{J}_2 consist of trigraphs of the form $J_2(n) - X$ for all $n \geq 1$ and all $X \subseteq \{u, v\}$. Let $\mathcal{J} = \mathcal{J}_1 \cup \mathcal{J}_2$.




Figure 7.2: J_1 and J_2

Theorem 7.3. *A connected trigraph is claw-free and Berge if and only if it is obtained by simplicial summing thickenings of trigraphs in $\mathcal{C} \cup \mathcal{L} \cup \mathcal{I} \cup \mathcal{J}$.*

We need a few lemmas in order to prove this theorem. A *1-separation* of a multigraph H is a pair (H_1, H_2) of edge-disjoint proper subgraphs of H such that $H_1 \cup H_2 = H$ and $|V(H_1) \cap V(H_2)| = 1$.

Suppose $G = \Omega(H, s, \{S_e\})$. Then a 1-separation (H_1, H_2) of H is called *trivial* if there exists $i \in \{1, 2\}$ such that $H_i = K_2$ and S_f is a spot, where f is the only edge of H_i .

Lemma 7.4. *Suppose $G = \Omega(H, s, \{S_e\})$ and suppose H has a nontrivial 1-separation (H_1, H_2) . Then G is a simplicial sum of two trigraphs.*

Proof. Let x be the common vertex of H_1, H_2 . For $i = 1, 2$, let H'_i be obtained from H_i by adding a new vertex x_i and a new edge xx_i . Let s_i be the signing of H'_i which agrees with s on H_i , and $s_i(xx_i) = 1$. Since all blocks of H'_i (other than xx_i) are blocks of H , (H'_i, s_i) is an even structure. Let S_{xx_i} be a spot and let $G_i = \Omega(H'_i, s_i, \{S_e : e \in E(H'_i)\})$. Since separation (H_1, H_2) is nontrivial, G_i must have ≥ 3 vertices. Now it is straightforward to verify that x_i is a simplicial vertex of G_i ($i = 1, 2$) and G is the simplicial sum of G_1 and G_2 over x_1 and x_2 . ■

Lemma 7.5. *Let H be a thickening of G .*

- (i) *H is claw-free if and only if G is claw-free.*
- (ii) *H is Berge if and only if G is Berge.*

Proof. Part (ii) is (6.4) of [9] and part (i) is easy to verify, as pointed out in [11]. ■

A trigraph G is *quasi-line* if $N(v)$ is the union of two strong cliques for every $v \in V(G)$. It is easy to see that if G is quasi-line then G is claw-free. A trigraph G is *cobipartite* if $V(G)$ is the union of two strong cliques. Clearly, if G cobipartite then G is quasi-line and thus is claw-free. It is also clear that every connected cobipartite trigraph with ≥ 2 vertices is a thickening of a two-vertex trigraph. Thus every cobipartite trigraph is Berge.

Proof of Theorem 7.3. To prove the backward implication, by Lemma 7.2 and Lemma 7.5, we only need to consider trigraphs $G \in \mathcal{C} \cup \mathcal{L} \cup \mathcal{I} \cup \mathcal{J}$. If $G \in \mathcal{C}$ then the result follows from Theorem 7.1. If $G \in \mathcal{I}$ then G is claw-free [11] and Berge [9]. If $G \in \mathcal{L} \cup \mathcal{J}$ then G is quasi-line and thus G is claw-free. If $G \in \mathcal{J}$, then deleting simplicial vertices from G results in a cobipartite trigraph, which implies that G is Berge. Finally, assume $G \in \mathcal{L}$ and $G^{\geq 0} = L(B)$ is the line graph of a bipartite multigraph B . We need to show that G is Berge. Since no semiadjacent pairs are contained in a triangle, every hole of G must come from a cycle of B and thus G contains no odd holes. If G has an antihole $v_1 v_2 \dots v_n v_1$ with $n \geq 7$, then we consider the restriction of G on v_1, \dots, v_6 . If $\theta(v_i v_{i+1}) = -1$ for all $i = 1, \dots, 5$, then the graph X formed by $\{v_i v_j : \theta(v_i v_j) \geq 0\}$ would be the complement of a path on six vertices, which is one of the minimal non-line-graphs. This is impossible since X is an induced subgraph of $L(B)$. So $\theta(v_i v_{i+1}) = 0$ holds for some i , which makes v_i, v_{i+1}, x_k a triangle for some k . This contradiction (two semiadjacent vertices are contained in a triangle) shows that G contains no antihole of length ≥ 7 . Thus G is Berge, which completes the proof of the backward direction.

To prove the forward implication, by Theorem 7.1, we assume $G = \Omega(H, s, \{S_e\})$. Since G is connected, H is connected as well. By Lemma 7.4, we also assume that all 1-separations of H are trivial. Let U be the set of all degree-one vertices u of H for which if e is the only edge incident with u then S_e is a spot. We assume $V(H) \neq U$ because otherwise $H = K_2$ and $G = K_1$ and thus the result holds. Let $H_0 = H - U$. Note that H_0 is connected, as H is connected. Moreover, by its construction, H_0 does not have a 1-separation. Thus either $H_0 = K_1$ or H_0 is a block of H .

Suppose H_0 is K_1 or K_2 . It follows that H is a tree with 1, 2, or 3 edges. Moreover, S_e is a

thickening of a linear interval strip for at most one e , and every other S_e is a spot. In all cases, it is routine to check that G is a thickening of a trigraph in \mathcal{I} .

Suppose H_0 is a loop e . Let $S_e = (G_e, \{z\}) \in \mathcal{C}'$ and let G_e be a thickening of $C \in \mathcal{C}$. If H has ≥ 2 edges then H consists of e and a pendent edge f with S_f a spot. It follows that $G = G_e$, which is a thickening of a trigraph in \mathcal{C} . So e is the only edge of H and $G = G_e - z$. If z is not the only vertex of X_{a_i} (here we use the notation in the definition of \mathcal{C}') then G is also a thickening of C . If z is the unique vertex of X_{a_i} then G is cobipartite. In this case G is a thickening of a two-vertex trigraph and thus G is a thickening of a trigraph in \mathcal{I} .

Suppose none of the last two cases occurs. Then H_0 is a block in which every edge is on a cycle of length ≥ 2 . Let s_0 be the restriction of s on H_0 . Then (H_0, s_0) is either in $\mathcal{F}_1 \cup \mathcal{F}_2$ or evenly signed. First we assume (H_0, s_0) is evenly signed. Then (H, s) is also evenly signed. Moreover, S_e is a thickening of a spring for every edge in $E_0 = \{e \in E(H_0) : s(e) = 0\}$, and S_e is a spot for every other edge of H . Let S'_e be a spring for each $e \in E_0$ and let $S'_e = S_e$ for every other edge of H . Then G is a thickening of $G' = \Omega(H, s, \{S'_e\})$. Now we only need to show that $G' \in \mathcal{L}$. Let H' be obtained from H by subdividing each edge in E_0 exactly once. Then H' is bipartite. It follows from the construction of Ω that adjacent pairs of G' are exactly adjacent pairs of the line graph $L(H')$. In addition, all semiadjacent pairs of G' come from a spring, and thus no such pair is contained in a triangle. Therefore, G' belongs to \mathcal{L} , as required.

It remains to consider the case $(H_0, s_0) \in \mathcal{F}_1 \cup \mathcal{F}_2$. If $(H_0, s_0) \in \mathcal{F}_1$, then H is obtained from K_4 by adding parallel edges and adding pendent edges to distinct vertices. Moreover, every S_e is a spot. It follows that G is an ordinary graph (meaning that G has no semiadjacent pairs) and this graph is exactly $L(H)$. Now it is clear that G is a thickening of $L(\text{si}(H))$, which belongs to \mathcal{J}_1 . So we assume $(H_0, s_0) \in \mathcal{F}_2$. Let $V(H_0) = \{x_1, x_2, y_1, \dots, y_m\}$ ($m \geq 1$) such that x_i ($i = 1, 2$) is adjacent to all other vertices. Like before, we assume that H_0 has no parallel edges, except for two possible edges e_0, e_1 between x_1, x_2 , and such that $s(e_0) = 0$ and $s(e_1) = 1$. We also assume that S_{e_0} is a spring, if e_0 is present. Suppose H is obtained by adding pendent edges to y_1, \dots, y_n ($n \geq 0$) and to k of x_1, x_2 ($0 \leq k \leq 2$). If e_0 is present, then G is a thickening of $J_2(n)$, where $\theta(uv) = 0$. So assume that e_0 is not in H , and thus $G = L(H)$. For $i = 1, 2$, let Q_i be the clique of G formed by edges of H incident with x_i . Let $Q'_i = Q_i - \{x_1x_2, x_1y_1, \dots, x_1y_n\}$. If $Q'_1 \neq \emptyset$ is neither complete nor anticomplete to $Q'_2 \neq \emptyset$, then again G is a thickening of $J_2(n)$ with $\theta(uv) = 0$. In the remainder cases (which are: some Q'_i is empty, or $Q'_1 \neq \emptyset$ is complete or anticomplete to $Q'_2 \neq \emptyset$), if $n = 0$ then G is a thickening of K_3 , and if $n \geq 1$ then $G = J_2(n) - X$ for some $X \subseteq \{u, v\}$. ■

8 Claw-free box-perfect graphs

In this section we prove the following.

Theorem 8.1. *A claw-free perfect graph is box-perfect if and only if it is S_3 -free.*

We divide the proof into several lemmas. Let G be a trigraph. We call G a *sun* if $G \in \text{tri}(S_3)$. We call G an *incomparability* trigraph if $G^{\geq 0}$ is an incomparability graph. We call G *elementary* if it is a thickening of a trigraph in \mathcal{L} . We remark that when an elementary trigraph has no semiadjacent pairs then they are exactly *elementary graphs* discussed in [18].

Lemma 8.2. *Let G be a connected Berge trigraph. If G is $\{\text{claw}, \text{sun}\}$ -free then G is obtained by simplicial summing incomparability trigraphs and elementary trigraphs.*

Proof. Since G is connected, Berge, and claw-free, by Theorem 7.3, G is obtained by simplicial summing thickenings of trigraphs in $\mathcal{C} \cup \mathcal{L} \cup \mathcal{I} \cup \mathcal{J}$. Therefore, we may assume that G is a thickening of a trigraph $G_0 \in \mathcal{C} \cup \mathcal{L} \cup \mathcal{I} \cup \mathcal{J}$. If $G_0 \in \mathcal{L}$ then G is elementary and we are done. If $G_0 \in \mathcal{C}$ then $G_0|\{a_1, a_2, a_3, x_1, x_2, x_3\}$ (here we are using the notation in the definition of \mathcal{C}) is a sun and thus G contains a sun, which is impossible. So we assume that $G_0 \in \mathcal{I} \cup \mathcal{J}$. In the following we prove that G is an incomparability trigraph.

Suppose $G_0 \in \mathcal{I}$. Then vertices of G_0 can be ordered as v_1, \dots, v_n such that if $i < j < k$ and $\theta_0(v_i v_k) \geq 0$ then $\theta_0(v_i v_j) = \theta_0(v_j v_k) = 1$. Using the notation in the definition of thickening, we let $X_{v_i} = \{x_{i,j} : j = 1, \dots, n_i\}$ ($1 \leq i \leq n$). Now we define a binary relation \prec on $V(G)$ such that $x_{i_1, j_1} \prec x_{i_2, j_2}$ if $\theta(x_{i_1, j_1} x_{i_2, j_2}) = -1$ and (i_1, j_1) is lexicographically smaller than (i_2, j_2) . We claim that \prec is transitive. Suppose $x_{i_1, j_1} \prec x_{i_2, j_2} \prec x_{i_3, j_3}$. Since each X_{v_i} is a strong clique, we must have $i_1 < i_2 < i_3$. It follows that $\theta_0(v_{i_1} v_{i_2}) \leq 0$ and $\theta_0(v_{i_2} v_{i_3}) \leq 0$. As a result, $\theta_0(v_{i_1} v_{i_3}) = -1$ and thus $\theta(x_{i_1, j_1} x_{i_3, j_3}) = -1$, which proves our claim. This claim implies that the complement of $G^{\geq 0}$ is the comparability graph of poset $(V(G), \prec)$, which proves that $G^{\geq 0}$ is an incomparability graph and thus G is an incomparability trigraph.

Now suppose $G_0 \in \mathcal{J}$. We claim that G_0 is a thickening of a trigraph in \mathcal{I} . This claim clearly implies that G is a thickening of trigraph in \mathcal{I} , and thus the last paragraph proves that G is an incomparability trigraph.

Before proving the claim we make an observation. It is clear that every cobipartite trigraph is a thickening of a trigraph that has exactly two vertices and that the two vertices are semiadjacent. Since this two-vertex trigraph is in \mathcal{I} , our claim holds if G_0 is cobipartite.

We first consider the case $G_0 \in \mathcal{J}_1$. If G_0 has two or more cubic vertices then G_0 contains an induced S_3 . So G_0 contains at most one cubic vertex and in this case G_0 is cobipartite. Next we assume $G_0 \in \mathcal{J}_2$ and let $G_0 = J_2(n) - X$ (see the definition of \mathcal{J}_2). Let $x_i y_i z_i$ ($1 \leq i \leq n$) denote the vertical triangles of $J_2(n)$, where $y_i \in Q_1$ and $z_i \in Q_2$. If $n \geq 3$ then $G_0|\{w, x_1, y_1, z_1, y_2, z_3\}$ is a sun. So we have $n \leq 2$. Now it is straightforward to verify that either G_0 is cobipartite, or G_0 contains a sun (found in a similar way), or $G_0 = J_2(2) - \{u, v\}$. In the last case, G_0 is a thickening of the trigraph $G^* = (\{t_1, t_2, t_3, t_4, t_5\}, \theta^*)$, where $\theta^*(t_i t_{i+1}) = 1$ ($i = 1, 2, 3, 4$), $\theta^*(t_2 t_4) = 0$, and $\theta^*(t_i t_j) = -1$ for all other pairs. This completes the proof of our claim and also completes the proof of the lemma. ■

Although simplicial sum was defined for trigraphs, this operation can be naturally inherited by ordinary graphs. Moreover, we have the following.

Lemma 8.3. *The simplicial sum of two ESP graphs is ESP.*

Proof. Let G be the simplicial sum of G_1 and G_2 over x_1 and x_2 , where G_i is an ESP graph with a simplicial vertex x_i for $i = 1, 2$. Let Λ be a set of maximal cliques of G . Note that $N_{G_1}(x_1) \cup N_{G_2}(x_2)$ is the only maximal clique of G that contains edges between $N_{G_1}(x_1)$ and $N_{G_2}(x_2)$. For $i = 1, 2$, let Λ_i consist of members of Λ that are cliques in G_i . Since G_i is ESP, Λ_i has an equitable subpartition $(\Lambda_{i1}, \Lambda_{i2})$. If Λ does not contain the clique $N_{G_1}(x_1) \cup N_{G_2}(x_2)$, then $(\Lambda_{11} \cup \Lambda_{21}, \Lambda_{12} \cup \Lambda_{22})$ is clearly an equitable subpartition of Λ . Now assume that Λ contains $N_{G_1}(x_1) \cup N_{G_2}(x_2)$. For $i = 1, 2$, let

$\Lambda'_i = \Lambda_i \cup \{\{x_i\} \cup N_{G_i}(x_i)\}$. Note that $d_{\Lambda'_i}(x_i) = 1$. Since G_i is ESP, Λ'_i has an equitable subpartition $(\Lambda'_{i1}, \Lambda'_{i2})$. Without loss of generality, suppose $d_{\Lambda'_{i1}}(x_i) = 1$ and $d_{\Lambda'_{i2}}(x_i) = 0$ for $i = 1, 2$. Let Λ' be obtained from $\Lambda'_{11} \cup \Lambda'_{21}$ by replacing the two cliques containing x_1 or x_2 by $N_{G_1}(x_1) \cup N_{G_2}(x_2)$; and set $\Lambda'' = \Lambda'_{12} \cup \Lambda'_{22}$. Then (Λ', Λ'') is an equitable subpartition of Λ . ■

Lemma 8.4. *Let Λ be a set of cliques of a graph G , for which $V(G)$ is partitioned into two cliques X, Y . Then G has a multiset Λ' of cliques such that*

- (i) $|\Lambda'| = |\Lambda|$ and $d_{\Lambda'}(v) = d_\Lambda(v)$, for all $v \in V(G)$;
- (ii) members of Λ' can be enumerated as $Q_1, \dots, Q_{|\Lambda|}$ such that every $v \in X$ appears in the first $d_\Lambda(v)$ terms and every $v \in Y$ appears in the last $d_\Lambda(v)$ terms.

Proof. For each $i = 1, \dots, |\Lambda|$, let $X_i = \{x \in X : i \leq d_\Lambda(x)\}$ and $Y_i = \{y \in Y : i \geq |\Lambda| - d_\Lambda(y) + 1\}$. Then for every i , $Q_i = X_i \cup Y_i$ is a clique since $d_\Lambda(x) + d_\Lambda(y) \leq |\Lambda|$ holds for all non-adjacent $x \in X$ and $y \in Y$. Now it is clear that $\Lambda' = \{Q_1, \dots, Q_{|\Lambda|}\}$ satisfies the requirements. ■

Lemma 8.5. *Elementary graphs are ESP.*

Proof. Let elementary graph H be obtained by thickening a trigraph G , where $G^{\geq 0}$ is the line graph of a bipartite multigraph B and such that semiadjacent pairs of G are not contained in any triangle. Let (Z_1, Z_2) be a partition of $V(B)$ into two stable sets. Let u_1v_1, \dots, u_nv_n be the semiadjacent pairs of G . Let $(X_v : v \in V(G))$ be the partition of $V(H)$ over which G is thickened. For $i = 1, \dots, n$, let $H_i = H[X_{u_i} \cup X_{v_i}]$. Since no semiadjacent pairs of G are contained in a triangle, it is easy to see that for each maximal clique C of H , either C is a maximal clique of some H_i or $C = \cup\{X_v : v \in Q\}$ for some maximal strong clique Q of G . On the other hand, since $G^{\geq 0} = L(B)$, for each maximal clique Q of G there exists a vertex z of B such that members of Q are precisely edges of B that are incident with z . We will say that Q and $C = \cup\{X_v : v \in Q\}$ come from z . Note that, if Q, Q' are maximal cliques of G with $u_i \in Q - Q'$ and $v_i \in Q' - Q$ for some i , then Q and Q' come from vertices that both belong to Z_1 or both belong to Z_2 .

Let Λ be a set of maximal cliques of H . We need to show that Λ admits an equitable subpartition. For $i = 1, \dots, n$, let $\Lambda^{(i)} = \{C \in \Lambda : C \subseteq V(H_i)\}$. Let $\Lambda^{(0)} = \Lambda - \Lambda^{(1)} \dots - \Lambda^{(n)}$. We assume by Lemma 8.4 that members of each $\Lambda^{(i)}$ are enumerated as $C_1^{(i)}, \dots, C_{n_i}^{(i)}$ such that every $x \in X_{u_i}$ appears in the first $d_{\Lambda^{(i)}}(x)$ terms and every $x \in X_{v_i}$ appears in the last $d_{\Lambda^{(i)}}(x)$ terms. In the following we define a partition (Λ_1, Λ_2) of Λ . To verify that (Λ_1, Λ_2) is an equitable subpartition of Λ we only need to verify $\min\{d_{\Lambda_1}(x), d_{\Lambda_2}(x)\} \geq \lfloor d_\Lambda(x)/2 \rfloor$, for all $x \in V(H)$.

We first consider $\Lambda^{(0)}$. If $C \in \Lambda^{(0)}$ then C comes from a vertex z of B . In this case we put C into Λ_i if $z \in Z_i$ ($i = 1, 2$). Since each $v \in V(G)$ is contained in at most two maximal cliques, we deduce that $d_\Lambda(x) \leq 2$ for all $x \in V(H) - V(H_1) - \dots - V(H_n)$. For these x , our partition of $\Lambda^{(0)}$ guarantees that $\min\{d_{\Lambda_1}(x), d_{\Lambda_2}(x)\} \geq \lfloor d_\Lambda(x)/2 \rfloor$.

For cliques in each $\Lambda^{(i)}$ ($i = 1, 2, \dots, n$) we consider three cases. If none of X_{u_i}, X_{v_i} is contained in any clique of $\Lambda^{(0)}$, then $d_\Lambda(x) = d_{\Lambda^{(i)}}(x)$ for all $x \in V(H_i)$. Moreover, for each $x \in V(H_i)$, since cliques containing x appear consecutively in the sequence $C_1^{(i)}, \dots, C_{n_i}^{(i)}$, putting $C_j^{(i)}$ into Λ_1 for all odd j and putting $C_j^{(i)}$ into Λ_2 for all even j lead to $\min\{d_{\Lambda_1}(x), d_{\Lambda_2}(x)\} \geq \lfloor d_\Lambda(x)/2 \rfloor$.

If exactly one of X_{u_i}, X_{v_i} is contained in a clique of $\Lambda^{(0)}$, we assume by symmetry that X_{u_i} is contained in a clique $C \in \Lambda^{(0)}$. We also assume without loss of generality that C has been placed

into Λ_2 . For each $x \in V(H_i)$, since cliques containing x appear consecutively in the sequence $X_{u_i}, C_1^{(i)}, \dots, C_{n_i}^{(i)}$, putting $C_j^{(i)}$ into Λ_1 for all odd j and putting $C_j^{(i)}$ into Λ_2 for all even j lead to $\min\{d_{\Lambda_1}(x), d_{\Lambda_2}(x)\} \geq \lfloor d_{\Lambda}(x)/2 \rfloor$.

If both X_{u_i}, X_{v_i} are contained in cliques, say C, D , of $\Lambda^{(0)}$, by discussion in the first paragraph of this proof, we assume that both C, D have been placed into Λ_2 . We consider two subcases. Suppose n_i is odd. For each $x \in V(H_i)$, since cliques containing x appear consecutively in the sequence $X_{u_i}, C_1^{(i)}, \dots, C_{n_i}^{(i)}, X_{v_i}$, putting $C_j^{(i)}$ into Λ_1 for all odd j and putting $C_j^{(i)}$ into Λ_2 for all even j lead to $\min\{d_{\Lambda_1}(x), d_{\Lambda_2}(x)\} \geq \lfloor d_{\Lambda}(x)/2 \rfloor$. Now suppose n_i is even. For each $x \in V(H_i)$, note that cliques containing x appear consecutively in the sequence $C_1^{(i)}, X_{u_i}, C_2^{(i)}, \dots, C_{n_i}^{(i)}, X_{v_i}$, unless $x \in C_1^{(i)} \cap X_{v_i}$. In this case we put $C_j^{(i)}$ into Λ_2 for all odd $j > 1$ and we put the rest into Λ_1 . For each $x \in V(H_i) - (C_1^{(i)} \cap X_{v_i})$, it is clear that $\min\{d_{\Lambda_1}(x), d_{\Lambda_2}(x)\} \geq \lfloor d_{\Lambda}(x)/2 \rfloor$. For each $x \in C_1^{(i)} \cap X_{v_i}$, we have $d_{\Lambda}(x) = 1 + n_i$. Our partition yields $d_{\Lambda_2}(x) = n_i/2$, which also leads to $\min\{d_{\Lambda_1}(x), d_{\Lambda_2}(x)\} \geq \lfloor d_{\Lambda}(x)/2 \rfloor$. ■

Proof of Theorem 8.1. The forward implication is clear, so we only need to consider the backward implication. Let G be perfect and $\{claw, S_3\}$ -free. By Lemma 8.2, each component of G is obtained by simplicial summing incomparability graphs and elementary graphs. By Theorem 4.2 and Lemma 8.5, incomparability graphs and elementary graphs are ESP. Thus G is ESP by Lemma 8.3, which proves that G is box-perfect by Theorem 3.7. ■

References

- [1] C. Berge, The q -perfect graphs. II, *Matematiche (Catania)* **47** (1992), 205-211.
- [2] M. Burić and J. Uhry, Parity graphs, *Ann. of Discrete Math.* **66** (1982), 1-26.
- [3] K. Cameron, *Polyhedral and Algorithmic Ramifications of Antichains*, Ph.D. Thesis, University of Waterloo, 1982. (Supervisor: J. Edmonds)
- [4] K. Cameron, A min-max relation for the partial q -colourings of a graph. II. Box perfection, *Discrete Math.* **74** (1989), 15-27.
- [5] K. Cameron and J. Edmonds, Coflow polyhedra, *Discrete Math.* **101** (1992), 1-21.
- [6] P. Camion, Characterization of totally unimodular matrices, *Proc. Amer. Math. Soc.* **16** (1965), 1068-1073.
- [7] X. Chen, G. Ding, and W. Zang, The box-TDI system associated with 2-edge connected spanning subgraphs, *Discrete Appl. Math.* **157** (2009), 118-125.
- [8] M. Chudnovsky, G. Cornuéjols, X. Liu, P. Seymour, and K. Vušković, Recognizing Berge graphs, *Combinatorica* **25** (2005), 143-186.
- [9] M. Chudnovsky and M. Plumettaz, The structure of claw-free perfect graphs, *J. Graph Theory* **75** (2014), 203-230.
- [10] M. Chudnovsky, N. Robertson, P. Seymour, and R. Thomas, The strong perfect graph theorem, *Ann. of Math. (2)* **164** (2006), 51-229.

- [11] M. Chudnovsky and P. Seymour, Claw-free graphs. V. Global structure, *J. Combin. Theory Ser. B* **98** (2008), 1373-1410.
- [12] M. Chudnovsky and P. Seymour, Even pairs in Berge graphs, *J. Combin. Theory Ser. B* **99** (2009), 370-377.
- [13] W. Cook, On box totally dual integral polyhedra, *Math. Programming* **34** (1986), 48-61.
- [14] J. Edmonds and R. Giles, A min-max relation for submodular functions on graphs, *Ann. of Discrete Math.* **1** (1977), 185-204.
- [15] T. Gallai, Transitiv orientierbare graphen, *Acta Math. Acad. Sci. Hungar.* **18** (1967), 25-66.
- [16] L. Lovász, Normal hypergraphs and the perfect graph conjecture, *Discrete Math.* **2** (1972), 253-267.
- [17] L. Lovász, Perfect graphs, in: *Selected Topics in Graph Theory II* (L. Beineke and R. Wilson, eds.), Academic Press, 1983, pp. 55-87.
- [18] F. Maffray and B. Reed, A description of claw-free perfect graphs, *J. Combin. Theory Ser. B* **75** (1999), 134-156.
- [19] H. Sachs, On the Berge conjecture concerning perfect graphs, in: *Combinatorial Structures and Their Applications*, Gordon and Breach, New York, 1970, pp. 377-384.
- [20] A. Schrijver, Min-max results in combinatorial optimization, in: *Mathematical Programming: The State of the Art*, Springer, Berlin, 1983, pp. 439-500.
- [21] A. Schrijver, *Theory of Linear and Integer Programming*, John Wiley & Sons, New York, 1986.
- [22] A. Schrijver, *Combinatorial Optimization: Polyhedra and Efficiency*, Springer-Verlag, Berlin, 2003.