




Chinese Pharmaceutical Association  
Institute of Materia Medica, Chinese Academy of Medical Sciences

Acta Pharmaceutica Sinica B

[www.elsevier.com/locate/apsb](http://www.elsevier.com/locate/apsb)  
[www.sciencedirect.com](http://www.sciencedirect.com)



ORIGINAL ARTICLE

# Garlic-derived compound *S*-allylmercaptocysteine inhibits hepatocarcinogenesis through targeting LRP6/Wnt pathway



Jia Xiao<sup>a,b,c,†</sup>, Feiyue Xing<sup>b,†</sup>, Yingxia Liu<sup>a,†</sup>, Yi Lv<sup>b</sup>, Xiaogang Wang<sup>d</sup>,  
Ming-Tat Ling<sup>f</sup>, Hao Gao<sup>g</sup>, Songying Ouyang<sup>h</sup>, Min Yang<sup>a</sup>, Jiang Zhu<sup>i</sup>,  
Yu Xia<sup>b</sup>, Kwok-Fai So<sup>e,\*\*</sup>, George L. Tipoe<sup>c,\*</sup>

<sup>a</sup>State Key Discipline of Infectious Diseases, Shenzhen Third People's Hospital, Shenzhen 518112, China

<sup>b</sup>Department of Immunobiology, Institute of Tissue Transplantation and Immunology, Jinan University, Guangzhou 510632, China

<sup>c</sup>School of Biomedical Sciences, The University of Hong Kong, Hong Kong, China

<sup>d</sup>Department of Cell Biology & Institute of Biomedicine, Jinan University, Guangzhou 510632, China

<sup>e</sup>GMH Institute of Central Nervous System Regeneration, Jinan University, Guangzhou 510632, China

<sup>f</sup>Australian Prostate Cancer Research Centre-Queensland and Institute of Health and Biomedical Innovation, Queensland University of Technology, Brisbane, Queensland, Australia

<sup>g</sup>Institute of Traditional Chinese Medicine and Natural Products, College of Pharmacy, Jinan University, Guangzhou 510632, China

<sup>h</sup>Biomedical Research Center of South China, College of Life Sciences, Fujian Normal University, Fuzhou 350117, China

<sup>i</sup>JM Medical (Shenzhen), LLC, Shenzhen 518112, China

Received 24 June 2017; received in revised form 31 August 2017; accepted 8 September 2017

## KEY WORDS

*S*-allylmercaptocysteine;  
HCC;  
Wnt;

**Abstract** Whether and how garlic-derived *S*-allylmercaptocysteine (SAMC) inhibits hepatocellular carcinoma (HCC) is largely unknown. In the current study, the role of low-density lipoprotein receptor (LDLR)-related protein 6 (LRP6) in HCC progression and the anti-HCC mechanism of SAMC was examined in clinical sample, cell model and xenograft/orthotopic mouse models. We demonstrated that SAMC inhibited

*Abbreviations:* Axin1, axis inhibition protein 1; DKK-1, Dickkopf Wnt signaling pathway inhibitor 1; DVL2, disheveled 2; FADD, Fas-associated protein with death domain; HCC, hepatocellular carcinoma; KD, knock-down; LDH, lactate dehydrogenase; LRP6, low-density lipoprotein receptor (LDLR)-related protein 6; MCL-1, myeloid cell leukemia-1; NAFLD, non-alcoholic fatty liver disease; PCNA, proliferating cell nuclear antigen; SAC, *S*-allylcysteine; SAMC, *S*-allylmercaptocysteine; SPR, surface plasmon resonance; TCF/LEF, T-cell factor/lymphoid enhancing factor; TSA, thermal shift assay;  $T_m$ , melting temperature

\*Corresponding author. Tel.: +852 3917 9185; fax: +852 2817 0857.

\*\*Corresponding author. Tel./fax: +86 20 8522 3563.

E-mail addresses: [hmaskf@hku.hk](mailto:hmaskf@hku.hk) (Kwok-Fai So), [tgeorge@hku.hk](mailto:tgeorge@hku.hk) (George L. Tipoe).

<sup>†</sup>These authors made equal contributions to this work.

Peer review under responsibility of Institute of Materia Medica, Chinese Academy of Medical Sciences and Chinese Pharmaceutical Association.

<http://dx.doi.org/10.1016/j.apsb.2017.10.003>

2211-3835 © 2018 Chinese Pharmaceutical Association and Institute of Materia Medica, Chinese Academy of Medical Sciences. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

LRP6;  
Human;  
Nude mice

cell proliferation and tumorigenesis, while induced apoptosis of human HCC cells without influencing normal hepatocytes. SAMC directly interacted with Wnt-pathway co-receptor LRP6 on the cell membrane. LRP6 was frequently over-expressed in the tumor tissue of human HCC patients (66.7% of 48 patients) and its over-expression only correlated with the over-expression of  $\beta$ -catenin, but not with age, gender, tumor size, stage and metastasis. Deficiency or over-expression of LRP6 in hepatoma cells could partly mimic or counteract the anti-tumor properties of SAMC, respectively. *In vivo* administration of SAMC significantly suppressed the growth of Huh-7 xenograft/orthotopic HCC tumor without causing undesirable side effects. In addition, stable down-regulation of LRP6 in Huh-7 facilitated the anti-HCC effects of SAMC. In conclusion, LRP6 can be a potential therapeutic target of HCC. SAMC is a promising specific anti-tumor agent for treating HCC subtypes with Wnt activation at the hepatoma cell surface.

© 2018 Chinese Pharmaceutical Association and Institute of Materia Medica, Chinese Academy of Medical Sciences. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Currently, liver cancer is the sixth most common malignant disease and the second leading cause of cancer death worldwide. There are approximately 50.5% of new liver cancer cases in China in each year, ~75% of which is hepatocytes-derived hepatocellular carcinoma (HCC)<sup>1</sup>. The survival rate after the onset of HCC symptoms is generally less than one year and as to date no effective clinical therapeutic strategy with desirable effects has been developed<sup>2</sup>. Therefore, elucidating the molecular mechanisms on the initiation and progression of HCC is critical for the control of this fatal disease.

The canonical Wnt/ $\beta$ -catenin pathway is aberrantly activated in HCC<sup>3</sup>. Activation of the Wnt pathway is through the binding of Wnt family proteins (*e.g.*, Wnt3a) to the cell surface receptors low-density lipoprotein receptor (LDLR)-related protein 5 (LRP5) and/or LRP6. After that, phosphorylated receptors recruit disheveled homologue proteins, *e.g.*, disheveled 2/3 (DVL2/3), and axis inhibition protein 1 (axin1) to stabilize and promote the nuclear translocation of  $\beta$ -catenin, which acts as a transactivator of T-cell factor/lymphoid enhancing factor (TCF/LEF) transcription factors to regulate the expression of key genes for cell proliferation, differentiation, and tumorigenesis<sup>4</sup>. In the liver, many temporal roles of the Wnt/ $\beta$ -catenin pathway have been identified during its development and maintenance of physiological homeostasis<sup>5</sup>. Emerging evidence suggests that dysregulated signaling of the Wnt/ $\beta$ -catenin pathway lead to hepatic carcinogenesis<sup>6–8</sup>. Recently, LRP6 has been identified as a novel nutritional therapeutic target for several liver diseases, including non-alcoholic fatty liver disease (NAFLD) and hyperlipidemia<sup>9</sup> while another study also found that up-regulation of LRP6 was associated with enhanced hepatic carcinogenesis and cell invasion<sup>10</sup>. Therefore, we hypothesized that LRP6 might be a direct target for nutraceutical agents with anti-HCC properties.

Garlic is used as a medicinal food for its anti-bacterial, immunoregulatory and anti-tumor properties in many countries for more than 2000 years<sup>11</sup>. Epidemiological studies indicate an association between garlic consumption and decreased risk of gastrointestinal tract cancers<sup>11</sup>. *S*-allylmercaptocysteine (SAMC) is a water-soluble active compound derived from aged garlic. We have demonstrated its potent hepato-protective properties and mechanisms in acute liver injury and NAFLD<sup>12,13</sup>. Its anti-tumor effects have been demonstrated in colon cancer<sup>14</sup>, prostate

cancer<sup>15,16</sup>, bladder cancer<sup>17</sup>, breast cancer<sup>18</sup>, and gastric cell cancer<sup>19</sup>. A very recent study found that SAMC induced apoptosis in human HepG2 cell through targeting the cross-talk between the transforming growth factor- $\beta$  and the mitogen-activated protein kinase pathways<sup>20</sup>. However, mechanistic data regarding the detailed anti-HCC functions of SAMC, particularly its “immediate receptor” when in contact with the tumor cell, is lacking. Therefore, in the current study, we aimed to investigate the anti-tumor effects and mechanisms of SAMC in human and mouse HCC cell lines and xenograft/orthotopic models, with emphasis in its direct target on the cell membrane.

## 2. Materials and methods

### 2.1. Patient samples and analysis

Use of human tissue samples in this project was approved by the Ethical Committee of Shenzhen Third People's Hospital. All patients were given formal notification and written consent on the use of the clinical specimens for research. Forty-eight pairs of HCC tissues and their corresponding non-tumorous liver tissues (1 cm away from the tumor), as well as 6 liver tissues from healthy people, were employed for analyses. The clinicopathological features of all these patients are listed in [Supplementary Information Table S1](#).

### 2.2. Generation of LRP6 rescue and over-expressed constructs

The cloning and generation of a codon-modified shRNA-resistant LRP6 (LRP6 rescue) construct was conducted as previously reported<sup>21</sup>.

### 2.3. GST-E-cadherin pull-down assay

The GST-E-cadherin pull-down assay was performed as previously described<sup>22</sup>. Western blotting was performed using an antibody to  $\beta$ -catenin.

### 2.4. Surface plasmon resonance (SPR) and thermal shift assay (TSA)

Analysis of direct binding between SAMC and LRP6 protein was performed by using SPR and TSA as previously described<sup>23</sup>. Apparent equilibrium dissociation constants ( $K_d$ ) were then

calculated as the ratio of  $k_d/k_a$ . In TSA, we used 0.04 mg/mL of recombinant human LRP6 protein with or without 0.2 mmol/L of SAMC in phosphate buffered saline (PBS). Data were analyzed with the differential scanning fluorimetry analysis tool (Excel based) using the curve-fitting software XLfit 5 (ID Business Solutions Ltd., Bridgewater, NJ, USA)<sup>23</sup>.

### 2.5. Xenograft and orthotopic HCC nude mice model

Male nude mice (Athymic *nu/nu*, 5–6 weeks, 20–25 g) were purchased from Guangdong Medical Laboratory Center (Guangdong, China). For the establishment of a subcutaneous (s.c.) xenograft HCC model, mice were s.c. injected at the dorsal region with  $1 \times 10^6/150 \mu\text{L}$  (low dose group) or  $4 \times 10^6/150 \mu\text{L}$  (high dose group) viable normal or *LRP6* knockdown Huh-7 cells. Seven days after Huh-7 injection, 300 mg/kg SAMC was treated by daily oral gastric lavage feeding ( $n=6$  for each group of mice)<sup>16</sup>. Tumor volume was assessed with digital calipers at days 8, 13, 18, 23, and 28 post Huh-7 injection. The tumor volume was calculated using the formula:  $\pi/6 \times \text{larger diameter} \times \text{smaller diameter}^2$ <sup>23</sup>.

For the orthotopic HCC model, a single tumor nodule could be observed in the liver after 6 days of an injection of  $2 \times 10^6$  wild-type or *LRP6* knock-down Huh-7 cells into the left liver lobe of nude mice. Then mice received daily oral gastric lavage feeding of 300 mg/kg SAMC or saline ( $n=10$  for each group of mice). Median survival analysis was conducted using the Kaplan–Meier methods in GraphPad Prism v6.0 software (GraphPad Software, San Diego, CA, USA)<sup>24</sup>. Mice were subjected to anesthesia and then killed if moribund. The total observation duration was 70 days. Tumor nodule was collected for the intra-tumor expression check of key proteins.

### 2.6. Statistical analysis

Data from each group were expressed as mean  $\pm$  standard error of the mean (SEM). Statistical comparisons between groups were done using the Kruskal–Wallis test followed by Dunn's *post hoc* test to detect differences in all groups. Clinical data were used Fisher's exact test to compare and calculate *P*-values. A  $P < 0.05$  was statistically significant (Prism 5.0, Graphpad software, Inc., San Diego, CA, USA).

Full methods and any associated references are available in [supplementary Information](#).

## 3. Results

### 3.1. *LRP6* is frequently over-expressed in human HCC samples

It was shown that the mRNA level of *LRP6* was significantly higher in both hepatoma cell lines (Hep3B and Huh-7) than normal cell line LO-2 ( $P < 0.001$ ). Interestingly, Huh-7 expressed a higher *LRP6* level than Hep3B ( $P=0.02$ , Fig. 1A). Then we examined its transcriptional level in the liver tissues of healthy subjects ( $n=6$ ), non-cancerous sections ( $n=48$ ) and cancerous sections ( $n=48$ ) of HCC patients (Fig. 1B), whose clinicopathological characteristics were presented in [Supplementary Information Table S1](#). *LRP6* transcripts were frequently and significantly up-regulated ( $P < 0.001$ ) in the cancerous liver tissue sections of patients than their corresponding non-cancerous tissues. Quantitative PCR showed that *LRP6* mRNA expression was elevated in 32 out of 48 (66.7%) cancerous liver tissues of HCC patients (defined as a 2-fold expressional elevation). Five

representative Western blot results were presented in Fig. 1C. Immunohistochemistry results indicated obvious over-expression of LRP6 protein in the cytoplasmic part of tumor cells while the non-cancerous liver tissue only showed relatively lower LRP6 signals (Fig. 1D). In addition, clinicopathological correlation analysis exhibited that the association between LRP6 over-expression and  $\beta$ -catenin over-expression was statistically significant in those patients ( $P=0.0039$ , [Supplementary Information Table S2](#)).


### 3.2. SAMC selectively inhibited cell proliferation of hepatoma cells

After 24-h incubation, SAMC significantly reduced the cell viability of both Hep3B and Huh-7 cells in a dose-dependent manner. The dose of 250  $\mu\text{mol/L}$  showed intermediate inhibitory effects (Fig. 2A, B, and [Supplementary Information Fig. S1](#)). The cell damaging effects of SAMC mostly occurred in the first 24 h of treatment since the inhibitory results of both 24-h and 48-h incubation were similar ([Supplementary Information Fig. S1](#)). Thus, we used 250  $\mu\text{mol/L}$ /1 mmol/L SAMC and 24 h as the optimal *in vitro* treatment conditions.

For normal hepatocytes LO-2, only the higher dose of SAMC (1 mmol/L) reduced its viability from ~100% to ~87% (Fig. 2C). This observation was further confirmed by the release of lactate dehydrogenase (LDH) from these three kinds of cells (Fig. 2D). To study the effects of SAMC on cell proliferation, we stained cells with Ki-67, a cell proliferation marker, after 24-h incubation of SAMC. SAMC reduced the cell number and Ki-67 fluorescent signal density in a dose-dependent manner in both Hep3B and Huh-7 cells. However, both doses of SAMC did not significantly alter the LO-2 cell number (Fig. 2E). In addition, SAMC treatments down-regulated the protein expressions of proliferation markers—proliferating cell nuclear antigen (PCNA) and survivin, and up-regulated the protein levels of tumor suppressor genes *P53* and *P21* in Hep3B and Huh-7 cells (Fig. 2F)<sup>25</sup>. Accordingly, SAMC did not affect the protein expression of these markers in LO-2 cells. Furthermore, SAMC strongly disrupted the cell cycle distribution of both Hep3B and Huh-7 cells *via* reducing the S phase percentage but increasing the G0/G1 phase percentage ([Supplementary Information Table S3](#)). Collectively, we found that SAMC specifically inhibited the proliferation of hepatoma cells without significantly influencing the normal hepatocytes.

### 3.3. SAMC induced apoptosis of hepatoma cells through both intrinsic and extrinsic apoptotic pathways

To further investigate the anti-tumor ability of SAMC, we tested its apoptosis-inducing effects on Hep3B and Huh-7 cells. As expected, SAMC significantly increased the apoptotic ratio of both hepatoma cells in a dose-dependent manner, with a similar increasing pattern in the cellular activities of both caspase-3/7 and caspase-8 (Fig. 3A–C). Additionally, SAMC up-regulated the protein expressions of cleaved poly(ADP-ribose) polymerase, cytochrome *c*, BAX1, and FAS-associated protein with death domain (FADD) while down-regulating the BCL-2 protein level, indicating that SAMC caused hepatoma cell apoptosis through both the intrinsic and extrinsic apoptotic pathways (Fig. 3D). Although SAMC treatment slightly increased the apoptotic cell number and caspase-8 activity in LO-2, the apoptotic-inductive effects of SAMC on LO-2 were significantly less than those of hepatoma cells ([Supplementary Information Fig. S2](#)). To further investigate the apoptotic inducing mechanisms of



**Figure 1** LRP6 was frequently over-expressed in HCC. (A) The basal mRNA expression of *LRP6* was highest in Huh-7 HCC cell line, midst in Hep3B HCC cell line, and lowest in normal human hepatocyte cell line LO-2 ( $n=4$  for each cell line). (B) The *LRP6* mRNA level was significantly higher in human HCC cancerous tissue ( $n=48$ ) than that in non-cancerous tissue ( $n=48$ ) or liver tissue from healthy human ( $n=6$ ). (C) Representative Western blot results of LRP6 from human HCC cancerous tissue and their corresponding non-cancerous tissues, showing that the protein level of LRP6 was frequently higher in HCC tumor than non-tumor area. (D) Representative immunohistochemistry results of LRP6 in a human liver section showing both cancerous and non-cancerous areas (Scale bars, 20  $\mu\text{m}$ ). Data are presented in means  $\pm$  SEM.

SAMC, we then checked the expressional changes of c-Myc and myeloid cell leukemia-1 (MCL-1), key mediators of intrinsic and extrinsic apoptotic pathways, respectively<sup>26,27</sup>. Their protein expressions were dose-dependently inhibited by SAMC incubation in Huh-7 cell (Supplementary Information Fig. S3A). When their endogenous expressions were over-expressed by corresponding plasmids, the basal and SAMC-induced apoptosis levels were significantly suppressed, with inhibited activity of caspase-3/7/8 (Supplementary Information Fig. S3B–F). Furthermore, over-expression of c-Myc partly counter-acted the intrinsic apoptotic induction effect of SAMC through elevating the protein levels of both cytochrome *c* and BAX1 (Supplementary Information Fig. S3G).


#### 3.4. SAMC reduced the tumorigenic ability of hepatoma cells

To further investigate the effects of SAMC on the tumorigenic ability of hepatoma cells, we measured the changes of cellular migration and adhesion in both Hep3B and Huh-7 cells with or without SAMC incubation. We found that SAMC significantly reduced their migration ability (Fig. 4A). It is well known that cell adhesion is a vital step in cell migration process. Thus, as shown in Fig. 4B, SAMC decreased the cellular adhesion ability on fibronectin in both Hep3B and Huh-7 cells.

Since the modulation of the Wnt pathway evidently influences the tumorigenic processes of many types of tumor cells, we tested the protein level changes of key markers in this pathway. It was demonstrated that SAMC down-regulated the protein levels of  $\beta$ -catenin, phosphorylated LRP6, total LRP6, and DVL2 in both Hep3B and Huh-7 cells. Consistently, SAMC elevated DKK-1 (Dickkopf Wnt signaling pathway inhibitor 1) protein level in a dose-dependent manner (Fig. 4C). Then we found that treatments with both concentrations of SAMC led to significant decreases in TCF/ $\beta$ -catenin reporter activity up to 45% and 85% for Hep3B cells and 77% and 90% for Huh-7 cells, respectively (Fig. 4D).

#### 3.5. SAMC directly targeted LRP6 on the hepatoma cell membrane


To confirm the hypothesis that LRP6 is the direct acting target of SAMC, we performed SPR using recombinant human LRP6 protein and dissolved SAMC. SAMC bound to LRP6 tightly in a dose-dependent manner (from 25 to 400 nmol/L) with a dissociation constant of  $K_d=5.65 \times 10^{-6}$  mol/L (Fig. 5A). To further confirm this result and to investigate whether SAMC influences the LRP6 protein through the Wnt pathway of modulating its protein structure, we used TSA and found that SAMC treatment increased the melting temperature ( $T_m$ ) of LRP6



**Figure 2** SAMC inhibited human hepatoma cell proliferation but not normal hepatocytes *in vitro*. (A)–(C) SAMC significantly reduced cell viability of human hepatoma cell lines Hep3B and Huh-7 at 250 μmol/L and 1 mmol/L without evident influence in human normal hepatocyte cell line LO-2 ( $n=4$ ). (D) Release of LDH (lactate dehydrogenase) was significantly higher in Hep3B and Huh-7 than LO-2 after SAMC incubation ( $n=4$ ; Scale bars, 100 μm). (E) Representative immuno-fluorescent images of Ki-67 protein in three cell lines after SAMC incubation. (F) Representative Western blot showing change of protein levels of PCNA, survivin, p53 and p21 in three cell lines after SAMC incubation. Data are presented in means  $\pm$  SEM. \*\* $P<0.01$  and \*\*\* $P<0.001$ , respectively between indicated groups (in panel D means comparison with the LO-2 group).

for  $\sim 4^\circ\text{C}$  (Fig. 5B). In addition, we labeled SAMC with Alexa Fluor 488 TFP ester and examined the time-lapse distribution of SAMC after its addition to the culture medium. As shown in Fig. 5C, 30 or 60 min after the treatment, green-fluorescent-labeled SAMC was distributed around a number of Huh-7 cell membranes. Two hours later, the membrane-bound SAMC green signals were intensified and concentrated, with less unbound signals in the medium. It should be noted that in some cells, SAMC penetrated

into the cytoplasmic part, which was consistent with the finding that the subcellular localization of LRP6 could be modulated by external treatment (*e.g.*, Wnt) from the cell membrane to the cytoplasm<sup>28</sup>. Conversely, time-lapse experiments using LO-2 cells showed less SAMC-positive signals around cells and in the culture medium (Supplementary Information Fig. S4). Collectively, our data demonstrated that SAMC directly targeted LRP6 on the cell membrane of hepatoma cells.



**Figure 3** SAMC induced apoptosis through both apoptotic pathways in human hepatoma cell lines *in vitro*. (A) Change of apoptotic ratio after two doses of SAMC incubation in human hepatoma cell lines Hep3B and Huh-7 ( $n=4$ ). (B) Change of caspase-3/7 activity after SAMC incubation in Hep3B and Huh-7 ( $n=4$ ). (C) Change of caspase-8 activity after SAMC incubation in Hep3B and Huh-7 ( $n=4$ ). (D) Representative Western blot results of protein level change of cleaved PARP, total PARP, cytochrome *c* (cyto *c*), BCL-2, BAX1, and FADD after SAMC incubation in both Hep3B and Huh-7. Data are presented in means  $\pm$  SEM. \*\* $P < 0.01$  and \*\*\* $P < 0.001$ , respectively between indicated groups.

To prove that the potent anti-HCC effect and direct interaction with LRP6 is specific to SAMC, we tested the precursor substance of SAMC during garlic aging—*S*-allylcysteine (SAC)—for its anti-HCC properties and LRP6 interactions (Supplementary Information Fig. S5A). It was found that, unlike SAMC, 250  $\mu\text{mol/L}$  and 1  $\text{mmol/L}$  SAC could not effectively inhibit Hep3B and Huh7 proliferation and migration (Supplementary Information Fig. S5B and C). Moreover, the interactions between SAC and LRP6 was loose, with a dissociation constant of  $K_d = 8.78 \times 10^{-2}$  mol/L from SPR and  $\sim 0.5$  °C difference of melting temperature from TSA (Supplementary Information Fig. S5D). This result was in line with a previous report that at least 5  $\text{mmol/L}$  of SAC was needed to significantly inhibit HCC cell growth<sup>29</sup>.


### 3.6. Down-regulation of LRP6 in hepatoma cells attenuated Wnt pathway signaling and induced apoptosis

To further study the role of LRP6 on HCC development and SAMC-mediated anti-tumor effects, we then knocked-down the endogenous expression of LRP6 by using specific shRNA in Huh-7 cells, which showed relatively higher level of LRP6 than Hep3B (Figs. 1A and 6A). Cell viability and apoptosis were reduced and increased by the


knock-down of LRP6, respectively. Importantly, addition of SAMC further enhanced these effects of LRP6 knock-down on Huh-7 cells (Fig. 6B). In addition, deficiency of LRP6 itself significantly attenuated the migration ability of Huh-7, which was slightly strengthened by SAMC co-treatment (Fig. 6C). Free  $\beta$ -catenin pool and TCF/LEF reporter activity were significantly reduced when LRP6 was knocked-down in Huh-7. Addition of SAMC in the culture medium further enhanced such effects (Fig. 6D).

### 3.7. Over-expression of shRNA-resistant LRP6 rescued Wnt signaling and cancer cell growth impaired by SAMC

To confirm that the above observations on Huh-7 cells were specifically attributed to the knock-down of LRP6, we synthesized a shRNA-resistant LRP6 (LRP6 rescue) construct to repeat those experiments<sup>21</sup>. Transfection of LRP6 rescue restored the basal protein expression level down-regulated by LRP6 shRNA in Huh-7 cells (Supplementary Information Fig. S6A). It also rescued the cell viability and apoptosis modulated by the deficiency of LRP6 to a control comparable level (Supplementary Information Fig. S6B and C). More importantly, over-expression of LRP6 rescue markedly restored the TCF/LEF reporter activity of Wnt



**Figure 4** SAMC reduced human hepatoma cell tumorigenesis and Wnt signaling *in vitro*. (A) Change of migrated cell number after SAMC treatment, examined by Transwell assay, in human hepatoma cell lines Hep3B and Huh-7 ( $n=4$ ). (B) Change in percentage of adhered cells after SAMC treatment in Hep3B and Huh-7 ( $n=4$ ). (C) Representative Western blot results of protein level change of  $\beta$ -catenin, phosphorylated LRP6 at ser1490, total LRP6, DVL2, and DKK1 after SAMC incubation in both Hep3B and Huh-7. (D) Change of Wnt signaling target TCF/LEF activity after SAMC treatment in Hep3B and Huh-7 ( $n=4$ ). Data are presented in means  $\pm$  SEM. \* $P < 0.05$ , \*\* $P < 0.01$  and \*\*\* $P < 0.001$ , respectively between indicated groups (in panel C and D, \* and # indicate a  $P < 0.05$  when compared with 250  $\mu$ mol/L and 1 mmol/L groups, respectively).



**Figure 5** SAMC directly interacted with cell membrane receptor LRP6 *in vitro*. (A) SPR analysis of the binding of SAMC to immobilized recombinant human LRP6 protein. A gradient of SAMC was applied to quantify the binding affinity (25–400 nmol/L). The  $K_d$  (mol/L) value is  $5.65 \times 10^{-5}$ . (B) The interaction of LRP6 with SAMC was detected by thermal shift assay. Black line: basal LRP6 melting curve. Red line: LRP6+50  $\mu\text{mol/L}$  SAMC melting curve. (C) Representative fluorescent images of cultured Huh-7 cells after 0, 30, 60, and 120 min incubation of SAMC which was conjugated with Alexa 488 TFP ester dye (Scale bars, 50  $\mu\text{m}$ ).


pathway, with or without Wnt3a conditioned medium (CM, [Supplementary Information Fig. S6D](#)). It was interesting that when LRP6 was over-expressed in normal hepatocyte line LO-2, cells gained sensitivity to SAMC-induced viability reduction and

apoptosis induction. Additionally, the over-expression of LRP6 significantly increased the migration ability of LO-2, when examined using Matrigel, which could be counter-acted by SAMC ([Supplementary Information Fig. S7](#)).


### 3.8. Over-expression of LRP6 antagonized the anti-tumor effects of SAMC

Since deficiency of *LRP6* partly mimicked the anti-tumor abilities of SAMC treatment, it is interesting to find out whether over-expression of endogenous *LRP6* is able to enhance Wnt signaling in hepatoma cells (Fig. 6E). It was found that over-expression of *LRP6* increased the free  $\beta$ -catenin pool and TCF/LEF reporter activity which were re-balanced to the control level by the co-treatment with SAMC (Fig. 6F). In addition, the protein expression of key Wnt targets, including c-Myc, cyclin D1 and axin2, were up-regulated by *LRP6* over-expression and then down-regulated by

SAMC, indicating that LRP6 was important for the signal transduction of Wnt-mediated cell cycle regulation (Supplementary Information Fig. S8A). We found that LRP6 was important for the tumorigenesis of Hep3B since its over-expression increased its basal viability (Fig. 6F) and transwell ability (data not shown), which were further strengthened by the addition of Wnt3a and attenuated by SAMC. At the apoptotic level, *LRP6* over-expression down-regulated pro-apoptotic markers cytochrome c and BAX1, up-regulated the anti-apoptotic marker BCL-2, suggesting that LRP6 could stabilize cell from programmed cell death, although the change of apoptotic ratio was not very evident (Fig. 6F and Supplementary Information Fig. S8B).



**Figure 6** Deficiency and over-expression of *LRP6* positively and negatively influenced the anti-HCC properties of SAMC *in vitro*, respectively. (A) Verification of *LRP6* shRNA transfection by detecting its mRNA and protein level change before and after the transfection in Huh-7 hepatoma cell line *in vitro*. (B)–(C) When *LRP6* was knocked-down, the cell viability and migration was inhibited, which was further potentiated by 250  $\mu$ mol/L SAMC incubation. The apoptosis ratio change showed inverse pattern ( $n=4$ ). (D) *LRP6* down-regulation suppressed Wnt signaling examined by free  $\beta$ -catenin pull-down and TCF/LEF reporter assays in the absence and presence of Wnt3a ligands. Incubation of 250  $\mu$ mol/L SAMC further enhanced such effects ( $n=4$ ). (E) *LRP6* protein level change after transfection of *LRP6* plasmid. (F) Over-expression of *LRP6* significantly increased the Wnt signaling examined by free  $\beta$ -catenin pull-down and TCF/LEF reporter assays in the absence and presence of Wnt3a ligands, which was attenuated by 250  $\mu$ mol/L SAMC incubation ( $n=4$ ). Over-expression of *LRP6* slightly reduced the basal apoptotic ratio of Huh-7 cells, which were significantly reversed by 250  $\mu$ mol/L SAMC incubation ( $n=4$ ). Over-expression of *LRP6* significantly increased cell viability ability of Huh-7, which was re-balanced by SAMC treatment ( $n=4$ ). Data are presented in means  $\pm$  SEM. \* $P<0.05$ , \*\* $P<0.01$  and \*\*\* $P<0.001$ , respectively between indicated groups. Lys, lysates.



**Figure 7** Down-regulation of *LRP6* and SAMC treatment significantly inhibits HCC tumor growth *in vivo*. Huh-7 cells (pooled clones) stably expressing control or *LRP6* were subcutaneously injected at the dorsal region with  $1 \times 10^6/150 \mu\text{L}$  (low dose group) or  $4 \times 10^6/150 \mu\text{L}$  (high dose group). (A) Gross specimen anatomy of xenograft tumors. (B) Measurements of tumor volume for 33 days with or without SAMC treatment ( $n=5$ ). (C) Immunohistochemical analysis of LRP6 level in control and *LRP6*-KD xenograft tumors (with or without SAMC treatment) with anti-*LRP6* antibody (Abcam) (Scale bars, 20  $\mu\text{m}$ ). (D)–(E) Levels of LRP6 and Wnt target protein expressions (cyclin D1, c-Myc, and axin2) in control and *LRP6*-KD xenograft tumors detected by Western blot analysis. Western blot analysis shows a decrease in total and free  $\beta$ -catenin in *LRP6*-KD tumors, which were further reduced by SAMC treatment. Another orthotopic HCC model was established by the injection of  $2 \times 10^6$  wild-type or *LRP6* knock-down Huh-7 cells into the left liver lobe of nude mice. The animals were treated by oral administration of 300 mg/kg SAMC. (E) Survival of animals was monitored daily for 70 day. (F) intratumor expression of LRP6, c-Myc, cyclin D1, axin2 and free  $\beta$ -catenin were measured by Western blot when tumor tissue was collected at day 50. Data are presented in means  $\pm$  SEM. \*\* $P < 0.01$ , \*\*\* $P < 0.001$  compared with control group; # $P < 0.05$ , ## $P < 0.01$  compared with SAMC group.

### 3.9. LRP6 was vital for the tumor growth and SAMC treatment efficacy *in vivo*

To investigate whether SAMC is effective in suppressing the tumor growth and the role of LRP6 *in vivo*, we used a Huh-7 xenograft animal model (two concentrations,  $1 \times 10^6$  per injection as the low dose group and  $5 \times 10^6$  per injection as the high dose group) with or without a stable down-regulation of LRP6 (LRP6-KD). Strikingly, tumors in both Huh-7 concentrations derived from cells with stable down-regulation of LRP6 grew substantially slower than those derived from control cells as evaluated by tumor volume calculation (Fig. 7A and B). Daily oral gastric lavage feeding with 300 mg/kg SAMC further suppressed the tumor growth, when compared with corresponding control groups (Fig. 7A and B). Gross examination of tumors from each group at necropsy also showed that LRP6-KD tumors were significantly smaller than those of control tumors at both Huh-7 treated concentrations. Treatment with SAMC further reduced the tumor sizes (Fig. 7A). Consistent with the *in vitro* data, knockdown of endogenous LRP6 decreased the level of Wnt signaling and target gene expressions (c-Myc, cyclin D1, and axin2), which were further inhibited by SAMC administration (Fig. 7C and D). To further validate the role of LRP6 and the efficacy of SAMC, we used an orthotopic HCC model through the injection of  $2 \times 10^6$  Huh-7 cells. Treatment with SAMC significantly increased survival of tumor-bearing animals compared with animals receiving saline. As expected, knockdown of endogenous LRP6 showed similar rescue effects with SAMC consumption (Fig. 7E). The protein expressional changes of Wnt pathway members were also similar with the results from the s.c. model (Fig. 7F). Taken together, these results demonstrated that SAMC inhibited hepatoma s.c./orthotopic tumor growth partly through the LRP6/Wnt pathway. It should be noted that both 30-day and 60-day of SAMC treatment did not pose any evident toxicity to the mice liver, suggesting it to be an acceptable and bio-safe product when used as a food supplement (Supplementary Information Fig. S9).

## 4. Discussion

Aberrant Wnt activation is frequently found in ~50% of HCC patients<sup>30</sup>. As in other forms of cancer, dysregulation of this pathway can be broadly divided into two categories—ligand-dependent disorders (e.g., Wnt protein, LRP and Frizzled receptors) and ligand-independent disorders (e.g., mutations of axin,  $\beta$ -catenin and TCF)<sup>31</sup>. Since receptor–drug interaction is an important basis for drug screening and discovery, the role of Wnt pathway receptors in cancer progression and drug therapy received much attention recently. For example, Frizzled3, -6 and -7 were found to be overexpressed in HCC<sup>32,33</sup>. Additional study confirmed that activated Wnt3/Frizzled7 could promote early carcinogenic process through promoting the acquisition of malignant phenotypes<sup>33</sup>. Thus, a soluble ectodomain of Frizzled7 was developed to inhibit the Wnt signaling, which sensitized HCC cells to chemotherapy drugs (e.g., doxorubicin)<sup>34</sup>. In the current study, we identified that LRP6 was frequently up-regulated in human HCC samples and cell lines, when compared with non-tumor liver tissue and normal hepatocytes, respectively. This finding was consistent with a pilot study showing that LRP6 over-expression in HCC promoted cell proliferation, cell migration and hyperactivation of the Wnt signaling<sup>10</sup>.

SAMC was found to be a tumor suppressor in several kinds of tumors. Although the “down-stream” anti-tumor events, such as promotion of apoptosis, inhibition of proliferation and cell cycle, and suppression of tumorigenesis, were entirely or partially reported in these studies, and there was no study that addressed these two basic questions: (1) What is the immediate molecule(s) that mediate(s) the anti-tumor signaling from SAMC?; and (2) Can SAMC specifically antagonize tumor cell proliferation? This study not only found that LRP6 was over-expressed in HCC cells, but we also identified the direct interaction between SAMC, but not its natural precursor SAC, and LRP6 *in vitro*, through the experiments done on SPR, TSA and time-lapse fluorescent imaging (Fig. 5 and Supplementary Information Figs. S4 and 5). More importantly, our results demonstrated that knockdown of LRP6 was capable of inhibiting hepatoma cell proliferation, tumorigenesis, Wnt signaling and inducing apoptosis, which could be further potentiated by the co-treatment with SAMC both in *in vitro* and *in vivo* models. These findings strongly supported our initial concepts about the crucial role of LRP6 in HCC progression and the receptor signaling used by SAMC. Consistently, when LRP6 was further over-expressed in hepatoma cells, the basal tumorigenic ability and Wnt signaling were elevated, which could also be attenuated by SAMC co-treatment (Figs. 6 and 7). Furthermore, when LRP6 was over-expressed in normal hepatocytes (LO-2), it enhanced proliferative and tumorigenic phenotypes, which could be reduced by SAMC treatment (Supplementary Information Fig. S7). Considering that after SAMC treatment, the change of LO-2 proliferation, apoptosis and survival markers (surviving and P53/P21) was not evident (Fig. 2 and Supplementary Information Fig. S2), these results were vitally important because they suggested that LRP6 might be one of the molecular pathways for the anti-tumor specificity of SAMC in HCC therapy.

Since basic normal Wnt signaling is indispensable for several physiological functions, an ideal LRP6/Wnt inhibitor should significantly reduce its signaling to inhibit the tumor growth without influencing the physiological homeostasis. We demonstrated in the current study that SAMC treatment in healthy mice did not influence normal liver histology nor caused any evident adverse effect (Supplementary Information Fig. S9). In our previous studies, the hepato-protective roles and possible toxicity of SAMC in several liver diseases have been mechanistically investigated<sup>12,13</sup>. Thus, our work introduces SAMC as a promising nutraceutical antitumor agent that can be further developed for HCC targeted therapy, since there is virtually no treatment for advanced HCC.

## Acknowledgments

Authors thank the contribution of Wakunaga Pharmaceutical Co., Ltd., Japan, for the free supply of pure form of SAMC and SAC. This study was supported by Foundation of Pearl River Science and Technology New Star (grant number 201506010087) and Basic Research Fund of Shenzhen City (JCYJ2015040211430633) to Jia Xiao Health Medical Research Fund (HMRF, No. 12133881), General Research Fund and Small Project Funding, University Research Committee, The University of Hong Kong, to George L. Tipoe National Natural Science Foundation of China (No. 81570552) to National Program on Key Basic Research Project of China (973 Program, 2014CB542205), Funds of Leading Talents of Guangdong (2013), Programme of Introducing Talents of Discipline to Universities (B14036) to Kwok-Fai So and National Health and Medical Research Council (1031221 & 1031228) to Ming-Tat Ling

## Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.apsb.2017.10.003>.

## References

- Theise ND, Chen CJ, Kew MC. Liver cancer. In: Stewart B, Wild C, editors. *World Cancer Report 2014*. Lyon: International Agency for Research on Cancer; 2014. p. 577–93.
- Hong Y, Huang J. Autoantibodies against tumor-associated antigens for detection of hepatocellular carcinoma. *World J Hepatol* 2015;**7**:1581–5.
- Monga SP.  $\beta$ -Catenin signaling and roles in liver homeostasis, injury, and tumorigenesis. *Gastroenterology* 2015;**148**:1294–310.
- Gough NR. Focus issue: wnt and  $\beta$ -catenin signaling in development and disease. *Sci Signal* 2012;**5**:eg2.
- Monga SP. Role and regulation of  $\beta$ -catenin signaling during physiological liver growth. *Gene Expr* 2014;**16**:51–62.
- Pez F, Lopez A, Kim M, Wands JR, Caron de Fromental C, Merle P. Wnt signaling and hepatocarcinogenesis: molecular targets for the development of innovative anticancer drugs. *J Hepatol* 2013;**59**:1107–17.
- Bogaerts E, Heindryckx F, Vandewynckel YP, van Grunsven LA, van Vlierberghe H. The roles of transforming growth factor- $\beta$ , Wnt, Notch and hypoxia on liver progenitor cells in primary liver tumours. *Int J Oncol* 2014;**44**:1015–22.
- Mercer KE, Hennings L, Ronis MJ. Alcohol consumption, Wnt/ $\beta$ -catenin signaling, and hepatocarcinogenesis. *Adv Exp Med Biol* 2015;**815**:185–95.
- Go GW. Low-density lipoprotein receptor-related protein 6 (LRP6) is a novel nutritional therapeutic target for hyperlipidemia, non-alcoholic fatty liver disease, and atherosclerosis. *Nutrients* 2015;**7**:4453–64.
- Tung EK, Wong BY, Yau TO, Ng IO. Upregulation of the Wnt co-receptor LRP6 promotes hepatocarcinogenesis and enhances cell invasion. *PLoS One* 2012;**7**:e36565.
- Nicastro HL, Ross SA, Milner JA. Garlic and onions: their cancer prevention properties. *Cancer Prev Res* 2015;**8**:181–9.
- Xiao J, Liong EC, Ling MT, Ching YP, Fung ML, Tipoe GL. S-Allylmercaptocysteine reduces carbon tetrachloride-induced hepatic oxidative stress and necroinflammation via nuclear factor  $\kappa$ B-dependent pathways in mice. *Eur J Nutr* 2012;**51**:323–33.
- Xiao J, Guo R, Fung ML, Liong EC, Chang RC, Ching YP, et al. Garlic-derived S-allylmercaptocysteine ameliorates nonalcoholic fatty liver disease in a rat model through inhibition of apoptosis and enhancing autophagy. *Evid Based Complement Altern Med* 2013;**2013**:642920.
- Xiao D, Pinto JT, Soh JW, Deguchi A, Gundersen GG, Palazzo AF, et al. Induction of apoptosis by the garlic-derived compound S-allylmercaptocysteine (SAMC) is associated with microtubule depolymerization and c-Jun NH<sub>2</sub>-terminal kinase 1 activation. *Cancer Res* 2003;**63**:6825–37.
- Chu Q, Ling MT, Feng H, Cheung HW, Tsao SW, Wang X, et al. A novel anticancer effect of garlic derivatives: inhibition of cancer cell invasion through restoration of E-cadherin expression. *Carcinogenesis* 2006;**27**:2180–9.
- Howard EW, Ling MT, Chua CW, Cheung HW, Wang X, Wong YC. Garlic-derived S-allylmercaptocysteine is a novel *in vivo* antimetastatic agent for androgen-independent prostate cancer. *Clin Cancer Res* 2007;**13**:1847–56.
- Hu H, Zhang XP, Wang YL, et al. Identification of a novel function of Id-1 in mediating the anticancer responses of SAMC, a water-soluble garlic derivative, in human bladder cancer cells. *Mol Med Rep* 2011;**4**:9–16.
- Zhang H, Wang K, Lin G, Chua CW, Luk SU, Wong YC, et al. Antitumor mechanisms of S-allyl mercaptocysteine for breast cancer therapy. *BMC Complement Altern Med* 2014;**14**:270.
- Lee Y, Kim H, Lee J, Kim K. Anticancer activity of S-allylmercaptocysteine on implanted tumor of human gastric cancer cell. *Biol Pharm Bull* 2011;**34**:677–81.
- Tong D, Qu H, Meng X, Jiang Y, Liu D, Ye S, et al. S-Allylmercaptocysteine promotes MAPK inhibitor-induced apoptosis by activating the TGF- $\beta$  signaling pathway in cancer cells. *Oncol Rep* 2014;**32**:1124–32.
- Liu CC, Prior J, Piwnica-Worms D, Bu G. LRP6 overexpression defines a class of breast cancer subtype and is a target for therapy. *Proc Natl Acad Sci U S A* 2010;**107**:5136–41.
- Ouyang S, Song X, Wang Y, Ru H, Shaw N, Jiang Y, et al. Structural analysis of the STING adaptor protein reveals a hydrophobic dimer interface and mode of cyclic di-GMP binding. *Immunity* 2012;**36**:1073–86.
- Llovet JM, Bustamante J, Castells A, Vilana R, Ayuso Mdel C, Sala M, et al. Natural history of untreated nonsurgical hepatocellular carcinoma: rationale for the design and evaluation of therapeutic trials. *Hepatology* 1999;**29**:62–7.
- Hu J, Dong A, Fernandez-Ruiz, Shan J, Kawa M, Martínez-Ansó E, et al. Blockade of Wnt signaling inhibits angiogenesis and tumor growth in hepatocellular carcinoma. *Cancer Res* 2009;**69**:6951–9.
- Liang X, Da M, Zhuang Z, Wu W, Wu Z, Wu Y, et al. Effects of Survivin on cell proliferation and apoptosis in MG-63 cells *in vitro*. *Cell Biol Int* 2009;**33**:119–24.
- Ladu S, Calvisi D, Conner F, Farina M, Factor VM, Thorgeirsson SS. E2F1 inhibits c-Myc-driven apoptosis via PIK3CA/Akt/mTOR and COX-2 in a mouse model of human liver cancer. *Gastroenterology* 2008;**135**:1322–32.
- Thomas LW, Lam C, Edwards SW. Mcl-1; the molecular regulation of protein function. *FEBS Lett* 2010;**584**:2981–9.
- Kagermeier-Schenk B, Wehner D, Özhan-Kizil G, Yamamoto H, Li J, Kirchner K, et al. Waif1/ST4 inhibits Wnt/ $\beta$ -catenin signaling and activates noncanonical Wnt pathways by modifying LRP6 subcellular localization. *Dev Cell* 2011;**21**:1129–43.
- Ng KT, Guo DY, Cheng Q, Geng W, Ling CC, Li CX, et al. A garlic derivative, S-allylcysteine (SAC), suppresses proliferation and metastasis of hepatocellular carcinoma. *PLoS One* 2012;**7**:e31665.
- Lachenmayer A, Alsinet C, Savic R, Cabellos L, Toffanin S, Hoshida Y, et al. Wnt-pathway activation in two molecular classes of hepatocellular carcinoma and experimental modulation by sorafenib. *Clin Cancer Res* 2012;**18**:4997–5007.
- Klaus A, Birchmeier W. Wnt signalling and its impact on development and cancer. *Nat Rev Cancer* 2008;**8**:387–98.
- Bengochea A, de Souza MM, Lefrancois L, Le Roux E, Galy O, Chemin I, et al. Common dysregulation of Wnt/Frizzled receptor elements in human hepatocellular carcinoma. *Br J Cancer* 2008;**99**:143–50.
- Nambotin SB, Tomimaru Y, Merle P, Wands JR, Kim M. Functional consequences of WNT3/Frizzled7-mediated signaling in non-transformed hepatic cells. *Oncogenesis* 2012;**1**:e31.
- Wei W, Chua MS, Grepper S, So SK. Soluble Frizzled-7 receptor inhibits Wnt signaling and sensitizes hepatocellular carcinoma cells towards doxorubicin. *Mol Cancer* 2011;**10**:16.