

1    **A Stilbene Synthase Gene (*SbSTS1*) is Involved in Host and Non-host Defense**

2    **Responses in *Sorghum bicolor*<sup>1</sup>**

3    Christine K.Y. Yu<sup>2</sup>, Karin Springob, Jürgen Schmidt, Ralph L Nicholson, Ivan K Chu,

4    Wing Kin Yip, and Clive Lo\*

5

6    Departments of Botany (C.K.Y.Y., W.K.Y., C.L.) and Chemistry (IKC), The

7    University of Hong Kong, Pokfulam Road, Hong Kong, China; Leibniz-Institut für

8    Pflanzenbiochemie (IPB), Weinberg 3, 06120 Halle/Saale, Germany (K.S., J.S.);

9    Department of Botany and Plant Pathology, Purdue University, West Lafayette, IN

10    47907 (RLN)

11

12    <sup>1</sup>This work was supported by a grant from the Research Grants Council of the Hong

13    Kong Special Administrative Region China (HKU 7349/03M).

14

15    <sup>2</sup>C.K.Y.Y. was supported by a Research Postgraduate Studentship from The University

16    of Hong Kong.

17

18    \* Corresponding author; e-mail clivelo@hkucc.hku.hk; fax +852-2858-3477.

19

1    **ABSTRACT**

2    A chalcone synthase (CHS)-like gene, *SbCHS8*, with high expressed sequence tag  
3    abundance in a pathogen-induced cDNA library was identified previously in sorghum.  
4    Genomic Southern analysis revealed that *SbCHS8* represents a single copy gene.  
5    *SbCHS8* expression was induced in sorghum mesocotyls following inoculation with  
6    *Cochliobolus heterotrophus* and *Colletotrichum sublineolum*, corresponding to  
7    non-host and host defense responses, respectively. However, the induction was  
8    delayed by approximately 24 h when compared to the expression of at least one of the  
9    other *SbCHS* genes. In addition, *SbCHS8* expression was not induced by light and  
10   did not occur in a tissue-specific manner. *SbCHS8*, together with *SbCHS2*, was  
11   over-expressed in transgenic *Arabidopsis tt4* mutants defective in CHS activities.  
12   *SbCHS2* rescued the ability of these mutants to accumulate flavonoids in seed coats  
13   and seedlings. In contrast, *SbCHS8* failed to complement the mutation, suggesting  
14   that the encoded enzyme does not function as a CHS. To elucidate their biochemical  
15   functions, recombinant proteins were assayed with different phenylpropanoid-CoA  
16   esters. Flavanones and stilbenes were detected in the reaction products of *SbCHS2*  
17   and *SbCHS8*, respectively. Taken together, our data demonstrated that *SbCHS2*  
18   encodes a typical CHS that synthesizes naringenin chalcone necessary for the  
19   formation of different flavonoid metabolites. On the other hand, *SbCHS8*, now

1 re-termed as *SbSTS1*, encodes an enzyme with stilbene synthase activities, suggesting  
2 that sorghum accumulates stilbene-derived defense metabolites in addition to the  
3 well-characterized 3-deoxyanthocyanidin phytoalexins.

## 1 INTRODUCTION

2 Sorghum (*Sorghum bicolor* L.) is well known for its adaptability to adverse  
3 environments such as hot and dry conditions. The plant is also a rich source of  
4 distinct natural products. For example, sorghum seedlings accumulate high levels of  
5 dhurrin, a cyanogenic glycoside derived from tyrosine (Busk and Møller, 2002). To  
6 preclude competition for resources, sorghum roots exude sorgoleone and derivatives,  
7 a group of hydrophobic *p*-benzoquinone compounds, which inhibit electron transfer in  
8 PSII (Czarnota et al., 2001). In response to pathogen infection, sorghum synthesizes  
9 a unique class of flavonoid phytoalexins, the 3-deoxyanthocyanidins, as an essential  
10 component in the plant's active defense mechanisms (Lo et al., 1999).

11

12 Chalcone synthase (CHS) catalyzes the first committed step in flavonoid  
13 biosynthesis. The enzyme is the prototype of the plant type III polyketide synthase  
14 (PKS) family including the closely related stilbene synthases (STSs), pyrone  
15 synthases, acridone synthases, valerophenone synthases, and benzalacetone synthases  
16 (Springob et al., 2003), giving rise to the diversity of type III PKS-derived  
17 phytochemicals throughout the plant kingdom (Austin and Noel, 2003). Particularly  
18 interesting is the STS enzymes which utilize the same starter phenylpropanoid-CoA  
19 esters as the CHS enzymes and perform three condensations with malonyl-CoA

1 generating a common tetraketide intermediate, but result in the formation of the  
2 stilbene backbone following a completely different cyclization mechanism (Fig. 1).  
3 In different public databases, hundreds of plant DNA sequences are annotated as CHS  
4 genes based on sequence homology. However, these PKS genes may in fact have  
5 different metabolic roles, such as stilbene-forming activities, which can only be  
6 uncovered by experimental characterizations (Springob et al., 2003).

7

8 We have previously described a family of eight CHS genes, *SbCHS1-SbCHS8*, in  
9 sorghum (Lo et al., 2002). *SbCHS1* to *SbCHS7* (AF152548-AF152554) are highly  
10 conserved (at least 97.5% sequence identity at amino acid level) and closely related to  
11 the maize *C2* and *Whp* genes encoding CHS enzymes. *SbCHS8* (AY069951), on the  
12 other hand, is only 81-82% identical to *SbCHS1-SbCHS7* at amino acid level and  
13 appears to be more distantly related as revealed by phylogenetic analysis (Lo et al.,  
14 2002). These findings suggested that *SbCHS8* was duplicated from the ancestral  
15 form of *SbCHS1-SbCHS7* and diverged in protein coding sequence. In *in silico*  
16 analysis, *SbCHS8* was found to have significantly higher expressed sequence tag  
17 (EST) abundance in a pathogen-induced library (Lo et al., 2002). This EST library  
18 was prepared from 2-week-old seedlings 48 h after inoculation with the anthracnose  
19 pathogen *Colletotrichum sublineolum* (University of Georgia). Accumulation of

1 3-deoxyanthocyanidin was consistently detected in sorghum tissues inoculated with  
2 this fungal pathogen (Lo et al., 1999; Snyder and Nicholson, 1990), leading to our  
3 speculation that *SbCHS8* is involved in the biosynthesis pathway (Lo et al., 2002).

4

5 In this study, we used the well-established mesocotyl inoculation system  
6 (Hipskind et al., 1996; Lo and Nicholson, 1998) to investigate *SbCHS8* gene  
7 expression in sorghum. In addition, we attempted to define the biochemical  
8 functions of the encoded protein through analysis of transgenic *Arabidopsis* flavonoid  
9 mutants and *in vitro* activity assays of recombinant proteins. Our data demonstrate  
10 that *SbCHS8*, in fact, encodes a STS enzyme and gene expression was activated  
11 during host and non-host defense responses. Possible metabolites derived from the  
12 activity of the sorghum STS enzyme are discussed.

13

## 14 **RESULTS**

### 15 **Genomic Southern analysis of *SbCHS* genes**

16 For genomic Southern analysis, total DNA samples from 3 different sorghum  
17 cultivars (BTx623, Sc748-5, and DK46) were digested to completion with *EcoR* I,  
18 *Hind* III, or *Xba* I. A *SbCHS8*-specific PCR fragment containing part of the coding  
19 sequence and 3'-untranslated (UTR) region was used as a hybridization probe.

1 Results indicated that *SbCHS8* is a single copy gene and there are no RFLPs among  
2 the different cultivars examined (Fig. 2A). In contrast, a number of signals with  
3 varying intensities and sizes were detected when the digested DNA samples were  
4 hybridized with a CHS universal probe (Fig. 2B), which was derived from a  
5 conserved region in the *SbCHS1-SbCHS7* coding sequences. RFLPs were observed  
6 among the different cultivars following *Hind* III digestion. For example, Sc748-5  
7 displayed a hybridization pattern distinct from the other two cultivars (Fig. 2B).

8

9 **Northern analysis of *SbCHS* gene expression**

10 Sorghum cultivar DK46 accumulates anthocyanin pigments in mesocotyls of  
11 etiolated seedlings upon light induction (Lo and Nicholson, 1998). Total RNA  
12 samples were prepared from mesocotyl tissue at various time points following light  
13 exposure. Northern analysis revealed that *SbCHS8* gene expression was not  
14 inducible by light (Fig. 3A). In contrast, expression of at least one of the  
15 *SbCHS1-SbCHS7* genes was detected when the universal probe was used in the  
16 hybridizations. These data indicated that *SbCHS8* is not involved in the  
17 light-induced anthocyanin biosynthesis pathway. The expression of *SbCHS8* was  
18 then investigated in different sorghum tissues. RNA samples were collected from  
19 roots and leaves of 6-d-old etiolated seedlings and 1-month old plants, as well as

1 developing panicles. As shown in Fig 3B, *SbCHS8* transcripts were not detectable in  
2 any of these tissues during normal growth conditions, indicating that the gene is not  
3 expressed in a tissue-specific manner.

4

5 To study the expression of *SbCHS8* during defense responses, etiolated seedlings

6 of DK46 were inoculated with either *Cochliobolus heterostrophus*, a maize pathogen

7 but nonpathogenic to sorghum, or *Colletotrichum sublineolum*, the causal agent of

8 sorghum anthracnose. The inoculated seedlings were either kept in the dark or

9 placed under constant light. Total RNA samples from various time points were

10 analyzed by northern hybridizations. Transcripts of *SbCHS* genes, including

11 *SbCHS8*, were detected in all the inoculation conditions examined (Fig. 3C-H).

12 However, pathogen-induced accumulation of *SbCHS8* transcripts was delayed

13 compared to transcripts detected by the universal *SbCHS* probe. For example,

14 transcripts of *SbCHS8* were not detected until 24 h after inoculation with *C.*

15 *heterotrophus* under dark conditions while transcripts of at least one of the other

16 *SbCHS* genes were detected within 3 h (Fig. 3D). Similarly, *SbCHS8* gene

17 expression was not observed until 72 h after inoculation with *C. sublineolum* under

18 dark conditions while the expression of at least one of the other *SbCHS* genes was

19 observed within 36 h (Fig. 3F). Although *SbCHS8* is not light inducible, the

1 pathogen-induced gene expression appeared to be enhanced under light. Thus,  
2 transcripts of *SbCHS8* were detected 12 h earlier in *C. heterotrophus*-inoculated plants  
3 and 24 h earlier in *C. sublineolum*-inoculated plants under light compared to the  
4 respective infected plants kept in the dark (Fig. 3C-F).

5

6 The expression of *SbCHS8* was also examined in two sorghum inbred lines,  
7 BTx623 and Sc748-5, with differential physiological and biochemical responses to  
8 the anthracnose pathogen *C. sublineolum* (Lo et al., 1999). Transcripts of *SbCHS8*  
9 were detected in Sc748-5 (resistant) plants with an accumulation pattern (Fig. 3G)  
10 similar to that observed in DK46 plants after inoculation with *C. sublineolum* (Fig.  
11 3E). In contrast, *SbCHS8* transcript accumulation was delayed and less intense in  
12 the inoculated BTx623 (susceptible) plants (Fig. 3H). On the other hand, the  
13 patterns of the accumulation of *SbCHS* transcripts detected by the universal probe  
14 were similar in both cultivars following fungal inoculation (Fig. 3G-H).

15

#### 16 **Transgenic analysis of *Arabidopsis tt4* mutants**

17 The complementation of *Arabidopsis* transparent testa (*tt*) mutants by maize  
18 genes demonstrated the convenience of this system to establish the function of  
19 uncharacterized coding sequences with homology to flavonoid structural genes (Dong

1 et al., 2001). *Arabidopsis tt4* mutants are deficient in CHS activities, resulting in the  
2 absence of flavonoid-derived metabolites in different tissues. *SbCHS2* and *SbCHS8*  
3 genes were expressed under the control of the cauliflower mosaic virus (CaMV) 35S  
4 promoter in *tt4* plants. *SbCHS2* was selected as a representative of the highly  
5 conserved *SbCHS1-SbCHS7* genes. Expression of the sorghum genes in transgenic  
6 *tt4* mutants was confirmed by northern analysis in 10 to 14-day-old T1 seedlings (data  
7 not shown). Three independent lines with strong expression for each transgene were  
8 selected for phenotypic studies.

9

10 Transgenic *tt4* mutants expressing *SbCHS2* produced T1 seeds with brown  
11 pigmentation characteristic of wild type seeds (Fig. 4A), indicating the accumulation  
12 of tannins in seed coats. In addition, these transgenic seedlings showed anthocyanin  
13 pigments in cotyledons and hypocotyls when germinated in medium devoid of  
14 nitrogen sources, a sensitive condition previously employed to induce the anthocyanin  
15 biosynthesis pathway in *Arabidopsis* (Dong et al., 2001; Hsieh et al., 1998). In  
16 contrast, seed coats of *SbCHS8*-expressing *tt4* plants remained yellow in both T1 and  
17 T2 generations and the transgenic seedlings failed to accumulate anthocyanin under  
18 nitrogen deficiency (Fig. 4A). These results demonstrated that *SbCHS2* was able to  
19 fully complement the *tt4* mutation in *Arabidopsis* and hence the gene product is a

1 functional CHS enzyme. In contrast, *SbCHS8* does not encode CHS that could  
2 otherwise rescue the deficiencies in flavonoid biosynthesis in the *tt4* mutants.

3

4 To further characterize the flavonoids synthesized by the transgenic  
5 *Arabidopsis tt4* mutants, HPLC experiments were performed using acid hydrolyzed  
6 methanol extracts prepared from 14-d-old seedlings. Expression of *SbCHS2* in  
7 transgenic *tt4* plants resulted in the accumulation of the flavonols quercetin and  
8 kaempferol which were not present in the extracts prepared from non-transformed  
9 mutants (Fig. 4B). The flavonoid profile, monitored at  $A_{360}$ , of these transgenic  
10 plants was near identical to that of the wild type plant, Landsberg *erecta* (*Ler*),  
11 confirming the complete complementation of *tt4* mutation by *SbCHS2*. In contrast,  
12 accumulation of these flavonols was not detected in the *SbCHS8*-expressing *tt4* plants,  
13 further suggesting that this sorghum enzyme does not function as a CHS *in planta*.

14

15 **Biochemical analysis of SbCHS recombinant proteins**

16 *SbCHS2* and *SbCHS8* were over-expressed in *E. coli* and purified by  
17 immobilized metal affinity chromatography to generate electrophoretically  
18 homogenous recombinant proteins (data not shown). Purified protein samples were  
19 incubated with  $^{14}\text{C}$ -malonyl CoA and different phenylpropanoid-CoA esters.

1 Recombinant proteins of *Cassia alata* CHS (CalCHS1; Samappito et al., 2002) and  
2 *Rheum tataricum* STS (RtSTS1; Samappito et al., 2003) were included as reference  
3 enzymes in our assays. The resulting radioactive products were resolved by  
4 reversed-phase thin layer chromatography (RP-TLC). With cinnamoyl-CoA and  
5 *p*-coumaroyl-CoA as start substrates, the radiolabeled RP-TLC profiles of the  
6 SbCHS2 reaction were the same as those of CalCHS1 (Fig. 5A). Surprisingly, the  
7 SbCHS8 reaction profiles were almost identical to those of RtSTS1 (Fig. 5A).  
8 SbCHS2 and SbCHS8 assays resulted in the production of flavanones (pinocembrin  
9 and naringenin) and stilbenes (pinosylvin and resveratrol), respectively. Flavanones  
10 were presumably detected due to spontaneous isomerization of the respective  
11 chalcones. *Bis*-noryangonin (BNY)-type and *p*-coumaroyltriacyclic acid lactone  
12 (CTAL)-type derailed pyrone byproducts were also identified in most of the assays  
13 (Fig. 5A). In addition, SbCHS8 was found to produce small amounts of flavanones  
14 (Fig. 5B, pinocembrin to pinosylvin ratio = 5.5: 100; naringenin to resveratrol ratio=  
15 2.0: 100). The CHS side activity of SbCHS8 was lower than that of RtSTS1 as  
16 reflected from their product ratios. Cross-reaction between CHS and STS enzymes  
17 has been demonstrated in *in vitro* reactions previously (Samappito et al 2002, 2003;  
18 Yamaguchi et al, 1999). Similarly, trace levels of pinosylvin were detected in the  
19 SbCHS2 and CalCHS1 assays with cinnamoyl-CoA while no resveratrol was detected

1 with *p*-coumaroyl-CoA (Fig. 5B). We also used caffeoyl-CoA and feruloyl-CoA in  
2 the assays but the BNY-type and CTAL-type pyrones were formed predominantly  
3 (data not shown), suggesting that these starter-CoAs are not physiologically relevant  
4 substrates (Samappito et al., 2003).

5 To unambiguously identify the reaction products, recombinant proteins were  
6 incubated with unlabeled malonyl-CoA and starter CoA esters in scaled-up reactions.  
7 The product mixtures obtained in these experiments were analyzed by combined  
8 LC/electrospray ionization (ESI)-MS/MS in selected reaction monitoring (SRM)  
9 mode using the reactions leading to key ions. Under positive ESI conditions,  
10 flavanones were detected by reactions leading to a key ion at *m/z* 153  
11 (trihydroxybenzoyl moiety) as well as the respective phenylpropanoyl cations:  
12 cinnamoyl cation at *m/z* 131 and *p*-coumaroyl cation at *m/z* 147 (Fig. 5C; Samappito  
13 et al., 2002). The mass spectral behavior of stilbenes under negative ESI conditions  
14 is characterized by the loss of ketene units. Resveratrol was confirmed by measuring  
15 the two key reactions *m/z* 227 [M-H]<sup>-</sup> to *m/z* 185 [M-H-CH<sub>2</sub>CO]<sup>-</sup> and *m/z* 227 [M-H]<sup>-</sup>  
16 to *m/z* 143 [M-H-2CH<sub>2</sub>CO]<sup>-</sup>, respectively (Fig. 5C, Stecher et al., 2001; Samappito et  
17 al., 2003). Similarly, pinosylvin was identified by the reaction *m/z* 211 [M-H]<sup>-</sup> to  
18 *m/z* 169 [M-H-CH<sub>2</sub>CO]<sup>-</sup> (Fig. 5C). Taken together, our results clearly demonstrated  
19 that SbCHS8 encodes an enzyme with STS activities.

1

2 **DISCUSSION**

3 *SbCHS8* was initially annotated as a CHS-like gene having high EST abundance  
4 in a cDNA library prepared from infected sorghum plants with the accumulation of  
5 3-deoxyanthocyanidin phytoalexins (Lo et al., 2002). However, we demonstrated  
6 *SbCHS8* is not involved in flavonoid biosynthesis *in planta* as it failed to complement  
7 the *tt4* mutation in *Arabidopsis* (Fig. 4). Instead, the recombinant *SbCHS8* protein  
8 synthesized pinosylvin and resveratrol as major products *in vitro* using  
9 cinnamoyl-CoA and *p*-coumaroyl-CoA as starter molecules, respectively (Fig. 5) and  
10 the sorghum gene was, therefore, re-termed as *SbSTS1*.

11

12 To our knowledge, *SbSTS1* represents the first example of a STS gene in  
13 monocots. The gene is not constitutively expressed, but inducible following fungal  
14 inoculation. Related enzymes performing STS-like cyclizations, e.g. bibenzyl  
15 synthases, have been isolated from a *Phalaenopsis* orchid (Preisig-Muller et al., 1995).  
16 STS enzymes occur only in limited numbers of unrelated plant species. Resveratrol  
17 STS enzymes were originally described in grapes and peanuts, which accumulate  
18 elevated levels of the stilbene following pathogen inoculations and elicitor treatments  
19 (Schröder et al., 1988; Wiese et al., 1994). Recently, a root-specific STS cDNA was

1 reported in *Rheum*, a medicinal plant in the Polygonaceae family, with resveratrol  
2 accumulation in roots (Samappito et al. 2003). A second category of STS,  
3 pinosylvin STS enzymes, is largely associated with pine trees. These enzymes  
4 utilize cinnamoyl-CoA as the starter ester to synthesize pinosylvin which is found in  
5 the heartwood or serving as phytoalexins in sapwoods and needles (Preisig-Müller et  
6 al., 1999). In *Psilotum nudum*, two pinosylvin STS enzymes were recently identified  
7 through *in vitro* enzyme assays of the recombinant proteins, although stilbenes or their  
8 derivatives have not been isolated from this primitive vascular plant (Yamazaki et al.,  
9 2001).

10

11 The expression of STS genes is often induced by a variety of abiotic and biotic  
12 stresses, such as elicitor treatment, pathogen inoculation, wounding, UV irradiation,  
13 and post-harvest wilting procedures (Preisig-Müller et al., 1999; Verari et al., 2001).  
14 Constitutive expression of STS genes was described in young seedlings of grapes,  
15 presumably representing a pre-existing defense mechanism (Sparvoli et al, 1994). In  
16 sorghum, *SbSTS1* gene expression was not detected under non-induced conditions in  
17 all tissues examined (Fig. 3). Our results also revealed that *SbSTS1* is a late  
18 component during both non-host (against *C. heterotrophus*) and host (against *C.*  
19 *sublineolum*) defense responses, comparing to the expression of at least one of the

1 *SbCHS* genes (Fig. 3). In inoculated plants kept in the dark, a condition in which  
2 flavonoid metabolism was not induced by light, transcripts of *SbSTS1* gene were not  
3 detected until 24 h or 72 h after inoculation with *C. heterotrophus* or *C. sublineolum*,  
4 respectively (Fig.3D, F), during which significant amounts of 3-deoxyanthocyanidins  
5 had accumulated (data not shown). The late induction of *SbSTS1* expression  
6 provided further evidence that the enzyme is not involved in the biosynthesis of  
7 3-deoxyanthocyanidins in sorghum. Nevertheless, *SbSTS1* and *SbCHS* genes are  
8 components involved in both non-host and host defense responses. Interestingly,  
9 earlier and stronger induction of the *SbSTS1* gene was detected in cultivar Sc748-5  
10 compared to cultivar BTx623, following inoculation with *C. sublineolum* (Fig. 3G-H).  
11 In the mesocotyl inoculation system, we have previously observed that fungal  
12 development in cultivar Sc748-5 (resistant host) was essentially contained during  
13 early stages of pathogenesis (Lo et al., 1999). In contrast, the fungal pathogen was  
14 able to colonize cultivar BTx623 (susceptible host) with the proliferation of primary  
15 and secondary hyphae. The differential expression of *SbSTS1* in the incompatible  
16 interaction suggests that the enzyme plays a key role in the expression of resistance  
17 against *C. sublineolum*.  
18  
19 An intriguing question remains regarding the identities of the sorghum defense

1 metabolites derived from SbSTS1 enzyme activities. In members of the Poaceae,  
2 resveratrol has been isolated from endophyte-infected grasses such as fescue, ryegrass,  
3 barley, sleepygrass, and bluegrass (Powell et al., 1994). Piceatannol, with an  
4 additional hydroxyl group at the 5'-position, was identified as a sugarcane  
5 phytoalexin after stalk inoculation with *Colletotrichum falcatum* (Brinker and Seigler,  
6 1993). However, neither resveratrol nor piceatannol were detected in  
7 acid-hydrolyzed extracts prepared from transgenic *Arabidopsis* or infected sorghum  
8 under our standard LC/MS-MS conditions (data not shown). It is likely that the  
9 immediate product(s) of SbCHS8 had been further modified *in planta*. The most  
10 common stilbene derivative piceid is a 3-*O*-glucoside of resveratrol, but the aglycone  
11 would be easily detected following acid hydrolysis. In Scots pine, pinosylvin is  
12 modified by an SAM dependent *O*-methyltransferase (PMT) to pinosylvin  
13 3-*O*-methyl ether following ozone or fungal elicitor treatment (Chiron et al., 2000).  
14 The recombinant PMT protein showed *in vitro* activities toward a broad range of  
15 substrates including resveratrol (Chiron et al., 2000). *O*-methyl ethers are common  
16 derivatives of flavonoid-related secondary metabolites. In fact, the two major  
17 3-deoxyanthocyanidin components luteolinidin and apigeninidin also exist as  
18 *O*-methyl ethers in sorghum (Lo et al., 1996; Lo and Nicholson, 1998). Whether a  
19 stilbene-*O*-methyl ether accumulates in inoculated sorghum plants is now under

1 investigation. Alternatively, SbSTS1 may utilize substrates other than  
2 phenylpropanoid-CoA esters leading to the formation of a more complex secondary  
3 metabolite in sorghum. Dayan et al. (2003) demonstrated recently that an STS-type  
4 reaction is involved in the biosynthesis of the sorghum root exudate sorgoleone.  
5 Sorgoleone and its derivatives are benzoquinone containing aliphatic tails of 15 or 17  
6 carbons with various degrees of unsaturation (Netzly et al., 1988). Thus, the “STS”  
7 involved would accept acyl-CoA esters of C16 and C18 fatty acids as starter  
8 molecules (Dayan et al., 2003). Examination on the activities of the recombinant  
9 SbSTS1 enzyme towards CoA esters of different fatty acids as well as other  
10 phenylpropanoids should help define its precise biochemical role in nature.  
11 Furthermore, recent advances in metabolic profiling technologies (von  
12 Roepenack-Lahaye et al., 2004) should allow one to identify novel natural products in  
13 plants in a more robust and efficient manner.

14

## 15 MATERIALS AND METHODS

### 16 Sorghum growth conditions and fungal inoculations

17 All sorghum seeds and fungal strains used in this study were described  
18 previously (Lo and Nicholson, 1998; Lo et al., 1999). For genomic DNA isolation,  
19 sorghum plants were grown in a greenhouse (16 h light, 8 h dark). For inoculation

1 experiments, sorghum seed were planted in rolls of germination paper and kept in the  
2 dark for 4 days at 28°C as described previously (Lo et al., 1996). Etiolated seedlings  
3 with elongated mesocotyls were then inoculated with conidial suspensions of *C.*  
4 *heterotrophus* or *C. sublineolum* at  $5.5 \times 10^4$  or  $1.0 \times 10^6$  conidia ml<sup>-1</sup>, respectively.  
5 Tween 20 was used as a wetting agent (100 µl 100 ml<sup>-1</sup>) in the inoculum. The  
6 resulting suspensions were misted onto the etiolated seedlings with an atomizer, and  
7 the plants were incubated at 100% RH at room temperature for at least 24 h.  
8

## 9 **DNA isolation and Southern blotting**

10 Genomic DNA samples were extracted from 4-week-old sorghum plants. Leaf  
11 tissues (1 g) were ground to a fine powder in liquid nitrogen and transferred to  
12 microfuge tubes containing the DNA extraction buffer (100 mM Tris-HCl, pH 8.0; 50  
13 mM EDTA, pH 8.0; 500 mM NaCl; 10 mM mercaptoethanol). 20% (w/v) Sodium  
14 dodecyl sulfate (SDS) (1 ml) was added to each tube and the mixtures were  
15 incubated at 65°C for 10 minutes. 5 M potassium acetate (5 ml) was then added and  
16 the tubes were incubated at 4°C for 20 min. The final mixtures were centrifuged at  
17 4,000 rpm for 20 min and the supernatants were transferred into tubes containing 10  
18 ml of isopropanol. After incubation at -20°C for 30 min, DNA samples were  
19 centrifuged at 14,000 rpm for 20 min. The pellets were washed in 70% ethanol,

1 air-dried, and resuspended in 0.5 ml of TE buffer (50 mM Tris-HCl, pH 8.0; 10 mM  
2 EDTA, pH 8.0). DNA samples (20 µg) were digested to completion with selected  
3 restriction enzymes. The digested DNA was separated by electrophoresis on a 0.8%  
4 agarose gel, depurinated, denatured, and blotted in 20× SSC (3 M NaCl, 0.3 M  
5 sodium citrate) by downward capillary transfer for at least 16 h onto a GeneScreen  
6 Plus nylon membrane (PerkinElmer, Boston, MA) then covalently cross-linked to the  
7 membrane with a UVP CL-1000 UV crosslinker (UVP, Cambridge, England).

8

9 **RNA extraction and northern blotting**

10 Sorghum tissues (1 g) were ground into a fine powder with liquid nitrogen and  
11 extracted with 1 ml of Trizol reagent (Invitrogen, Carlsbad, CA) in microfuge tubes.  
12 Chloroform (200 µl) was added to each tube and the resulting mixtures were  
13 centrifuged at 14,000 rpm for 10 min. The supernatants were transferred to new  
14 tubes containing 500 µl of isopropanol and 60 µl of 3 M sodium acetate. The  
15 mixtures were then centrifuged at 14,000 rpm for 10 minutes. The pellets were  
16 washed with 70% ethanol, air-dried, and re-suspended in 30 µl of RNase-free water.  
17 Fifteen micrograms of total RNA from each sample were denatured and fractionated  
18 on a 1% formaldehyde gel in 1× FA buffer, pH 7.0 (20 mM MOPS; 5 mM sodium  
19 acetate; 1 mM EDTA) and transferred to nylon membranes as described above.

1 Equal loading of RNA on gels was confirmed by ethidium bromide staining.

2

3 **Southern and northern hybridizations**

4 Individual membranes were pre-hybridized in hybridization buffer (1 M sodium

5 chloride; 0.1% dextran sulfate; 1% SDS; 100  $\mu$ g ml<sup>-1</sup> salmon sperm DNA) for 1 h at

6 65°C. The membranes were then hybridized in the same buffer containing different

7 denatured <sup>32</sup>P-labeled DNA probes for at least 16 h at the same temperature. The

8 hybridized membranes were washed twice in 2 $\times$  SSC, 0.1% SDS for 20 min at 65°C,

9 and twice in 0.2 $\times$  SSC, 0.1% SDS for 20 min at 65°C. High stringency washes in

10 0.1 $\times$  SSC at 65°C were performed when necessary. After washing, the membranes

11 were exposed to FUJI 100NIF X-ray films (Fuji Photo Ltd., Tokyo, Japan) with

12 intensifying screens at -80°C.

13

14 **Hybridization probes**

15 PCR fragments were generated for use as probes in the hybridization

16 experiments. The CHS8 probe (394 bp) was amplified from a full-length cDNA

17 clone (Lo et al., 2002) using primers derived from the 3'-UTR as well as part of the

18 coding region (Forward: 5'-GGC AAC ATG TCA AGC GTT TG-3'; Reverse: 5'-CCA

19 CTG CAC TGT GTT GAC TTG-3'). The CHS-U probe (643 bp) was amplified

1 from a full-length *SbCHS2* cDNA clone (L Pratt, University of Georgia, Athens, GA)  
2 using primers derived from a region conserved in *SbCHS1-SbCHS7* (Forward: 5'-  
3 CGC TGG ACG CCC GCC AGG ACA -3'; Reverse: 5' - GG GTG CGC CAC CCA  
4 GAA GAT). The hybridization probes were gel-purified (Qiagen, Valencia, CA)  
5 and labeled with <sup>32</sup>P-dCTP using the Rediprime II kit following the manufacturer's  
6 instructions (Amersham Biosciences, Piscataway, NJ).

7

8 **Complementation of *Arabidopsis tt4* mutants**

9 Full-length *SbCHS2* and *SbCHS8* cDNA fragments were each cloned into the  
10 *Bam*H I and *Xho* I sites of 103c-SK (E. Lam, Rutgers University, New Brunswick,  
11 NJ), an over-expression vector containing the CaMV 35S promoter and the nopaline  
12 synthase 3'-terminator. The resulting plasmids were cloned into the *Eco*R I and  
13 *Hind* III sites of the binary vector pCAMBIA 1300 (CAMBIA, Australia) to generate  
14 pCAM1300-*SbCHS2* and pCAM1300-*SbCHS8* for *Arabidopsis* transformation.

15

16 The *Arabidopsis tt4* mutants (CS8605) were obtained from the *Arabidopsis*  
17 Biological Resource Center (The Ohio State University, Columbus, OH). They are  
18 of the *Ler* genetic background and have a yellow seed coat color. The plant  
19 expression vectors were transformed into the mutants by the floral dip method

1 (Clough and Bent 1998). For selection of transformants, harvested seeds were  
2 surface-sterilized with 70% ethanol and 100% chlorox, followed by rinsing three  
3 times in sterilized water. The sterilized seeds were germinated on Murashige and  
4 Skoog (MS) (Sigma, MO, USA) agar plates containing 3% (v/v) sucrose, 25  $\mu\text{g ml}^{-1}$   
5 of hygromycin, and 500  $\mu\text{g ml}^{-1}$  of carbenicillin. After 2 weeks of selection,  
6 hygromycin-resistant plants ( $T_0$  plants) were transplanted to soil and placed in a  
7 growth chamber (25°C, 16 h light, 8 h dark).  $T_1$  seeds from individual  $T_0$  lines were  
8 collected and examined for seed coat color. For nitrogen deficiency assays,  $T_1$  seeds  
9 were germinated on MS plates without nitrogen sources. Accumulation of  
10 anthocyanins on cotyledons was observed in 4 to 5 days.

11

12 **HPLC analysis of transgenic *Arabidopsis tt4* mutants**

13  $T_1$  and  $T_2$  lines expressing *SbCHS2* or *SbCHS8* were grown on MS agar  
14 containing 3% (v/v) sucrose and hygromycin (25  $\mu\text{g ml}^{-1}$ ). Plant materials (0.5-1.0 g)  
15 were collected from 10 to 14 d-old seedlings and ground to a fine powder in liquid  
16 nitrogen. Methanol (300  $\mu\text{l}$ ) containing 1% (v/v) HCl was then added to the tissue  
17 powder. Acid hydrolysis was carried by addition of an equal volume of 2N HCl,  
18 followed by incubation at 70°C for 1 h. The hydrolyzed samples were evaporated to  
19 dryness under nitrogen and resuspended in 100  $\mu\text{l}$  of acidified methanol. Final

1 sample preparations (20  $\mu$ l) were injected onto a HP 1100 series HPLC system  
2 (Agilent Technologies, USA) equipped with a Nucleosil 100-5 C18 column (5  $\mu$ m,  
3 250 $\times$ 4 mm, Agilent Technologies). Chromatographic separation was performed  
4 using a solvent system of 1% acetic acid (v/v) (A) and acetonitrile (B) with a linear  
5 gradient of 20-60% B over 25 min. Flow rate was maintained at 0.6 ml  $\text{min}^{-1}$  and  
6 the elution was monitored by a diode-array detector (200-600 nm). Flavonol  
7 standards were obtained from Sigma (St Louis, MO).

8

9 **Over-expression of SbCHS proteins in *E. coli* and enzyme assays**

10 To express the sorghum proteins in *E. coli*, cDNAs were cloned into the *Nde* I  
11 and *Bam*H I sites of the pET14b vector (Novagen, San Diego, CA) containing a  
12 hexahistidine N-terminal fusion tag. To engineer the restriction sites in the inserts,  
13 PCR amplifications were performed using gene-specific primers [SbCHS2-F (5'-AGT  
14 CAT ATG GCC GGC GCG ACT GTG ACC-3') and SbCHS2-R (5'-AGT GGA TCC  
15 TCA GGC GGT GAT GGC CGC-3'); SbCHS8-F (5'-AGT CAT ATG ACG ACT  
16 GGG AAG GTA ACA-3') and SbCHS8-R (5'-GAT GGA TCC TCA TGC AGC CAC  
17 TGT GGT-3')] with the corresponding full-length cDNA clones as templates and the  
18 enzyme *Pfu* polymerase (Promega, Madison, WI). The resulting plasmids were each  
19 transformed into *E. coli* BL21-CodonPlus (DE3)-RIL cells (Stratagene). After 20 h

1 induction with 0.4 mM isopropyl -1-thio-  $\beta$  -D -galactopyranoside at 28°C, the  
2 recombinant proteins were purified from the bacterial cultures following procedures  
3 essentially as described previously (Samappito et al., 2002). The reference enzymes  
4 CalCHS1 and RtSTS1 were expressed and purified according to Samappito et al.  
5 (2002 and 2003).

6 The standard enzyme assays contained 100 mM HEPES buffer (pH 7.0), 20  $\mu$ M  
7 starter CoA, 15  $\mu$ M [ $2-^{14}\text{C}$ ] malonyl-CoA (24,000 dpm), and 1.0  $\mu$ g protein in a 50- $\mu$ l  
8 reaction. Starter CoAs (cinnamoyl-CoA, *p*-coumaroyl-CoA, caffeoyl-CoA,  
9 feruloyl-CoA), prepared essentially as described (Stöckigt and Zenk 1975), were  
10 kindly provided by Dagmar Knöfel (Department of Secondary Metabolism, IPB,  
11 Halle, Germany). The assay mixtures were incubated for 30 min at 30°C. The  
12 reactions were stopped by addition of 5  $\mu$ l of 50% (v/v) acetic acid and extracted with  
13 200  $\mu$ l of ethyl acetate. The organic phase was dried and separated by TLC (RP18)  
14 and developed in MeOH-H<sub>2</sub>O-acetic acid (75:25:1). The  $^{14}\text{C}$ -labelled products were  
15 visualized by phosphoimaging and quantification was performed with the  
16 ImageQuant software (Molecular Dynamics). Reaction products were identified by  
17 the use of authentic standards as well as comparison to published profiles of CalCHS1  
18 and RtSTS1 reactions (Samappito et al., 2002, 2003). To confirm the identities of  
19 flavanone and stilbene products, scaled-up reactions were performed containing 10  $\mu$ g

1 recombinant proteins, 50  $\mu$ M starter CoA, and 100  $\mu$ M malonyl-CoA in a total  
2 volume of 200  $\mu$ l for LC/ESI-MS/MS analysis in SRM mode. Positive and negative  
3 ESI mass spectra were obtained from a Finnigan MAT TSQ 7000 instrument  
4 (electrospray voltage, 4.5 kV; heated capillary temperature, 220 °C; sheath and  
5 auxiliary gas, nitrogen) coupled with a Surveyor MicroLC system equipped with an  
6 RP18-column (5  $\mu$ m, 1x100 mm, SepServ, Berlin). For all measurements a gradient  
7 system ranging from  $\text{H}_2\text{O}:\text{CH}_3\text{CN}$  90:10 (each containing 0.2% acetic acid) to 10:90  
8 over 15 min, followed by isocratic elution with a 10:90 mixture of both solvents for  
9 10 min, was used at a flow rate of 50  $\mu$ l min<sup>-1</sup>. Argon was used as collision gas and  
10 the collision pressure was at  $1.8 \times 10^{-3}$  Torr.

11

12 **ACKNOWLEDGEMENTS**

13 We thank Dr. Lee Pratt (University of Georgia) for providing the sorghum cDNA  
14 clones and Dr. Eric Lam (Rutgers University) for the use of his laboratory facilities at  
15 the University of Hong Kong as well as his suggestions on the manuscript. We are  
16 also indebted to Verona Dietl (IPB) for expression and purification of the recombinant  
17 proteins, Dagmar Knöfel (IPB) for providing the starter CoAs, and Christine Kuhnt  
18 (IPB) for recording the LC/ESI SRM measurements.

19

1    **LITERATURE CITED**

2    **Austin MB, Noel AP** (2003) The chalcone synthase superfamily of type III  
3    polyketide synthases. *Nat Prod Rep* **20**:79-110

4

5    **Brinker AM, Seigler DS** (1993) Time course of piceatannol accumulation in resistant  
6    and susceptible sugarcane stalks after inoculation with *Colletotrichum falcatum*.  
7    *Physiol Mol Plant Pathol* **42**: 169-176

8

9    **Busk PK, Møller BL** (2002) Dhurrin synthesis in sorghum is regulated at the  
10   transcriptional level and induced by nitrogen fertilization in older plants. *Plant  
11   Physiol* **129**: 1222-1231

12

13   **Chiron H, Drouet A, Claudot A-C, Eckerskorn C, Trost M, Heller W, Ernst D,**

14   **Sandermann H** (2000) Molecular cloning and functional expression of a  
15   stress-induced multifunctional *O*-methyltransferase with pinosylvin methyltransferase  
16   activity from Scots pine (*Pinus sylvestris* L.) *Plant Mol Biol* **44**: 733-745

17

18   **Clough SJ, Bent AF** (1998) Floral dip: a simplified method for *Agrobacterium*-  
19   mediated transformation of *Arabidopsis thaliana*. *Plant J* **16**:735-743

1

2   **Czarnota MA, Paul RN, Dayan FE, Numal CI, Weston LA** (2001) Mode of action,  
3   localization of production, chemical nature, and activity of sorgoleone: a potent PSII  
4   inhibitor in *Sorghum* spp. root exudates.   Weed Technol **15**:813-825

5

6   **Dayan FE, Kagan IA, Rimando AM** (2003) Elucidation of the biosynthetic pathway  
7   of the allelochemical sorgoleone using retrobiosynthetic NMR analysis.   J Biol  
8   Chem **278**: 28607-28611

9

10   **Dong X, Braun EL, Grotewold E** (2001) Functional conservation of plant secondary  
11   metabolic enzymes revealed by complementation of arabidopsis flavonoid mutants  
12   with maize genes.   Plant Physiol **127**: 46-57

13

14   **Hipskind J, Goldsbrough, PB, Urmeev F, Nicholson RL** (1996) Synthesis of  
15   3-deoxyanthocyanidin phytoalexins in sorghum does not occur via the same pathway  
16   as 3-hydroxylated anthocyanidins and phlobaphenes. Maydica **41**:155-166

17

18   **Hsieh M-H, Lam H-M, van de Loo FJ, Coruzzi G** (1998) A PII-like protein in  
19   *Arabidopsis*: putative role in nitrogen sensing.   Proc Natl Acad Sci USA **95**:

1 13965-13970

2

3 **Lo C, Coolbaugh RC, Nicholson RL** (2002) Molecular characterization and *in silico*

4 expression analysis of a chalcone synthase gene family in sorghum. *Physiol Mol*

5 *Plant Pathol* **61**: 179-188

6

7 **Lo S-C, Weiergang I, Bonham C, Hipskind J, Wood K, Nicholson RL** (1996)

8 Phytoalexin accumulation in sorghum: Identification of a methyl ether of luteolinidin.

9 *Physiol Mol Plant Pathol* **49**:21-31

10

11 **Lo S-CC, de Verdier K, Nicholson RL** (1999) Accumulation of

12 3-deoxyanthocyanidin phytoalexins and resistance to *Colletotrichum sublineolum* in

13 sorghum. *Physiol Mol Plant Pathol* **55**: 263-273

14

15 **Lo S-CC, Nicholson RL** (1998) Reduction of light-induced anthocyanin

16 accumulation in inoculated sorghum mesocotyls: implications for a compensatory

17 role in the defense response. *Plant Physiol* **116**: 979-989

18

19 **Netzly DH, Riopel JL, Ejeta G, Butler LG** (1988) Germination stimulants of

1      witchweed (*Striga asiatica*) from hydrophobic root exudates of sorghum (*Sorghum*  
2      *bicolor*).   Weed Sci 36: 441-446

3

4      **Powell RG, TePaske MR, Platner RD, White JF, Clement SL** (1994) Isolation of  
5      resveratrol from *Festuca versuta* and evidence for the widespread occurrence of this  
6      stilbene in the Poaceae.   Phytochemistry 35: 335-338

7

8      **Preisig-Müller R, Gnau P, Kindl H** (1995) The inducible 9,  
9      10-dihydrophenanthrene pathway: characterization and expression of bibenzyl  
10     synthase and S-adenosylhomocysteine hydrolase.   Arch Biochem Biophys 317:  
11     201-7

12

13     **Preisig-Müller R, Schwkendiek A, Brehm I, Reif H-J, Kindl H** (1999)  
14     Characterization of a pine multigene family containing elicitor-responsive stilbene  
15     synthase genes.   Plant Mol Biol 39: 221-229

16

17     **Samappito S, Page J, Schmidt J, De-Eknamkul W, Kutchan TM** (2002) Molecular  
18     characterization of root-specific chalcone synthases from *Cassia alata*.   Planta  
19     216:64-71

1

2     **Samappito S, Page J, Schmidt J, De-Eknamkul W, Kutchan TM** (2003) Aromatic  
3     and pyrone polyketides synthesized by a stilbene synthase from *Rheum tatarium*.  
4     Phytochemistry **62**:313-323

5

6     **Schröder G, Brown JWS, Schröder J** (1988) Molecular analysis of resveratrol  
7     synthase: cDNA, genomic clones and relationship with chalcone synthase.   Eur J  
8     Biochem **172**: 161-169

9

10    **Snyder BA, Nicholson RL** (1990) Synthesis of phytoalexins as a site specific  
11    response to fungal ingress.   Science **248**: 1637-1639

12

13    **Sparvoli F, Martin C, Scienza A, Gavazzi G, Tonelli C** (1994) Cloning and  
14    molecular analysis of structural genes involved in flavonoid and stilbene biosynthesis  
15    in grape (*Vitis vinifera* L.).   Plant Mol Biol **24**: 743-755

16

17    **Springob K, Nakajima J-I, Yamazaki M, Saito K** (2003) Recent advances in the  
18    biosynthesis and accumulation of anthocyanins.   Nat Prod Rep **20**: 288-303

19

1    **Stecher G, Huck CW, Popp M, Bonn GK** (2001) Determination of flavonoids and  
2    stilbenes in red wine and related biological products by HPLC and  
3    HPLC-ESI-MS-MS.    *Fresenius J Anal Chem* **371**: 73-80

4

5    **Stöckigt J, Zenk MH** (1975) Chemical syntheses and properties of  
6    hydroxycinnamoyl-coenzyme A derivatives.    *Z Naturforsch C* **30**: 352-358

7

8    **Versari A, Parpinello GP, Tornielli GB, Ferrarini R, Giulive C** (2001) Stilbene  
9    compounds and stilbene synthase expression during ripening, wilting, UV treatment  
10   in grape cv. Corina.    *J Agric Food Chem* **49**: 5531-5536

11

12   **von Roepenack-Lahaye E, Degenkolb T, Zerjeski M, Franz M, Roth U,**  
13   **Wessjohann L, Schmidt J, Scheel D, Clemens S** (2004) Profiling of *Arabidopsis*  
14   secondary metabolites by capillary liquid chromatography coupled to electrospray  
15   ionization quadrupole time-of-flight mass spectrometry.    *Plant Physiol* **134**: 548-559

16

17   **Wiese W, Vornam B, Krause E, Kindl H** (1994) Structural organization and  
18   differential expression of three stilbene synthase genes on a 13 kb grapevine DNA  
19   fragment.    *Plant Mol Biol* **26**: 667-677

1

2 **Yamaguchi T, Kurosaki F, Suh D-Y, Sankawa U, Nishioka M, Akiyama T,**

3 **Shibuya M, Ebizuka Y** (1999) Cross-reaction of chalcone synthase and stilbene

4 synthase overexpressed in *Escherichia coli*. FEBS Lett **460**: 457-461

5

6 **Yamazaki Y, Suh D-Y, Sitthithaworn W, Ishiguro K, Kobayashi Y, Shibuya M,**

7 **Ebizuka Y, Sankawa U** (2001) Diverse chalcone synthase superfamily enzymes from

8 the most primitive vascular plant, *Psilotum nudum*. Planta **214**: 75-84

1    **FIGURE LEGENDS**

2    **Figure 1.** Reaction steps catalyzed by CHS and STS. Cinnamoyl-CoA and

3    *p*-coumaroyl-CoA are the common start substrates for CHS and STS enzymes.

4    Chalcones are usually converted to flavanones spontaneously *in vitro*.

5

6    **Figure 2.** Genomic Southern analysis of *SbCHS* genes. Total DNA samples were

7    prepared from the indicated cultivars and digested to completion with *Eco*R I (E),

8    *Hind* III (H), or *Xba* I (X). A, Southern blot containing the digested DNA samples

9    were hybridized with a *SbCHS8*-specific probe, CHS8. A single hybridization signal

10    was detected in each digested sample following film exposure for 3 d. B, A

11    universal CHS fragment, CHS-U, for the *SbCHS1-SbCHS7* genes was used to probe a

12    blot containing *Hind* III digested DNA samples. A number of hybridization signals

13    of varying intensities were detected following overnight film exposure and RLFPs

14    were observed among the cultivars.

15

16    **Figure 3.** Northern analysis of *SbCHS* gene expression. A, Etiolated seedlings

17    (4-d old) of cultivar DK46 were exposed to light and RNA samples were prepared

18    from tissues collected at the indicated time point (h). B, To examine tissue specific

19    expression (B), RNA samples were collected from 6-d-old roots (YR), 1-month-old

1 roots (MR), 6-d-old leaves (YL), 1-month-old leaves (ML), and developing panicles  
2 (P). For infection experiments, 4-d-old etiolated seedlings were inoculated and RNA  
3 samples were prepared from tissues collected at the indicated time points (h). C-D,  
4 DK46 plants were inoculated with *C. heterotrophus* and kept under light or in the dark.  
5 E-F, DK 46 plants were inoculated with *C. sublineolum* and kept under light or dark.  
6 G-H, Inbred cultivars BTx623 (susceptible) and Sc748-5 (resistant) were inoculated  
7 with *C. sublineolum*. Hybridization probes (CHS8 and CHS-U) were the same as  
8 those used in the Southern experiments.

9

10 **Figure 4.** Analysis of transgenic *Arabidopsis tt4* mutants. A, Complementation of  
11 seed coat color and anthocyanin pigmentation. Seed coats of wild type plants (*Ler*)  
12 are brown due to tannin deposition. *Ler* seedlings accumulate anthocyanin in  
13 cotyledons and hypocotyls when germinated on MS medium without nitrogen sources  
14 (MS-N). *Arabidopsis tt4* mutants produced seeds with yellow color and failed to  
15 accumulate anthocyanin under nitrogen deficiency. Note the complementation of *tt4*  
16 phenotypes in T1 lines of SbCHS2-expressing plants (*tt4* + *SbCHS2*). On the other  
17 hand, *SbCHS8* did not restore the *tt4* mutations in T1 transgenic plants (*tt4* +  
18 *SbCHS8*). Same phenotypes were observed in the T2 lines of *SbCHS8*-expressing  
19 plants (data not shown). B, HPLC profiles of transgenic *tt4* plants.

1 Acid-hydrolyzed extracts were prepared from T1 lines and analyzed by HPLC with  
2 elution monitoring at  $A_{360}$ . Note the detection of peaks representing quercetin (Q,  
3 19.0 min) and kaempferol (K, 23.0 min) in *Ler* and *tt4 + SbCHS2* plants.


4


5 **Figure 5.** Enzyme assays of recombinant CHS proteins. A, RP-TLC analysis of  
6 products extracted from enzyme assays of recombinant proteins (CalCHS1, SbCHS2,  
7 SbCHS8, and RtSTS1). Assays were performed with 1.0  $\mu$ g of purified protein,  
8 radiolabeled malonyl-CoA and either cinnamoyl-CoA or *p*-coumaroyl-CoA.


9 Positions of flavanones (Pc, pinocembrin; N, naringenin), stilbenes (Ps, pinosylvin; R,  
10 resveratrol), and the BNY-type and CTAL-type pyrone byproducts (BNY-P and  
11 CTAL-P) are indicated. Inset: SDS-PAGE analysis of recombinant proteins

12 visualized with Coomassie Brilliant Blue R250. Lane 1, SbCHS2 crude cell lysate;  
13 Lane 2, purified SbCHS2; Lane 3, SbCHS8 crude cell lysate; Lane 4, purified  
14 SbCHS8. B, Ratios of flavanone to stilbene products in the assay reactions.



15  $^{14}\text{C}$  -labelled products were quantified after phosphoimaging and ratios were  
16 calculated based on average values from three independent assays. C, LC/ESI-SRM  
17 analysis of reaction products. Flavanone and stilbene products were confirmed by  
18 LC/MS-MS in SRM mode. RT, retention time. CE, collision energy. Structures  
19 of the starter-CoAs, flavanones, and stilbenes are shown in Fig. 1.








pp059337F03-HT.tif



pp059337F04\_RGB.tif



**B**

	Flavanone: Stilbene ratio			
	CalCHS1	SbCHS2	SbCHS8	RtSTS1
Cinnamoyl-CoA (Pc: Ps)	100: 1.8	100: 2.5	5.5: 100	26: 100
<i>p</i> -Coumaroyl-CoA (N: R)	100: 0	100: 0	2.0: 100	8.7: 100

**C**

Compound	RT (min)	CE (eV)	SRM
Pc	15.53	-25 -25	m/z 257 [M+H] <sup>+</sup> → m/z 131 m/z 257 [M+H] <sup>+</sup> → m/z 153
N	12.83	-25 -25	m/z 273 [M+H] <sup>+</sup> → m/z 147 m/z 273 [M+H] <sup>+</sup> → m/z 153
Ps	14.44	+25	m/z 211 [M-H] <sup>-</sup> → m/z 169
R	11.42	+25 +25	m/z 227 [M-H] <sup>-</sup> → m/z 185 m/z 227 [M-H] <sup>-</sup> → m/z 143

pp059337F05-HT.tif