



Contents lists available at ScienceDirect

Biochemical and Biophysical Research Communications

journal homepage: www.elsevier.com/locate/ybbrc



Male germ cell-specific protein Trs4 binds to multiple proteins

Yu-Qiang Shi <sup>a,b,1</sup>, Yin-Chuan Li <sup>a,1</sup>, Xiao-Qian Hu <sup>a</sup>, Tao Liu <sup>a</sup>, Shang-Ying Liao <sup>a</sup>, Jian Guo <sup>a</sup>, Lin Huang <sup>a</sup>, Zhao-Yuan Hu <sup>a</sup>, Alan Y.B. Tang <sup>c</sup>, Kai-Fai Lee <sup>c</sup>, William S.B. Yeung <sup>c</sup>, Chun-Sheng Han <sup>a,\*</sup>, Yi-Xun Liu <sup>a,\*</sup>

<sup>a</sup>State Key Laboratory of Reproductive Biology, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, PR China

<sup>b</sup>College of Bio-Engineering, Weifang University, Weifang, Shandong Province 261061, PR China

<sup>c</sup>Department of Obstetrics and Gynaecology, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Pokfulam, Hong Kong, PR China

ARTICLE INFO

Article history:  
Received 31 July 2009  
Available online xxx

Keywords:  
Trs4  
IQ calmodulin-binding motif  
Testis  
Cryptorchidism  
Spermatogenesis

ABSTRACT

Temperature-related sequence 4 (*Trs4*) has been identified as a testis-specific gene with expression sensitive to the abdominal temperature changes induced by artificial cryptorchidism. In murine testes, *Trs4* mRNA was detected in round spermatids and its protein was localized mainly in the elongating spermatids as well as in the acrosomes and tails of mature spermatozoa. Using a yeast two-hybrid screening system, we identified *Rshl-2*, *Gstmu1*, and *Ddc8* as putative binding partners of the *Trs4* protein in mouse testes. Their interactions were confirmed by in vivo and in vitro binding assays. Further studies demonstrated that *Ddc8*, a newly identified gene with unknown functions, displayed a similar expression pattern with *Trs4* in mouse testes. In particular, *Trs4*, *Ddc8*, and *Rshl-2* proteins were co-localized to the tails of mature spermatozoa. These results suggested that *Trs4* might be involved in diverse processes of spermiogenesis and/or fertilization through interactions with its multiple binding partners.

© 2009 Published by Elsevier Inc.

Introduction

Spermatogenesis is a complex process involving mitosis in spermatogonia, meiosis in spermatocytes and dramatic morphological changes to spermatids leading to the production of ciliated mature spermatozoa. This is a highly regulated process in which many testis-specific genes are involved [1]. It is estimated that about 4% of the genes in the mouse genome are testis-specific: most are transcribed in round spermatids but are not translated until the proteins are needed in later stages [2]. A number of highly specialized strategies for gene regulation are adopted during spermatogenesis, including a unique chromatin reorganization program, the use of distinct promoter elements and specific transcription factors [3]. Defective spermatogenesis is a major cause of human male infertility [4], so elucidating the functions of these testis-specific genes will not only help in understanding the mechanism of spermatogenesis, but might also allow the development of novel treatments for male infertility [2].

Spermatocytes and spermatids are very sensitive to many internal and external stresses. For example, they undergo apoptosis in response to hyperthermia [5,6]. Spermatogenesis occurs in the testes at temperature 4–5 °C lower than the body core temperature in most mammals. Patients with testes in the abdomen cavity (crypt-

orchidism) suffer from spermatogenic impairments, which can be restored by bringing the cryptorchid testes surgically back to the scrotum [7]. We and others have used animal models to study the effect of artificial cryptorchidism or heat shock on spermatogenesis [8,9] and two novel heat-sensitive genes from rat spermatids, *T6-441* and *Afaf* were identified [10,11]. Using a cryptorchid rat model, we have isolated an expressed sequence tag (EST) in rat testes, named *Trs4* (temperature-related sequence 4) [12]. Here we report the cloning and expression pattern of *Trs4* and its interaction with proteins including *Rshl-2*, *Ddc8*, and *Gstmu1* in mouse testes. We hypothesize that *Trs4* might be involved in diverse processes of spermiogenesis and/or fertilization through interactions with its multiple binding partners.

Methods

**Animal experiments.** Sprague–Dawley (SD) rats and CD-1 mice were obtained from the Experimental Animal Center, Chinese Academy of Sciences. All animals were treated in accordance with the NIH Guide for the Care and Use of Laboratory Animals and all protocols were approved by the Committee of Animal Care and Use of the Institute of Zoology, Chinese Academy of Sciences. The procedure for artificial cryptorchidism was performed as described before [12].

**Reagents.** If not stated otherwise, all reagents for cell culture were purchased from Invitrogen (Carlsbad, CA) and Sigma–Aldrich (St. Louis, MO). The Matchmaker library construction and screening

\* Corresponding authors. Fax: +86 10 64807105 (C.-S. Han), +86 10 64807038 (Y.-X. Liu).

E-mail addresses: hancs@ioz.ac.cn (C.-S. Han), liuyx@ioz.ac.cn (Y.-X. Liu).

<sup>1</sup> These authors contributed equally to this work.

kit, SMART RACE cDNA Amplification Kit and Matchmaker Co-IP kit were purchased from Clontech (BD Biosciences, San Jose, CA). Anti-green fluorescent protein (GFP) monoclonal antibody was purchased from Santa Cruz Biotechnology (Santa Cruz, CA). Anti-Flag monoclonal antibody was purchased from Sigma-Aldrich.

**Rapid amplification of 5'- and 3'-cDNA ends.** Total RNAs from adult SD rat and CD-1 mice testes were extracted to synthesize complementary DNA (cDNA). Rapid amplification of 5' and 3' complementary DNA ends (RACE) was performed using SMART RACE cDNA Amplification Kits. Domain and motif analysis followed the annotations of the NCBI Conserved Domain Database (<http://www.ncbi.nlm.nih.gov/Structure/cdd/cdd.shtml>). Multiple sequence alignment was performed using the EBI ClustalW service (<http://www.ebi.ac.uk/Tools/webservices/services/clustalw>).

**Recombinant protein expression and polyclonal antibody production.** The mouse Trs4 cDNA fragment (amino acids 31–101) and mouse Ddc8 cDNA fragment (amino acids 182–282) were subcloned in-frame into the pET21b(+) vector with the C-terminal tagged with His6 peptide. His-tagged fusion protein was then expressed in *Escherichia coli* strain BL21 and purified using a HisTrap FF column (Amersham Pharmacia, Piscataway, NJ). The antibodies were produced and purified as described [11].

**Northern blotting, in situ hybridization, and reverse transcription-polymerase chain reaction (RT-PCR).** Total RNAs from multiple adult rat tissues were extracted for Northern blotting and RT-PCR analysis. The primer pairs, 5'-TTG CTT CCA TCG GGA GAC ATA G-3' and 5'-CGT GTT TAT CCC ACC TGA CCA T-3' were used to synthesize probes for Northern blotting and in situ hybridization as described [10]. The same primers were also used to detect mRNA expression in rat testes by RT-PCR. Sertoli and Leydig cells from adult mice were prepared by two-step enzymatic digestion and cultured as described [13]. Total RNAs from mouse testes at different stages of development, the cultured primary testicular cells and the GC2-spd spermatocyte cell lines were extracted for RT-PCR analysis. The primer pairs 5'-aaatgg aggttg gag tgg ga-3' and 5'-ATA CGA GCA GCG GAT CTA TG-3', 5'-GAC CAC CTT AGG ACA AGA TTG-3' and 5'-TCC TGC CAC TCT GCC TTT AAC-3', were used to detect *Trs4* and *Ddc8* mRNA expression in mouse testes by RT-PCR, respectively.

**Confocal fluorescence and immunohistochemistry.** The whole coding sequence of mouse *Trs4* cDNA was cloned in-frame into pEGFP-N1 to construct a *Trs4*-GFP recombinant plasmid. The *Trs4*-GFP fusion protein was expressed in HeLa cells using Lipofectamine 2000 transfection reagent. Anti-*Trs4* antibody (1:500) and TRITC-conjugated anti-rabbit secondary antibody (1:200) were used for indirect immunostaining. To investigate the localization of *Trs4* and *Ddc8* proteins in mature spermatozoa, mouse cauda epididymidis was isolated and incubated in DMEM medium. The dissociated sperm were collected and smeared on glass slides. After fixing and blocking, the slides were incubated with anti-*Trs4* or anti-*Ddc8* antibodies (1:400) or preimmune rabbit serum (negative control). Subsequently, the slides were incubated with FITC-conjugated anti-rabbit secondary antibody (1:200). The nuclei were counterstained with 4',6-diamidino-2-phenylindol (DAPI) and the images were captured using a laser confocal microscope (Zeiss). Immunohistochemistry was performed as described [11].

**Yeast two-hybrid screening.** Yeast two-hybrid screening was performed using the Matchmaker library construction and screening kit with some modifications [14]. Two bait plasmids, T1 encoding the N-terminal 407 amino acids of mouse *Trs4* (22–428) and T2 encoding the C-terminal 500 amino acids of mouse *Trs4* (281–780) were constructed.

**Coimmunoprecipitation.** To confirm their interactions in vivo, cDNA fragments of mouse *Trs4* bait plasmid T2 and its interacting proteins were subcloned into pFlag-CMV4 and pEGFP-N1 vectors to construct Flag and GFP-tagged plasmids, respectively. The Flag-*Trs4* was overexpressed in HEK293T cells with each of its

GFP-tagged interacting proteins using Lipofectamine 2000. Total protein lysates of the cells were incubated with anti-GFP antibody or rabbit immunoglobulin (IgG) for 2 h at 4 °C, followed by incubation with protein A coated agarose beads overnight at 4 °C. The agarose beads and captured protein complexes were washed six times and suspended in SDS sample buffer for immunoblotting with anti-GFP (1:1000) or anti-Flag (1:2000). In vitro binding assays were performed using the TNT<sup>®</sup> Quick Coupled Transcription/Translation Systems (Promega, Madison, WI) and Matchmaker Co-IP kit as described before [14].

**Western blot analysis.** Equal amount of protein (50–100 µg total protein/lane) were loaded and separated by SDS-PAGE. After the proteins were transferred to polyvinylidene fluoride (PVDF) membranes, the membranes were blocked and incubated with primary antibodies, followed by incubation with HRP-conjugated secondary antibody. Proteins were visualized by Enhanced Chemiluminescence kit (Pierce, Rockford, IL).

**Statistics.** Band intensities were analyzed and normalized with internal controls. Values are represented as means ± SEM of three separate experiments. Statistical analysis was performed using SPSS (version 13.0; SPSS Inc., Chicago, IL) and one-way ANOVA was used to analyze the data in different groups; *P* < 0.05 was assumed significant.

## Results

### Cloning and sequence analysis of *Trs4*


The sequence of rat *Trs4* (GenBank Accession No.: DQ132434) was obtained by RACE and sequence analysis showed that rat *Trs4* has a full length of 2370 nucleotide acids encoding a putative protein of 790 amino acids. Multiple sequence alignment indicated that the proteins are highly conserved across species. Importantly, a putative IQ calmodulin-binding motif and an ubiquitin-like structure (DUF2021 in the PFAM database; <http://pfam.sanger.ac.uk/>) were found in all the species tested (Supplementary Fig. 1).

### Expression profile of *Trs4* mRNA in murine testes


*Trs4* was specifically expressed in the rat testis and no expression was found in other tissues including the ovary, kidney, heart, liver, spleen, lung, brain and stomach by RT-PCR (Fig. 1A). This expression pattern was further confirmed by Northern blotting in rat tissues (Fig. 1B). A single transcript of about 3000 nucleotides was visualized on the blot, consistent with the result of RACE analysis. The developmental profiles of *Trs4* transcripts in testes were examined by Northern blotting in rats (Fig. 1C) and RT-PCR in mice (Fig. 1D), respectively. *Trs4* mRNA was first detected at 28 days post partum (dpp) in rats and 21 dpp in mice, at the time when round spermatids are produced in the first wave of spermatogenesis [15]. No expression was found in Sertoli cells, Leydig cells or GC2-spd cells cultured in vitro (Fig. 1D). In situ hybridization results confirmed that *Trs4* mRNA was localized in round spermatids of stages VII–VIII seminiferous tubules in mouse testes (Fig. 1E). Furthermore, the *Trs4* transcript in rat testis was found to be sensitive to intra-abdominal temperature and was decreased significantly after artificial cryptorchidism (Supplementary Fig. 2), which is consistent with our previous report [12].

### *Trs4* protein is localized in early elongating spermatids and mature spermatozoa

HeLa cells were transfected with *Trs4*-GFP fusion plasmid for detecting specificity of *Trs4* antibody and its subcellular localiza-



**Fig. 1.** Expression of *Trs4* mRNA in rat and mouse testes. (A) *Trs4* mRNA was exclusively expressed in rat testes using RT-PCR analysis of multiple tissues. (B) Northern blotting on multiple tissues detected one transcript of about 3000 nucleotides in rat testes. (C) In postnatal rat testes, *Trs4* transcript was first detected at 28 days post partum (dpp) and thereafter by Northern blotting. (D) In mice, *Trs4* expression was first detected at 21 dpp. No expression of *Trs4* transcript was detected in the Sertoli cells, Leydig cells or GC2-spd cells by RT-PCR. (E) *Trs4* mRNA was visualized in round spermatids of mouse testes at stages VII–VIII seminiferous tubules using in situ hybridization analysis. G3PDH RNA and 28S rRNA were used as internal controls for RNA loadings. Scale bars = 50  $\mu$ m.



**Fig. 2.** Localization of *Trs4* protein in mouse testes. (A) The specificity of *Trs4* polyclonal antibody was tested in HeLa cells transfected with a *Trs4*-GFP fusion plasmid. The staining signals of *Trs4*-EGFP (green) and the anti-*Trs4* antibody (red) are in complete alignment in the transfected cells. No staining was developed in non-transfected cells. (B) A single band of 94 KDa was detected in Western blotting, consistent with the predicted size of *Trs4*. All lanes were loaded with 50  $\mu$ g protein extracts from mouse testes (lane 1), rat testes (lane 2), mouse epididymis (lane 3) and rat epididymis (lane 4). (C) Immunohistochemical data indicated that *Trs4* protein was mainly localized in the elongating spermatids of stages VIII–IX seminiferous tubules. (D) *Trs4* protein was also detected in the acrosomes and tails of mature spermatozoa prepared from the cauda epididymidis. Sperm-smear slides were incubated with anti-*Trs4* antibody and FITC-conjugated anti-rabbit IgG (green). The nuclei of HeLa cells and mature spermatozoa were stained with DAPI (blue) and the images were captured using a laser confocal microscope (Zeiss). Scale = 20  $\mu$ m. (For interpretation of color mentioned in this figure legend the reader is referred to the web version of the article.)



tion. As shown in Fig. 2A, the GFP-fusion protein was exclusively localized in the cytoplasm of transfected HeLa cells and co-localized with Trs4 immunostaining. Furthermore, no immunostaining signal of Trs4 was detected in untransfected HeLa cells. Specificity of the Trs4 antibody was also confirmed as a single band of protein with predicted molecular weight of 94 KDa detected in murine total testicular lysates using Western blotting (Fig. 2B). Immunohistochemical data indicated that the Trs4 protein was localized in the elongating spermatids of stages VIII–IX seminiferous tubules (Fig. 2C) and in the acrosomes and tails of mature spermatozoa prepared from the cauda epididymidis (Fig. 2D). A similar localization of Trs4 in rat testes was also observed using immunohistochemistry (data not shown).

### Trs4 protein interacts with Rshl-2, Gstm1, and Ddc8 proteins

Using a yeast two-hybrid system, putative interacting proteins including Rshl-2, Ddc8, Gstm1, and Odf-1 were identified using the T2 bait plasmid of the Trs4 protein. No interacting protein was identified when T1 was used as a bait. To eliminate potential false positives, *in vivo* and *in vitro* binding assays were performed subsequently. *In vivo* binding assays revealed that the Trs4 protein could coimmunoprecipitate with Rshl-2, Ddc8 and Gstm1 but not with Odf-1 from the lysates of HEK293T cells transfected with Flag-Trs4 and GFP-tagged interacting proteins (Fig. 3A). The results were further confirmed by the *in vitro* binding assays, which demonstrated that <sup>35</sup>S-Met-labeled Myc-Trs4 protein could coimmunoprecipitate with Rshl-2, Ddc8, and Gstm1 proteins except for Odf-1 (Fig. 3B).

### Spatiotemporal expression of Ddc8


The Ddc8 transcript was uniquely expressed in testes. It was first detected in the mouse testis at 21 dpp and no expression was found in Sertoli cells, Leydig cells or GC2-spd cells (Fig. 4A and B). Ddc8 mRNA was present in round spermatids of the stages VII–VIII seminiferous tubules (Fig. 4C) as revealed by *in situ* hybridization. The protein was also localized in the tails of elongated spermatids and spermatozoa of stages VII–VIII seminiferous tubules (Fig. 4D) and mature spermatozoa prepared from the cauda epididymidis (Fig. 4E) as detected by immunostaining.

## Discussion


We investigated the expression pattern of the testis-specific gene *Trs4* and its protein interactions with other binding partners. The expression of *Trs4* was examined using RT-PCR, Northern blotting, *in situ* hybridization and immunostaining techniques. The results of these assays together indicated that *Trs4* is a male germ cell-specific gene whose mRNA is expressed in round spermatids and whose protein is expressed in elongating spermatids and mature spermatozoa. In particular, the protein was mainly located in the acrosomes and tails of mature spermatozoa indicative of its diverse functions during spermiogenesis and/or fertilization. Because the *Trs4* protein sequences are evolutionarily conserved and contain an ubiquitin-like domain and an IQ calmodulin-binding motif, we hypothesize that they might play a role in spermatogenesis through interactions with other proteins.

Using a yeast two-hybrid system, we demonstrated that the Trs4 protein interacted with Rshl-2, Gstm1, and Ddc8 in the mouse testis. These interactions were further confirmed by *in vitro* and *in vivo* binding assays. Rshl-1 and Rshl-2 (radial spoke-head-like proteins) are radial spoke proteins that regulate the activity of axonemal inner arm dynein for maintaining the structural integrity and movement of spermatozoa through protein phosphorylation and dephosphorylation [16]. The radial spokes is a T-shaped structure extending from the A-tubule of each outer doublet microtubule to the center of the axoneme of motile structures such as cilia, flagella and the flagella of spermatozoa [17]. In line with this, Trs4 was localized in the acrosomes and tails of isolated mature epididymal spermatozoa. Therefore, Trs4 protein might be involved in regulating the movement of sperm by interaction with the Rshl-2 protein.

Gstm1 belongs to the glutathione S-transferase family of enzymes that are responsible for metabolizing a broad range of xenobiotics and carcinogens. Gstm1 can convert organic compounds to thioethers, a reaction that is the first step in a detoxification process leading to mercapturic acid formation [18,19]. Furthermore, Gstm1 binds to some chemical substances such as steroids, thyroxin, bile acids and bilirubin in a noncatalytic manner to facilitate their transport [18,20]. Gstm1 modulates stress-mediated signals by repressing apoptosis signal-regulating kinase 1 (ASK1). This activity occurs independently of its catalytic activity in intracellular glutathione metabolism [21]. The expression level of Trs4 decreased in the cryptorchid condition or by treatment of the testes



**Fig. 3.** Interactions of Trs4 protein with Rshl-2, Gstm1, and Ddc8 proteins. (A) *In vivo* binding assays revealed that Trs4 interacts with Rshl-2, Ddc8, and Gstm1. Total lysates of HEK293T cells transfected with Flag-Trs4 protein and each of its putative interacting proteins tagged with GFP were immunoprecipitated with an anti-GFP antibody and detected with anti-GFP or anti-Flag antibodies. HEK293T cells cotransfected with Flag-Trs4 and GFP plasmids were used as controls. (B) *In vitro* binding assays further confirmed that the Trs4 protein could coimmunoprecipitate with Rshl-2, Ddc8, and Gstm1 proteins but not Odf-1 when <sup>35</sup>S-Met-labeled Myc-Trs4 protein was incubated with the interacting proteins.



**Fig. 4.** Expression of *Ddc8* in mouse testes. (A) *Ddc8* mRNA was exclusively detected in mouse testes by RT-PCR using primers for the testis-specific transcript. (B) In developing mouse testes, the expression of *Ddc8* was first detected in the mouse testes at 21 dpp. (C) *Ddc8* mRNA was visualized in round spermatids of stages VII–VIII seminiferous tubules using in situ hybridization assays. (D) Immunohistochemical results indicated that the *Ddc8* protein is localized in the tails of elongated spermatids and spermatozoa of stages VII–VIII seminiferous tubules. (E) *Ddc8* protein was also detected in the tail of mature spermatozoa prepared from epididymis. Scale bars = 50  $\mu$ m.

at 43 °C in a water bath [12]. These treatments increase the level of free radicals in mitochondria that might cause germ cell apoptosis [22]. We hypothesize that *Trs4* could participate in regulating the level of oxidative molecules through interaction with *Gstmu1*. However, this interaction might only facilitate *Trs4* protein transport in the cell.

*Ddc8*, a newly discovered gene with unknown function, was first identified as a testis-specific gene using differential-display reverse transcription (DDRT)-PCR analysis searching for genes expressed differentially between prepubertal and adult mouse testes [23]. Further studies revealed that *Ddc8*, a gene nested in its host gene family the tissue inhibitors of metalloproteinase 2 (TIMP-2), was not testis-specific. Moreover, *Ddc8* expression in non-neural and neural tissues mimicked that of TIMP-2 and was upregulated in response to traumatic brain injury [24]. Interestingly, a specific transcript of *Ddc8* consisting of exon-1, exon-2, and exon-3 was found to exist only in mouse testes, while the expression of exon-3 possessed a character of universal expression by RT-PCR and in situ hybridization assays [24]. Using primers specifically designed to detect the testis-specific transcript of *Ddc8*, it was found specifically expressed and regulated in a development-dependent manner in mouse testes. In addition, the mRNA of *Ddc8* was only found in round spermatids and its protein was localized in the tails of elongated spermatids and spermatozoa in stages VII–VII of seminiferous tubules. *Ddc8* and *Trs4* thus share a similar expression pattern in testes and in mature spermatozoa. Moreover, the expression of exon-3 of *Ddc8* in other tissues including brain, kidney and lung, was also confirmed by RT-PCR (data not shown). Our findings are consistent with data reported previously [24]. We sug-

gest that a specific transcript of *Ddc8* exists in testes and that its expression is tightly controlled during spermatogenesis.

In conclusion, *Trs4* was specifically expressed in germ cells from early elongating spermatids to mature spermatozoa and it was downregulated by hyperthermia. The *Trs4* protein interacted with *Rshl-2*, *Gstmu1*, and *Ddc8* proteins in mouse testes and co-localized to the tail of mature spermatozoa with *Rshl-2* and *Ddc8*. We suggest that *Trs4* alone or in combination with its interacting proteins could play important roles in the processes of spermiogenesis and fertilization.

#### Acknowledgments

This study was supported by the National Basic Research Program of China (2006CB944000, 2006CB944004), the National “973” Program (2006CB504001), the NSFC-RGC joint Project (30618005 and N\_HKU712/06), Beijing NSF (5073032), the CAS Knowledge Innovation Project (KSCX2-YW-R-5) and the National Natural Science Foundation of China (90508008).

#### Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.bbrc.2009.08.053.

#### References

- [1] K.C. Kleene, A possible meiotic function of the peculiar patterns of gene expression in mammalian spermatogenic cells, *Mech. Dev.* 106 (2001) 3–23.

- [2] N. Schultz, F.K. Hamra, D.L. Garbers, A multitude of genes expressed solely in meiotic or postmeiotic spermatogenic cells offers a myriad of contraceptive targets, *Proc. Natl. Acad. Sci. USA* 100 (2003) 12201–12206.
- [3] P. Sassone-Corsi, Unique chromatin remodeling and transcriptional regulation in spermatogenesis, *Science* 296 (2002) 2176–2178.
- [4] R.I. McLachlan, C. Mallidis, K. Ma, S. Bhasin, D.M. de Kretser, Genetic disorders and spermatogenesis, *Reprod. Fertil. Dev.* 10 (1998) 97–104.
- [5] T. Koji, Y. Hishikawa, Germ cell apoptosis and its molecular trigger in mouse testes, *Arch. Histol. Cytol.* 66 (2003) 1–16.
- [6] R.J. Levine, R.M. Mathew, C.B. Chenault, M.H. Brown, M.E. Hurtt, K.S. Bentley, K.L. Mohr, P.K. Working, Differences in the quality of semen in outdoor workers during summer and winter, *N. Engl. J. Med.* 323 (1990) 12–16.
- [7] E. Caroppo, C. Niederberger, S. Elhanbly, R. Schoor, L. Ross, G. D'Amato, Effect of cryptorchidism and retractile testes on male factor infertility: a multicenter, retrospective, chart review, *Fertil. Steril.* 83 (2005) 1581–1584.
- [8] J. Guo, S.X. Tao, M. Chen, Y.Q. Shi, Z.Q. Zhang, Y.C. Li, X.S. Zhang, Z.Y. Hu, Y.X. Liu, Heat treatment induces liver receptor homolog-1 expression in monkey and rat sertoli cells, *Endocrinology* 148 (2007) 1255–1265.
- [9] M. Chen, H. Cai, J.L. Yang, C.L. Lu, T. Liu, W. Yang, J. Guo, X.Q. Hu, C.H. Fan, Z.Y. Hu, F. Gao, Y.X. Liu, Effect of heat stress on expression of junction-associated molecules and upstream factors androgen receptor and Wilms' tumor 1 in monkey sertoli cells, *Endocrinology* 149 (2008) 4871–4882.
- [10] X.X. Song, Y.C. Li, Y.Q. Shi, X.Q. Hu, Z.Y. Hu, C.S. Han, Y.X. Liu, Cloning and characterization of a novel spermiogenesis-related gene, T6441, in rat testis, *Front. Biosci.* 11 (2006) 143–150.
- [11] Y.C. Li, X.Q. Hu, K.Y. Zhang, J. Guo, Z.Y. Hu, S.X. Tao, L.J. Xiao, Q.Z. Wang, C.S. Han, Y.X. Liu, Afaf, a novel vesicle membrane protein, is related to acrosome formation in murine testis, *FEBS Lett.* 580 (2006) 4266–4273.
- [12] C.X. Guo, T.S. Tang, X.M. Mu, S.H. Li, G.Q. Fu, H. Liu, Y.X. Liu, Cloning of novel temperature-related expressed sequence tags in rat testis during spermatogenesis, *Biochem. Biophys. Res. Commun.* 258 (1999) 401–406.
- [13] A.F. Karl, M.D. Griswold, Sertoli cells of the testis: preparation of cell cultures and effects of retinoids, *Methods Enzymol.* 190 (1990) 71–75.
- [14] H. Nian, C.H. Fan, S.Y. Liao, Y.Q. Shi, K.Y. Zhang, Y.X. Liu, C.S. Han, RNF151, a testis-specific RING finger protein, interacts with dysbindin, *Arch. Biochem. Biophys.* 465 (2007) 157–163.
- [15] M. Malkov, Y. Fisher, J. Don, Developmental schedule of the postnatal rat testis determined by flow cytometry, *Biol. Reprod.* 59 (1998) 84–92.
- [16] P. Padma, Y. Satouh, K. Wakabayashi, A. Hozumi, Y. Ushimaru, R. Kamiya, K. Inaba, Identification of a novel leucine-rich repeat protein as a component of flagellar radial spoke in the Ascidian *Ciona intestinalis*, *Mol. Biol. Cell* 14 (2003) 774–785.
- [17] P. Yang, D.R. Diener, C. Yang, T. Kohno, G.J. Pazour, J.M. Dienes, N.S. Agrin, S.M. King, W.S. Sale, R. Kamiya, J.L. Rosenbaum, G.B. Witman, Radial spoke proteins of *Chlamydomonas flagella*, *J. Cell. Sci.* 119 (2006) 1165–1174.
- [18] J. Guo, L. Zimniak, P. Zimniak, J.L. Orchard, S.V. Singh, Cloning and expression of a novel Mu class murine glutathione transferase isoenzyme, *Biochem. J.* 366 (2002) 817–824.
- [19] H. Raza, M.A. Robin, J.K. Fang, N.G. Avadhani, Multiple isoforms of mitochondrial glutathione S-transferases and their differential induction under oxidative stress, *Biochem. J.* 366 (2002) 45–55.
- [20] J.D. Hayes, D.J. Pulford, The glutathione S-transferase supergene family: regulation of GST and the contribution of the isoenzymes to cancer chemoprotection and drug resistance, *Crit. Rev. Biochem. Mol. Biol.* 30 (1995) 445–600.
- [21] S.G. Cho, Y.H. Lee, H.S. Park, K. Ryoo, K.W. Kang, J. Park, S.J. Eom, M.J. Kim, T.S. Chang, S.Y. Choi, J. Shim, Y. Kim, M.S. Dong, M.J. Lee, S.G. Kim, H. Ichijo, E.J. Choi, Glutathione S-transferase mu modulates the stress-activated signals by suppressing apoptosis signal-regulating kinase 1, *J. Biol. Chem.* 276 (2001) 12749–12755.
- [22] C.Q. Li, G.N. Wogan, Nitric oxide as a modulator of apoptosis, *Cancer Lett.* 226 (2005) 1–15.
- [23] R.D. Catalano, M. Vlad, R.C. Kennedy, Differential display to identify and isolate novel genes expressed during spermatogenesis, *Mol. Hum. Reprod.* 3 (1997) 215–221.
- [24] D.M. Jaworski, M. Beem-Miller, G. Lluri, R. Barrantes-Reynolds, Potential regulatory relationship between the nested gene DDC8 and its host gene tissue inhibitor of metalloproteinase-2, *Physiol. Genomics* 28 (2007) 168–178.

372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406