

INVERSES OF NEW HILBERT-PACHPATTE-TYPE INEQUALITIES

CHANG-JIAN ZHAO AND WING-SUM CHEUNG

Received 7 February 2006; Revised 3 June 2006; Accepted 5 June 2006

We establish a new inequality of Hilbert type for a finite double number of nonnegative sequences of real numbers and some interrelated results, which are inverse and general forms of Pachpatte's and Handley's results. An integral version and some interrelated results are also obtained. These results provide some new estimates on such types of inequalities.

Copyright © 2006 C.-J. Zhao and W.-S. Cheung. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

The various generalizations and sharpenings of Hilbert's double series inequality and its integral version are obtained by Pachpatte, Handley et al., Gao et al. (see [1, 3, 5, 7, 10, 12, 16–18]). An elegant survey on this kind of inequalities was provided by Yang and Rassias (see [13]). Moreover, Pachpatte [9] established a new Hilbert-type inequality and its integral version as follows.

THEOREM 1.1. *Let $\{a_m\}$, $\{b_n\}$ be two nonnegative sequences of real numbers defined for $m = 1, 2, \dots, k$ and $n = 1, 2, \dots, r$ with $a_0 = b_0 = 0$, and let $\{p_m\}$, $\{q_n\}$ be two positive sequences of real numbers defined for $m = 1, 2, \dots, k$, $n = 1, 2, \dots, r$, where k, r are natural numbers. Define $P_m = \sum_{s=1}^m p_s$ and $Q_n = \sum_{t=1}^n q_t$. Let ϕ and ψ be two real-valued nonnegative, convex, and submultiplicative functions defined on $\mathbb{R}_+ = [0, \infty)$. Then*

$$\sum_{m=1}^k \sum_{n=1}^r \frac{\phi(a_m)\psi(b_n)}{m+n} \leq M(k,r) \left(\sum_{m=1}^k (k-m+1) \left(p_m \phi \left(\frac{\nabla a_m}{p_m} \right) \right)^2 \right)^{1/2} \times \left(\sum_{n=1}^r (r-n+1) \left(q_n \psi \left(\frac{\nabla b_n}{q_n} \right) \right)^2 \right)^{1/2}, \quad (1.1)$$

2 Inverses of new Hilbert-Pachpatte-type inequalities

where

$$M(k, r) = \frac{1}{2} \left(\sum_{m=1}^k \left(\frac{\phi(P_m)}{P_m} \right)^2 \right)^{1/2} \left(\sum_{n=1}^r \left(\frac{\phi(Q_n)}{Q_n} \right)^2 \right)^{1/2}, \quad (1.2)$$

and $\nabla a_m = a_m - a_{m-1}$, $\nabla b_n = b_n - b_{n-1}$.

THEOREM 1.2. Let $f \in C^1[[0, x], \mathbb{R}^+]$, $g \in C^1[[0, y], \mathbb{R}^+]$ with $f(0) = g(0) = 0$ and let $p(\sigma)$, $q(\tau)$ be two positive functions defined for $\sigma \in [0, x]$ and $\tau \in [0, y]$. Let $P(s) = \int_0^s p(\sigma) d\sigma$ and $Q(t) = \int_0^t q(\tau) d\tau$ for $s \in [0, x]$ and $t \in [0, y]$, where x, y are positive real numbers. Let ϕ , and ψ be as in Theorem 1.1. Then

$$\int_0^x \int_0^y \frac{\phi(f(s))\psi(g(t))}{s+t} ds dt \leq L(x, y) \left(\int_0^x (x-s) \left(p(s) \phi \left(\frac{f'(s)}{p(s)} \right) \right)^2 ds \right)^{1/2} \times \left(\int_0^y (y-t) \left(q(t) \phi \left(\frac{g'(t)}{q(t)} \right) \right)^2 dt \right)^{1/2}, \quad (1.3)$$

where

$$L(x, y) = \frac{1}{2} \left(\int_0^x \left(\frac{\phi(P(s))}{P(s)} \right)^2 ds \right)^{1/2} \left(\int_0^y \left(\frac{\phi(Q(t))}{Q(t)} \right)^2 dt \right)^{1/2}, \quad (1.4)$$

and $'$ denotes the derivative of a function.

In [4], Handley et al. gave general versions of inequalities (1.1) and (1.3) as follows.

THEOREM 1.3. Let $\{a_{i,m_i}\}$ ($i = 1, 2, \dots, n$) be n sequences of nonnegative real numbers defined for $m_i = 1, 2, \dots, k_i$ with $a_{1,0} = a_{2,0} = \dots = a_{n,0} = 0$, and let $\{p_{i,m_i}\}$ be n sequences of positive real numbers defined for $m_i = 1, 2, \dots, k_i$, where k_i ($i = 1, 2, \dots, n$) are natural numbers. Set $P_{i,m_i} = \sum_{s=1}^{m_i} p_{i,s}$. Let ϕ_i ($i = 1, 2, \dots, n$) be n real-valued nonnegative convex and sub-multiplicative functions defined on \mathbb{R}_+ . Let $\alpha_i \in (0, 1)$, and set $\alpha'_i = 1 - \alpha_i$ ($i = 1, 2, \dots, n$), $\alpha = \sum_{i=1}^n \alpha_i$, and $\alpha' = \sum_{i=1}^n \alpha'_i = n - \alpha$. Then

$$\sum_{m_1=1}^{k_1} \dots \sum_{m_n=1}^{k_n} \frac{\prod_{i=1}^n \phi_i(a_{i,m_i})}{\left(\sum_{i=1}^n \alpha'_i m_i \right)^{\alpha'}} \leq M(k_1, \dots, k_n) \prod_{i=1}^n \left(\sum_{m_i=1}^{k_i} (k_i - m_i + 1) \left(p_{i,m_i} \phi_i \left(\frac{\nabla a_{i,m_i}}{P_{i,m_i}} \right) \right)^{1/\alpha'_i} \right)^{\alpha_i}, \quad (1.5)$$

where

$$M(k_1, \dots, k_n) = \frac{1}{(\alpha')^{\alpha'}} \prod_{i=1}^n \left(\sum_{m_i=1}^{k_i} \left(\frac{\phi_i(P_{i,m_i})}{P_{i,m_i}} \right)^{1/\alpha'_i} \right)^{\alpha'_i}, \quad (1.6)$$

and $\nabla a_{i,m_i} = a_{i,m_i} - a_{i,m_i-1}$ ($i = 1, 2, \dots, n$).

THEOREM 1.4. Let $f_i \in C^1[[0, k_i], \mathbb{R}_+]$, $i = 1, 2, \dots, n$, with $f_i(0) = 0$ ($i = 1, 2, \dots, n$), let $p_i(\sigma_i)$ be n positive functions defined for $\sigma_i \in [0, x_i]$ ($i = 1, \dots, n$). Set $P_i(s_i) = \int_0^{s_i} p_i(\sigma_i) d\sigma_i$ for $s_i \in [0, x_i]$, where x_i are positive real numbers. Let ϕ_i , α_i , α'_i , α , and α' be as in Theorem 1.3. Then

$$\int_0^{x_1} \cdots \int_0^{x_n} \frac{\prod_{i=1}^n \phi_i(f_i(s_i))}{\left(\sum_{i=1}^n \alpha'_i s_i\right)^{\alpha'}} ds_1 \cdots ds_n \leq L(x_1, \dots, x_n) \prod_{i=1}^n \left(\int_0^{x_i} (x_i - s_i) \left(p_i(s_i) \phi_i \left(\frac{f'_i(s_i)}{P_i(s_i)} \right) \right)^{1/\alpha_i} ds_i \right)^{\alpha_i}, \tag{1.7}$$

where

$$L(x_1, \dots, x_n) = \frac{1}{(\alpha')^{\alpha'}} \prod_{i=1}^n \left(\int_0^{x_i} \left(\frac{\phi_i(P_i(s_i))}{P_i(s_i)} \right)^{1/\alpha'_i} ds_i \right)^{\alpha'_i}. \tag{1.8}$$

The main purpose of the present paper is to establish their reverse versions, which are more extensive results for this type of inequalities. Our main results are given in the following theorems.

THEOREM 1.5. Let $\{a_{s_i, t_i, m_{s_i}, m_{t_i}}\}$ ($i = 1, 2, \dots, n$) be n sequences of nonnegative numbers defined for $m_{s_i} = 1, 2, \dots, k_{s_i}$, $m_{t_i} = 1, 2, \dots, k_{t_i}$, with $a_{s_i, t_i, 0, m_{t_i}} = a_{s_i, t_i, m_{s_i}, 0} = 0$ ($i = 1, 2, \dots, n$), where k_{s_i} and k_{t_i} ($i = 1, 2, \dots, n$) are natural numbers. Let $\{p_{s_i, t_i, m_{s_i}, m_{t_i}}\}$ be n sequences of positive real numbers defined for m_{s_i}, m_{t_i} . Set

$$P_{s_i, t_i, m_{s_i}, m_{t_i}} = \sum_{m_{\xi_i}=1}^{m_{s_i}} \sum_{m_{\eta_i}=1}^{m_{t_i}} p_{s_i, t_i, m_{\xi_i}, m_{\eta_i}} \quad (i = 1, 2, \dots, n). \tag{1.9}$$

Let ϕ_i ($i = 1, 2, \dots, n$) be n real-valued nonnegative concave and supermultiplicative functions defined on \mathbb{R}_+ . Let $\alpha_i \in (1, \infty)$. Set $\alpha'_i = 1 - \alpha_i$ ($i = 1, 2, \dots, n$), $\alpha = \sum_{i=1}^n \alpha_i$, and $\alpha' = \sum_{i=1}^n \alpha'_i = n - \alpha$. Define operators

$$\begin{aligned} \nabla_1 a_{s_i, t_i, m_{s_i}, m_{t_i}} &= a_{s_i, t_i, m_{s_i}, m_{t_i}} - a_{s_i, t_i, m_{s_i}-1, m_{t_i}}, \\ \nabla_2 a_{s_i, t_i, m_{s_i}, m_{t_i}} &= a_{s_i, t_i, m_{s_i}, m_{t_i}} - a_{s_i, t_i, m_{s_i}, m_{t_i}-1}, \\ \nabla_2 \nabla_1 a_{s_i, t_i, m_{s_i}, m_{t_i}} &= \nabla_2 (\nabla_1 a_{s_i, t_i, m_{s_i}, m_{t_i}}). \end{aligned} \tag{1.10}$$

Then

$$\begin{aligned} &\sum_{m_{s_1}=1}^{k_{s_1}} \sum_{m_{t_1}=1}^{k_{t_1}} \cdots \sum_{m_{s_n}=1}^{k_{s_n}} \sum_{m_{t_n}=1}^{k_{t_n}} \frac{\prod_{i=1}^n \phi_i(a_{s_i, t_i, m_{s_i}, m_{t_i}})}{\left((1/\alpha') \sum_{i=1}^n \alpha'_i m_{s_i} m_{t_i} \right)^{\alpha'}} \\ &\geq C(k_{s_1} k_{t_1}, \dots, k_{s_n} k_{t_n}) \\ &\times \prod_{i=1}^n \left(\sum_{m_{s_i}=1}^{k_{s_i}} \sum_{m_{t_i}=1}^{k_{t_i}} (k_{s_i} - m_{s_i} + 1) (k_{t_i} - m_{t_i} + 1) \left(p_{s_i, t_i, m_{s_i}, m_{t_i}} \phi_i \left(\frac{\nabla_2 \nabla_1 a_{s_i, t_i, m_{s_i}, m_{t_i}}}{P_{s_i, t_i, m_{s_i}, m_{t_i}}} \right) \right)^{1/\alpha_i} \right)^{\alpha_i}, \end{aligned} \tag{1.11}$$

4 Inverses of new Hilbert-Pachpatte-type inequalities

where

$$C(k_{s_1}k_{t_1}, \dots, k_{s_n}k_{t_n}) = \prod_{i=1}^n \left(\sum_{m_{s_i}=1}^{k_{s_i}} \sum_{m_{t_i}=1}^{k_{t_i}} \left(\frac{\phi_i(P_{s_i, t_i, m_{s_i}, m_{t_i}})}{P_{s_i, t_i, m_{s_i}, m_{t_i}}} \right)^{1/\alpha'_i} \right)^{\alpha'_i}. \quad (1.12)$$

THEOREM 1.6. Let $f_i(s_i, t_i)$ ($i = 1, 2, \dots, n$) be real-valued continuous functions defined on $[0, x_i] \times [0, y_i]$, where $x_i \in (0, \infty)$, $y_i \in (0, \infty)$, and with $f_i(0, t_i) = f_i(s_i, 0) = 0$ ($i = 1, 2, \dots, n$). Let $p_i(\sigma_i)$ and $q_i(\tau_i)$ be positive continuous functions defined for $\sigma_i \in (0, s_i)$, $\tau_i \in (0, t_i)$. Set

$$P_i(s_i, t_i) = \int_0^{t_i} \int_0^{s_i} p_i(\sigma_i) q_i(\tau_i) d\sigma_i d\tau_i. \quad (1.13)$$

For the function $f_i(s_i, t_i)$, denote the partial derivatives $(\partial/\partial s_i) f_i(s_i, t_i)$, $(\partial/\partial t_i) f_i(s_i, t_i)$, and $(\partial^2/\partial s_i \partial t_i) f_i(s_i, t_i)$ by $D_1 f_i(s_i, t_i)$, $D_2 f_i(s_i, t_i)$, and $D_2 D_1 f_i(s_i, t_i) = D_1 D_2 f_i(s_i, t_i)$, respectively. Let ϕ_i , α_i , α'_i , α , and α' be as in Theorem 1.5. Then

$$\begin{aligned} & \int_0^{x_1} \int_0^{y_1} \cdots \int_0^{x_n} \int_0^{y_n} \prod_{i=1}^n \frac{\phi_i(f_i(s_i, t_i))}{((1/\alpha') \sum_{i=1}^n \alpha'_i s_i t_i)^{\alpha'}} ds_1 dt_1 \cdots ds_n dt_n \\ &= G(x_1 y_1, \dots, x_n y_n) \\ & \times \prod_{i=1}^n \left(\int_0^{x_i} \int_0^{y_i} (x_i - s_i)(y_i - t_i) \left(p_i(s_i) q_i(t_i) \cdot \phi_i \left(\frac{D_2 D_1 f_i(s_i, t_i)}{p_i(s_i) q_i(t_i)} \right) \right)^{1/\alpha_i} ds_i dt_i \right)^{\alpha_i}, \end{aligned} \quad (1.14)$$

where

$$G(x_1 y_1, \dots, x_n y_n) = \prod_{i=1}^n \left(\int_0^{x_i} \int_0^{y_i} \left(\frac{\phi_i(P_i(s_i, t_i))}{P_i(s_i, t_i)} \right)^{1/\alpha'_i} ds_i dt_i \right)^{\alpha'_i}. \quad (1.15)$$

These results provide new estimates on these types of inequalities.

2. Proofs and remarks

Proof of Theorem 1.5. From the operators defined in Theorem 1.5, we obtain

$$\begin{aligned} a_{s_i, t_i, m_{s_i}, m_{t_i}} &= \sum_{m_{\xi_i}=1}^{m_{s_i}} \{ a_{s_i, t_i, m_{\xi_i}, m_{t_i}} - a_{s_i, t_i, m_{\xi_i}-1, m_{t_i}} \} \\ &= \sum_{m_{\xi_i}=1}^{m_{s_i}} \sum_{m_{\eta_i}=1}^{m_{t_i}} \{ a_{s_i, t_i, m_{\xi_i}, m_{\eta_i}} - a_{s_i, t_i, m_{\xi_i}-1, m_{\eta_i}} - a_{s_i, t_i, m_{\xi_i}, m_{\eta_i}-1} + a_{s_i, t_i, m_{\xi_i}-1, m_{\eta_i}-1} \} \\ &= \sum_{m_{\xi_i}=1}^{m_{s_i}} \sum_{m_{\eta_i}=1}^{m_{t_i}} \nabla_2 \nabla_1 a_{s_i, t_i, m_{\xi_i}, m_{\eta_i}}. \end{aligned} \quad (2.1)$$

From (2.1) and the hypotheses of Theorem 1.5, and in view of Jensen's inequality and inverse Hölder's inequality [6], we obtain

$$\begin{aligned}
& \phi_i(a_{s_i, t_i, m_{s_i}, m_{t_i}}) \\
&= \phi_i\left(\frac{P_{s_i, t_i, m_{s_i}, m_{t_i}} \sum_{m_{\xi_i}=1}^{m_{s_i}} \sum_{m_{\eta_i}=1}^{m_{t_i}} P_{s_i, t_i, m_{\xi_i}, m_{\eta_i}} (\nabla_2 \nabla_1 a_{s_i, t_i, m_{\xi_i}, m_{\eta_i}} / p_{s_i, t_i, m_{\xi_i}, m_{\eta_i}})}{\sum_{m_{\xi_i}=1}^{m_{s_i}} \sum_{m_{\eta_i}=1}^{m_{t_i}} P_{s_i, t_i, m_{\xi_i}, m_{\eta_i}}}\right) \\
&\geq \phi_i(P_{s_i, t_i, m_{s_i}, m_{t_i}}) \cdot \phi_i\left(\frac{\sum_{m_{\xi_i}=1}^{m_{s_i}} \sum_{m_{\eta_i}=1}^{m_{t_i}} P_{s_i, t_i, m_{\xi_i}, m_{\eta_i}} (\nabla_2 \nabla_1 a_{s_i, t_i, m_{\xi_i}, m_{\eta_i}} / p_{s_i, t_i, m_{\xi_i}, m_{\eta_i}})}{\sum_{m_{\xi_i}=1}^{m_{s_i}} \sum_{m_{\eta_i}=1}^{m_{t_i}} P_{s_i, t_i, m_{\xi_i}, m_{\eta_i}}}\right) \\
&\geq \frac{\phi_i(P_{s_i, t_i, m_{s_i}, m_{t_i}})}{P_{s_i, t_i, m_{s_i}, m_{t_i}}} \left(\sum_{m_{\xi_i}=1}^{m_{s_i}} \sum_{m_{\eta_i}=1}^{m_{t_i}} P_{s_i, t_i, m_{\xi_i}, m_{\eta_i}} \phi_i\left(\frac{\nabla_2 \nabla_1 a_{s_i, t_i, m_{\xi_i}, m_{\eta_i}}}{P_{s_i, t_i, m_{\xi_i}, m_{\eta_i}}}\right)\right) \\
&\geq \frac{\phi_i(P_{s_i, t_i, m_{s_i}, m_{t_i}})}{P_{s_i, t_i, m_{s_i}, m_{t_i}}} (m_{s_i} m_{t_i})^{\alpha'_i} \times \left(\sum_{m_{\xi_i}=1}^{m_{s_i}} \sum_{m_{\eta_i}=1}^{m_{t_i}} \left(P_{s_i, t_i, m_{\xi_i}, m_{\eta_i}} \phi_i\left(\frac{\nabla_2 \nabla_1 a_{s_i, t_i, m_{\xi_i}, m_{\eta_i}}}{P_{s_i, t_i, m_{\xi_i}, m_{\eta_i}}}\right)\right)^{1/\alpha_i}\right)^{\alpha_i}. \tag{2.2}
\end{aligned}$$

On the other hand, noticing $\alpha'_i < 0$ ($i = 1, 2, \dots, n$), $\lambda_i > 0$ ($i = 1, 2, \dots, n$), and applying the well-known means inequality, we have

$$\prod_{i=1}^n \lambda_i^{\alpha'_i} \geq \left(\frac{1}{\alpha'} \sum_{i=1}^n \alpha'_i \lambda_i\right)^{\alpha'}. \tag{2.3}$$

Hence

$$\begin{aligned}
& \frac{\prod_{i=1}^n \phi_i(a_{s_i, t_i, m_{s_i}, m_{t_i}})}{((1/\alpha') \sum_{i=1}^n \alpha'_i m_{s_i} m_{t_i})^{\alpha'}} \\
&\geq \prod_{i=1}^n \frac{\phi_i(P_{s_i, t_i, m_{s_i}, m_{t_i}})}{P_{s_i, t_i, m_{s_i}, m_{t_i}}} \left(\sum_{m_{\xi_i}=1}^{m_{s_i}} \sum_{m_{\eta_i}=1}^{m_{t_i}} \left(P_{s_i, t_i, m_{\xi_i}, m_{\eta_i}} \phi_i\left(\frac{\nabla_2 \nabla_1 a_{s_i, t_i, m_{\xi_i}, m_{\eta_i}}}{P_{s_i, t_i, m_{\xi_i}, m_{\eta_i}}}\right)\right)^{1/\alpha_i}\right)^{\alpha_i}. \tag{2.4}
\end{aligned}$$

Taking the sum of both sides of (2.4) over m_{s_i}, m_{t_i} ($i = 1, 2, \dots, n$) from 1 to k_{s_i}, k_{t_i} , respectively, and in view of inverse Hölder's inequality, we get

$$\begin{aligned}
& \sum_{m_{s_1}=1}^{k_{s_1}} \sum_{m_{t_1}=1}^{k_{t_1}} \cdots \sum_{m_{s_n}=1}^{k_{s_n}} \sum_{m_{t_n}=1}^{k_{t_n}} \frac{\prod_{i=1}^n \phi_i(a_{s_i, t_i, m_{s_i}, m_{t_i}})}{((1/\alpha') \sum_{i=1}^n \alpha'_i m_{s_i} m_{t_i})^{\alpha'}} \\
&\geq \prod_{i=1}^n \sum_{m_{s_i}=1}^{k_{s_i}} \sum_{m_{t_i}=1}^{k_{t_i}} \left(\frac{\phi_i(P_{s_i, t_i, m_{s_i}, m_{t_i}})}{P_{s_i, t_i, m_{s_i}, m_{t_i}}}\right) \left(\sum_{m_{\xi_i}=1}^{m_{s_i}} \sum_{m_{\eta_i}=1}^{m_{t_i}} \left(P_{s_i, t_i, m_{\xi_i}, m_{\eta_i}} \phi_i\left(\frac{\nabla_2 \nabla_1 a_{s_i, t_i, m_{\xi_i}, m_{\eta_i}}}{P_{s_i, t_i, m_{\xi_i}, m_{\eta_i}}}\right)\right)^{1/\alpha_i}\right)^{\alpha_i}
\end{aligned}$$

6 Inverses of new Hilbert-Pachpatte-type inequalities

$$\begin{aligned}
&\geq \prod_{i=1}^n \left(\sum_{m_{s_i}=1}^{k_{s_i}} \sum_{m_{t_i}=1}^{k_{t_i}} \left(\frac{\phi_i(P_{s_i,t_i,m_{s_i},m_{t_i}})}{P_{s_i,t_i,m_{s_i},m_{t_i}}} \right)^{1/\alpha'_i} \right)^{\alpha'_i} \\
&\quad \times \left(\sum_{m_{s_i}=1}^{k_{s_i}} \sum_{m_{t_i}=1}^{k_{t_i}} \left(\sum_{m_{\xi_i}=1}^{m_{s_i}} \sum_{m_{\eta_i}=1}^{m_{t_i}} \left(p_{s_i,t_i,m_{\xi_i},m_{\eta_i}} \phi_i \left(\frac{\nabla_2 \nabla_1 a_{s_i,t_i,m_{\xi_i},m_{\eta_i}}}{p_{s_i,t_i,m_{\xi_i},m_{\eta_i}}} \right) \right)^{1/\alpha_i} \right)^{\alpha_i} \\
&= C(k_{s_1} k_{t_1}, \dots, k_{s_n} k_{t_n}) \\
&\quad \times \prod_{i=1}^n \left(\sum_{m_{\xi_i}=1}^{k_{s_i}} \sum_{m_{\eta_i}=1}^{k_{t_i}} (k_{s_i} - m_{\xi_i} + 1)(k_{t_i} - m_{\eta_i} + 1) \left(p_{s_i,t_i,m_{\xi_i},m_{\eta_i}} \phi_i \left(\frac{\nabla_2 \nabla_1 a_{s_i,t_i,m_{\xi_i},m_{\eta_i}}}{p_{s_i,t_i,m_{\xi_i},m_{\eta_i}}} \right) \right)^{1/\alpha_i} \right)^{\alpha_i} \\
&= C(k_{s_1} k_{t_1}, \dots, k_{s_n} k_{t_n}) \\
&\quad \times \prod_{i=1}^n \left(\sum_{m_{s_i}=1}^{k_{s_i}} \sum_{m_{t_i}=1}^{k_{t_i}} (k_{s_i} - m_{s_i} + 1)(k_{t_i} - m_{t_i} + 1) \left(p_{s_i,t_i,m_{s_i},m_{t_i}} \phi_i \left(\frac{\nabla_2 \nabla_1 a_{s_i,t_i,m_{s_i},m_{t_i}}}{p_{s_i,t_i,m_{s_i},m_{t_i}}} \right) \right)^{1/\alpha_i} \right)^{\alpha_i}. \tag{2.5}
\end{aligned}$$

This completes the proof. \square

Remark 2.1. Let $a_{s_i,t_i,m_{s_i},m_{t_i}}$, m_{s_i} , m_{t_i} , k_{s_i} , k_{t_i} , $a_{s_i,t_i,0,m_{t_i}}$, $a_{s_i,t_i,m_{s_i},0}$, $P_{s_i,t_i,m_{s_i},m_{t_i}}$, ∇_1 , and ∇_2 be as in Theorem 1.5. Let ϕ_i , α_i , α'_i , α , and α' ($i = 1, 2, \dots, n$) be as in Theorem 1.3. Similar to the proof of Theorem 1.5, we obtain

$$\begin{aligned}
&\sum_{m_{s_1}=1}^{k_{s_1}} \sum_{m_{t_1}=1}^{k_{t_1}} \cdots \sum_{m_{s_n}=1}^{k_{s_n}} \sum_{m_{t_n}=1}^{k_{t_n}} \frac{\prod_{i=1}^n \phi_i(a_{s_i,t_i,m_{s_i},m_{t_i}})}{(\sum_{i=1}^n \alpha'_i m_{s_i} m_{t_i})^{\alpha'}} \\
&\leq \bar{C}(k_{s_1} k_{t_1}, \dots, k_{s_n} k_{t_n}) \\
&\quad \times \prod_{i=1}^n \left(\sum_{m_{s_i}=1}^{k_{s_i}} \sum_{m_{t_i}=1}^{k_{t_i}} (k_{s_i} - m_{s_i} + 1)(k_{t_i} - m_{t_i} + 1) \left(p_{s_i,t_i,m_{s_i},m_{t_i}} \phi_i \left(\frac{\nabla_2 \nabla_1 a_{s_i,t_i,m_{s_i},m_{t_i}}}{p_{s_i,t_i,m_{s_i},m_{t_i}}} \right) \right)^{1/\alpha_i} \right)^{\alpha_i}, \tag{2.6}
\end{aligned}$$

where

$$\bar{C}(k_{s_1} k_{t_1}, \dots, k_{s_n} k_{t_n}) = \frac{1}{(\alpha')^{\alpha'}} \prod_{i=1}^n \left(\sum_{m_{s_i}=1}^{k_{s_i}} \sum_{m_{t_i}=1}^{k_{t_i}} \left(\frac{\phi_i(P_{s_i,t_i,m_{s_i},m_{t_i}})}{P_{s_i,t_i,m_{s_i},m_{t_i}}} \right)^{1/\alpha'_i} \right)^{\alpha'_i}. \tag{2.7}$$

This is just a *general form* of inequality (1.5) and the *inverse form* of inequality (1.11).

Moreover, let $\{a_{i,m_i}\}$, m_i , k_i , $a_{i,0}$, p_{i,m_i} , ∇ , and P_{i,m_i} ($i = 1, 2, \dots, n$) be as in Theorem 1.3. Let ϕ_i , α_i , α'_i , α , and α' ($i = 1, 2, \dots, n$) be as in Theorem 1.5. Similar to the proof of Theorem 1.5, we obtain

$$\sum_{m_1=1}^{k_1} \dots \sum_{m_n=1}^{k_n} \frac{\prod_{i=1}^n \phi_i(a_{i,m_i})}{((1/\alpha') \sum_{i=1}^n \alpha'_i m_i)^{\alpha'}} \geq \overline{M}(k_1, \dots, k_n) \prod_{i=1}^n \left(\sum_{m_i=1}^{k_i} (k_i - m_i + 1) \left(p_{i,m_i} \phi_i \left(\frac{\nabla a_{i,m_i}}{p_{i,m_i}} \right) \right)^{1/\alpha_i} \right)^{\alpha_i}, \tag{2.8}$$

where

$$\overline{M}(k_1, \dots, k_n) = \prod_{i=1}^n \left(\sum_{m_i=1}^{k_i} \left(\frac{\phi_i(P_{i,m_i})}{P_{i,m_i}} \right)^{1/\alpha'_i} \right)^{\alpha'_i}. \tag{2.9}$$

This is just an *inverse form* of inequality (1.5) which was given by Handley et al. [4]. Let $n = 2$, $\alpha_1 = \alpha_2 = 2$, then $\alpha'_1 = \alpha'_2 = -1$, putting them in inequality (2.8), we have

$$\sum_{m_1=1}^{k_1} \sum_{m_2=1}^{k_2} \frac{\phi_1(a_{1,m_1}) \phi_2(a_{2,m_2})}{(m_1 + m_2)^{-2}} \geq \overline{M}(k_1, k_2) \left(\sum_{m_1=1}^{k_1} (k_1 - m_1 + 1) \left(p_{1,m_1} \phi_1 \left(\frac{\nabla a_{1,m_1}}{p_{1,m_1}} \right) \right)^{1/2} \right)^2 \times \left(\sum_{m_2=1}^{k_2} (k_2 - m_2 + 1) \left(p_{2,m_2} \phi_2 \left(\frac{\nabla a_{2,m_2}}{p_{2,m_2}} \right) \right)^{1/2} \right)^2, \tag{2.10}$$

where

$$\overline{M}(k_1, k_2) = 4 \left(\sum_{m_1=1}^{k_1} \left(\frac{\phi_1(P_{1,m_1})}{P_{1,m_1}} \right)^{-1} \right)^{-1} \left(\sum_{m_2=1}^{k_2} \left(\frac{\phi_2(P_{2,m_2})}{P_{2,m_2}} \right)^{-1} \right)^{-1}. \tag{2.11}$$

This is just an *inverse form* of inequality (1.1) which was given by Pachpatte [9].

Proof of Theorem 1.6. From the hypotheses of Theorem 1.6, we obtain

$$f_i(s_i, t_i) = \int_0^{s_i} \int_0^{t_i} D_2 D_1 f_i(\sigma_i, \tau_i) d\sigma_i d\tau_i. \tag{2.12}$$

8 Inverses of new Hilbert-Pachpatte-type inequalities

From (2.12) and using Jensen's integral inequality and Hölder's integral inequality [6], we obtain

$$\begin{aligned}
 \phi_i(f_i(s_i, t_i)) &= \phi_i\left(\frac{P_i(s_i, t_i) \int_0^{t_i} \int_0^{s_i} p_i(\sigma_i) q_i(\tau_i) (D_2 D_1 f_i(\sigma_i, \tau_i) / p_i(\sigma_i) q_i(\tau_i)) d\sigma_i d\tau_i}{\int_0^{t_i} \int_0^{s_i} p_i(\sigma_i) q_i(\tau_i) d\sigma_i d\tau_i}\right) \\
 &\geq \phi_i(P_i(s_i, t_i)) \cdot \phi_i\left(\frac{\int_0^{t_i} \int_0^{s_i} p_i(\sigma_i) q_i(\tau_i) (D_2 D_1 f_i(\sigma_i, \tau_i) / p_i(\sigma_i) q_i(\tau_i)) d\sigma_i d\tau_i}{\int_0^{t_i} \int_0^{s_i} p_i(\sigma_i) q_i(\tau_i) d\sigma_i d\tau_i}\right) \\
 &\geq \frac{\phi_i(P_i(s_i, t_i))}{P_i(s_i, t_i)} \int_0^{t_i} \int_0^{s_i} p_i(\sigma_i) q_i(\tau_i) \cdot \phi_i\left(\frac{D_2 D_1 f_i(\sigma_i, \tau_i)}{p_i(\sigma_i) q_i(\tau_i)}\right) d\sigma_i d\tau_i \\
 &\geq \frac{\phi_i(P_i(s_i, t_i))}{P_i(s_i, t_i)} (s_i t_i)^{\alpha'_i} \left(\int_0^{t_i} \int_0^{s_i} \left(p_i(\sigma_i) q_i(\tau_i) \cdot \phi_i\left(\frac{D_2 D_1 f_i(\sigma_i, \tau_i)}{p_i(\sigma_i) q_i(\tau_i)}\right)\right)^{1/\alpha_i} d\sigma_i d\tau_i\right)^{\alpha_i}.
 \end{aligned} \tag{2.13}$$

From the well-known inequality for means, we have

$$\begin{aligned}
 \prod_{i=1}^n \frac{\phi_i(f_i(s_i, t_i))}{((1/\alpha') \sum_{i=1}^n \alpha'_i s_i t_i)^{\alpha'}} &\geq \prod_{i=1}^n \frac{\phi_i(P_i(s_i, t_i))}{P_i(s_i, t_i)} \left(\int_0^{t_i} \int_0^{s_i} \left(p_i(\sigma_i) q_i(\tau_i) \cdot \phi_i\left(\frac{D_2 D_1 f_i(\sigma_i, \tau_i)}{p_i(\sigma_i) q_i(\tau_i)}\right)\right)^{1/\alpha_i} d\sigma_i d\tau_i\right)^{\alpha_i}.
 \end{aligned} \tag{2.14}$$

Integrating both sides of (2.14) over s_i, t_i from 1 to x_i, y_i ($i = 1, 2, \dots, n$) and in view of Hölder's integral inequality and Fubini's theorem, we observe that

$$\begin{aligned}
 &\int_0^{x_1} \int_0^{y_1} \cdots \int_0^{x_n} \int_0^{y_n} \prod_{i=1}^n \frac{\phi_i(f_i(s_i, t_i))}{((1/\alpha') \sum_{i=1}^n \alpha'_i s_i t_i)^{\alpha'}} ds_1 dt_1 \cdots ds_n dt_n \\
 &\geq \prod_{i=1}^n \int_0^{x_i} \int_0^{y_i} \frac{\phi_i(P_i(s_i, t_i))}{P_i(s_i, t_i)} \\
 &\quad \times \left(\int_0^{t_i} \int_0^{s_i} \left(p_i(\sigma_i) q_i(\tau_i) \cdot \phi_i\left(\frac{D_2 D_1 f_i(\sigma_i, \tau_i)}{p_i(\sigma_i) q_i(\tau_i)}\right)\right)^{1/\alpha_i} d\sigma_i d\tau_i\right)^{\alpha_i} ds_i dt_i \\
 &\geq \prod_{i=1}^n \left(\int_0^{x_i} \int_0^{y_i} \left(\frac{\phi_i(P_i(s_i, t_i))}{P_i(s_i, t_i)}\right)^{1/\alpha'_i} ds_i dt_i\right)^{\alpha'_i} \\
 &\quad \times \left(\int_0^{x_i} \int_0^{y_i} \left(\int_0^{t_i} \int_0^{s_i} \left(p_i(\sigma_i) q_i(\tau_i) \cdot \phi_i\left(\frac{D_2 D_1 f_i(\sigma_i, \tau_i)}{p_i(\sigma_i) q_i(\tau_i)}\right)\right)^{1/\alpha_i} d\sigma_i d\tau_i\right)^{\alpha_i} ds_i dt_i\right)^{\alpha_i}
 \end{aligned}$$

$$\begin{aligned}
 &= G(x_1 y_1, \dots, x_n y_n) \\
 &\quad \times \prod_{i=1}^n \left(\int_0^{x_i} \int_0^{y_i} (x_i - s_i)(y_i - t_i) \left(p_i(s_i) q_i(t_i) \cdot \phi_i \left(\frac{D_2 D_1 f_i(s_i, t_i)}{p_i(s_i) q_i(t_i)} \right) \right)^{1/\alpha_i} ds_i dt_i \right)^{\alpha_i}.
 \end{aligned} \tag{2.15}$$

The proof is complete. □

Remark 2.2. Let $f_i(s_i, t_i)$, x_i , y_i , $f_i(0, t_i)$, $f_i(s_i, 0)$, $D_1 f_i(s_i, t_i)$, $D_2 f_i(s_i, t_i)$, $D_2 D_1 f_i(s_i, t_i)$, $p_i(\sigma_i)$, $q_i(\tau_i)$, and $P_i(s_i, t_i)$ be as in Theorem 1.6. Let ϕ_i , α_i , α'_i , α , and α' be as in Theorem 1.3. Similar to the proof of Theorem 1.6, we have

$$\begin{aligned}
 &\int_0^{x_1} \int_0^{y_1} \cdots \int_0^{x_n} \int_0^{y_n} \prod_{i=1}^n \frac{\phi_i(f_i(s_i, t_i))}{(\sum_{i=1}^n \alpha'_i s_i t_i)^{\alpha'}} ds_1 dt_1 \cdots ds_n dt_n \\
 &\leq \bar{G}(x_1 y_1, \dots, x_n y_n) \\
 &\quad \times \prod_{i=1}^n \left(\int_0^{x_i} \int_0^{y_i} (x_i - s_i)(y_i - t_i) \left(p_i(s_i) q_i(t_i) \cdot \phi_i \left(\frac{D_2 D_1 f_i(s_i, t_i)}{p_i(s_i) q_i(t_i)} \right) \right)^{1/\alpha_i} ds_i dt_i \right)^{\alpha_i},
 \end{aligned} \tag{2.16}$$

where

$$\bar{G}(x_1 y_1, \dots, x_n y_n) = \frac{1}{(\alpha')^{\alpha'}} \prod_{i=1}^n \left(\int_0^{x_i} \int_0^{y_i} \left(\frac{\phi_i(P_i(s_i, t_i))}{P_i(s_i, t_i)} \right)^{1/\alpha'_i} ds_i dt_i \right)^{\alpha'_i}. \tag{2.17}$$

This is just a *general form* of inequality (1.7) and an *inverse form* of the inequality (1.14).

On the other hand, let f_i , $f_i(0)$, $f'_i(s_i)$, $p_i(\sigma_i)$, σ_i , k_i , s_i , x_i , $P_i(s_i)$ be as in Theorem 1.4, and let ϕ_i , α_i , α'_i , α , α' be as in Theorem 1.5. Then

$$\begin{aligned}
 &\int_0^{x_1} \cdots \int_0^{x_n} \frac{\prod_{i=1}^n \phi_i(f_i(s_i))}{(1/\alpha')(\sum_{i=1}^n \alpha'_i s_i)^{\alpha'}} ds_1 \cdots ds_n \\
 &\geq \bar{L}(x_1, \dots, x_n) \prod_{i=1}^n \left(\int_0^{x_i} (x_i - s_i) \left(p_i(s_i) \phi_i \left(\frac{f'_i(s_i)}{P_i(s_i)} \right) \right)^{1/\alpha_i} ds_i \right)^{\alpha_i},
 \end{aligned} \tag{2.18}$$

where

$$\bar{L}(x_1, \dots, x_n) = \prod_{i=1}^n \left(\int_0^{x_i} \left(\frac{\phi_i(P_i(s_i))}{P_i(s_i)} \right)^{1/\alpha'_i} ds_i \right)^{\alpha'_i}. \tag{2.19}$$

This is just an *inverse form* of inequality (1.7) which was given by Handley et al. [4].

10 Inverses of new Hilbert-Pachpatte-type inequalities

Let $n = 2$, $\alpha_1 = \alpha_2 = 2$, then $\alpha'_1 = \alpha'_2 = -1$, taking them in inequality (2.18), we have

$$\begin{aligned} & \int_0^{x_1} \int_0^{x_2} \frac{\phi_1(f_1(s_1))\phi_2(f_2(s_2))}{(s_1 + s_2)^{-2}} ds_1 ds_2 \\ & \geq \bar{L}(x_1, x_2) \left(\int_0^{x_1} (x_1 - s_1) \left(p_1(s_1) \phi_1 \left(\frac{f'_2(s_1)}{p_1(s_1)} \right) \right)^{1/2} ds_1 \right)^2 \\ & \quad \times \left(\int_0^{x_2} (x_2 - s_2) \left(p_2(s_2) \phi_2 \left(\frac{f'_2(s_2)}{p_2(s_2)} \right) \right)^{1/2} ds_2 \right)^2, \end{aligned} \quad (2.20)$$

where

$$\bar{L}(x_1, x_2) = 4 \left(\int_0^{x_1} \left(\frac{\phi_1(P_1(s_1))}{P_1(s_1)} \right)^{-1} ds_1 \right)^{-1} \left(\int_0^{x_2} \left(\frac{\phi_2(P_2(s_2))}{P_2(s_2)} \right)^{-1} ds_2 \right)^{-1}. \quad (2.21)$$

This is just an *inverse form* of inequality (1.3) which was given by Pachpatte [9].

For interrelated research on similar inequalities, one is referred to [2, 8, 11, 14, 15], and the references cited therein.

Acknowledgments

Research of the first author is supported by Zhejiang Provincial Natural Science Foundation of China (Y605065), National Natural Sciences Foundation of China (10271071), Foundation of the Education Department of Zhejiang Province of China (20050392), and the Academic Mainstay of Middle-age and Youth Foundation of Shandong Province of China (200203). Research of the second author is partially supported by the Research Grants Council of the Hong Kong SAR, China (Project no. HKU7040/03P, and HKU 7017/05P).

References

- [1] W.-S. Cheung, Ž. Hanjŝ, and J. Pečarić, *Some Hardy-type inequalities*, Journal of Mathematical Analysis and Applications **250** (2000), no. 2, 621–634.
- [2] W.-S. Cheung and Q.-H. Ma, *On certain new Gronwall-Ou-Iang type integral inequalities in two variables and their applications*, Journal of Inequalities and Applications **2005** (2005), no. 4, 347–361.
- [3] M. Gao and B. Yang, *On the extended Hilbert's inequality*, Proceedings of the American Mathematical Society **126** (1998), no. 3, 751–759.
- [4] G. D. Handley, J. J. Koliha, and J. E. Pečarić, *A Hilbert type inequality*, Tamkang Journal of Mathematics **31** (2000), no. 4, 311–315.
- [5] ———, *New Hilbert-Pachpatte type integral inequalities*, Journal of Mathematical Analysis and Applications **257** (2001), no. 1, 238–250.
- [6] G. H. Hardy, J. E. Littlewood, and G. Pólya, *Inequalities*, Cambridge University Press, Cambridge, 1934.
- [7] K. Jichang, *On new extensions of Hilbert's integral inequality*, Journal of Mathematical Analysis and Applications **235** (1999), no. 2, 608–614.

- [8] C. A. Okpoti, L.-E. Persson, and A. Wedestig, *Weight characterizations for the discrete Hardy inequality with kernel*, Journal of Inequalities and Applications **2006** (2006), Article ID 18030, 14 pages.
- [9] B. G. Pachpatte, *A note on Hilbert type inequality*, Tamkang Journal of Mathematics **29** (1998), no. 4, 293–298.
- [10] ———, *On some new inequalities similar to Hilbert’s inequality*, Journal of Mathematical Analysis and Applications **226** (1998), no. 1, 166–179.
- [11] P. Řehák, *Hardy inequality on time scales and its application to half-linear dynamic equations*, Journal of Inequalities and Applications **2005** (2005), no. 5, 495–507.
- [12] B. Yang, *On new generalizations of Hilbert’s inequality*, Journal of Mathematical Analysis and Applications **248** (2000), no. 1, 29–40.
- [13] B. Yang and T. M. Rassias, *On the way of weight coefficient and research for the Hilbert-type inequalities*, Mathematical Inequalities & Applications **6** (2003), no. 4, 625–658.
- [14] S. Yaotian and C. Zhihui, *General Hardy inequalities with optimal constants and remainder terms*, Journal of Inequalities and Applications **2005** (2005), no. 3, 207–219.
- [15] J. Yuan and G. S. Leng, *Inequalities for dual affine quermassintegrals*, Journal of Inequalities and Applications **2006** (2006), Article ID 50181, 7 pages.
- [16] C.-J. Zhao, *Generalization on two new Hilbert type inequalities*, Journal of Mathematics **20** (2000), no. 4, 413–416.
- [17] ———, *Inverses of disperse and continuous Pachpatte’s inequalities*, Acta Mathematica Sinica **46** (2003), no. 6, 1111–1116.
- [18] C.-J. Zhao and L. Debnath, *Some new inverse type Hilbert integral inequalities*, Journal of Mathematical Analysis and Applications **262** (2001), no. 1, 411–418.

Chang-Jian Zhao: Department of Information and Mathematics Sciences, College of Science,
China Jiliang University, Hangzhou 310018, China
E-mail address: chjzhao@cjlu.edu.cn

Wing-Sum Cheung: Department of Mathematics, The University of Hong Kong, Pokfulam Road,
Hong Kong
E-mail address: wscheung@hku.hk