

Article

Synthesis and Calcium Mobilization Activity of cADPR Analogues Which Integrate Nucleobase, Northern and Southern Ribose Modifications

Yue Zhou ¹, Peilin Yu ¹, Hongwei Jin ¹, Zhenjun Yang ¹, Jianbo Yue ², Liangren Zhang ^{1,*} and Lihe Zhang ¹

¹ State Key Laboratory of Natural and Biomimetic Drugs, School of Pharmaceutical Sciences, Peking University, Beijing 100191, China

² Department of Physiology, University of Hong Kong, Hong Kong, China

* Author to whom correspondence should be addressed; E-Mail: liangren@bjmu.edu.cn; Tel: +86-10-8280-2567, Fax: +86-10-8280-5063.

Received: 24 February 2012; in revised form: 31 March 2012 / Accepted: 5 April 2012 / Published: 10 April 2012

Abstract: Novel cADPR mimics, which integrate nucleobase, northern and southern ribose modifications were synthesized. The key steps of the synthesis were a Cu(I)-catalyzed Huisgen [3+2] cycloaddition and a microwave-assisted intramolecular pyrophosphorylation. Preliminary biological investigations showed that these cADPR mimics are membrane-permeating agonists of the calcium signaling pathway. The introduction of chlorine or fluorine at the 2'-position of the southern riboses led to a decrease of activity. The existence of a hydrophobic group on the 3'-OH of the southern riboses does not obviously alter the agonistic activity.


Keywords: cADPR analogue; nucleotide; synthesis; calcium mobilization

1. Introduction

Cyclic adenosine disphosphate ribose (cADPR, **1**, Figure 1) is a universal Ca²⁺ mobilizing secondary messenger first identified in the sea urchin egg system [1]. Since its discovery, numerous cell systems that utilize the cADPR/ryanodine receptor (RyR) Ca²⁺ signaling system to control Ca²⁺-dependent cellular responses, such as fertilization, secretion, contraction, proliferation and so on have been described [2,3]. A large number of proteins are involved in specifically shaping Ca²⁺ signals,

however, it is still unclear exactly how cADPR elicits calcium release, which includes whether specific cADPR binding proteins exist. Since the discovery of cADPR, a number of structurally diverse cADPR analogues have been synthesized and used to elucidate the molecular mechanism of calcium signaling [4–6]. The syntheses could be divided into chemo-enzymatic or chemical synthesis [7,8]. The structures include modifications of the pyrophosphate [9–12], purine [13–16] and southern and northern riboses [17–20] of cADPR. These analogues can agonize or antagonize cADPR/RyR calcium signal pathway.

Figure 1. Structures of cADPR analogues.



cADPR analogues with modifications on the northern and southern riboses are the most investigated. Shuto *et al.* found that the effect of modification of the northern ribose on calcium signaling depended on the cell system [21,22]. Our previous studies showed that the northern ribose of cADPR tolerated structural modifications to some extent. The agonistic activity is mostly maintained even if the northern ribose is replaced by ether or alkane linkages, such as in cIDPRE (2), and this modification makes analogues cell-permeant [18]. Potter *et al.* investigated the importance of the 2'-OH on the calcium signaling behavior of 8-substituted cADPR derivatives. They found that the 2'-OH group does not affect the Ca²⁺-mobilizing ability of cADPR itself, but that it is an important motif for the antagonistic activities of 8-substituted cADPR analogues [23]. The C2'-*endo/syn* conformation is crucial for agonistic or antagonistic activity in sea urchin egg homogenates [24]. The findings implied the coordinating effect of nucleobase and riboses on the activity of cADPR analogues.


The nucleobase of cADPR was simplified in our previous study; we have found that triazole-based cADPR analogues 3,4 are cell-permeating mild agonists of the cADPR/RyR calcium pathway [25]. To elucidate the structure-activity relationships of cADPR analogues in more detail, and provide probes for investigation of the molecular mechanism of cADPR regulated calcium pathways, we have designed and synthesized novel cADPR analogues which integrate three types of modifications of the nucleobase, northern and southern riboses (compounds 5). In this study, the nucleobase is replaced by a simplified triazole moiety, the northern ribose is replaced by an ether linkage and the southern ribose is replaced by 2'-deoxy or 2'-deoxy-2'-haloribofuranoses, respectively.

2. Results and Discussion

2.1. Chemistry


The synthesis could be generalized to three steps, *i.e.*, the synthesis of phosphorylated northern and southern moieties, the assembly of the triazole moiety and intramolecular cyclization (Scheme 1). The phosphorylation was performed before the coupling of the northern and southern moieties. Microwave-assisted intramolecular cyclization was used to form the pyrophosphates of the cADPR analogues.

Scheme 1. Retrosynthesis of cADPR analogues.



For the syntheses of the southern moieties, various strategies were adopted to construct 1-azido-2-modified sugars **16**, which are summarized in Scheme 2.

Scheme 2. Synthesis of 1-azido-2-modified sugars.




Reagents and conditions: (a) Ac_2O , pyridine, 1,4-dioxane, reflux, 3 h; (b) SnCl_4 , TMSN_3 , TMSOTf , CH_2Cl_2 , ice-bath; (c) K_2CO_3 , CH_3OH , rt, 12 h; (d) 33% HBr , AcOH , CH_2Cl_2 , rt, 3 h; (e) NaN_3 , DMF , 40 °C, 10 h; (f) $\text{NH}_3/\text{CH}_3\text{OH}$, rt, 10–12 h; (g) BzCl , pyridine, CH_2Cl_2 , ice-bath; (h) TBDMSCl , imidazole, DMF , 60–65 °C; (f) $\text{NH}_3/\text{CH}_3\text{OH}$, rt, 2–3 h.

1- β -D-Azido-2-deoxyribose (**9**) was prepared starting from 2-deoxyribose based on Greenberg's method [26]. The azidation of triacetylribose **7** gave a pair of anomers. The α : β isomer ratio was 1:8 and they could be separated by silica gel column chromatography. The fluoroarabinose **10a** was used directly from commercial material [27]. The chloroarabinose **10b** was synthesized by the triflation of commercial available 1,3,5-tri-*O*-benzoyl- α -D-ribofuranose, followed by treatment with LiCl in *N*-methylpyrrolidinone (NMP) [28]. Treatment of **10** with HBr/CH₃COOH in dichloromethane followed by the addition of sodium azide in DMF at 30 °C for 12 h gave **12** in a yield of 75%. Considering the instability of the anomeric bromide, intermediate **11** was used without purification. The 3-OH groups of **9** and **13** were protected by *tert*-butyldimethyl silyl (TBDMS) groups by reaction with TBDMSCl after their 5-OH groups were selectively protected as benzoyl groups, thus giving **15**. After treatment with NH₃/CH₃OH, compound **16** was obtained.

Cu(I)-catalyzed Huisgen [3+2] cycloaddition between terminal alkyne and azide groups was adopted to build the 1,2,3-triazole unit (Scheme 3). The terminal alkyne building block **17** which carried a *S,S*-diphenylphosphate group was prepared by the reported procedure [29]. Several classical Cu(I) catalysis systems were used to catalyze the Huisgen [3+2] cycloaddition reaction of **16** and **17**. Under optimal conditions, compound **18** was obtained in a yield of 85%. Compound **18** was phosphorylated by using POCl₃/DIPEA in CH₃CN, followed with purification by HPLC eluting with 0.05M triethylammonium bicarbonate (TEAB, pH = 7.5). It was found that the *S,S*-diphenylphosphate group was sensitive to alkaline conditions and was decomposed to *S*-phenylphosphate. Compound **20** was obtained as a triethylammonium salt in a yield of 54% for two steps.

Scheme 3. Synthesis of compound **5**.



Reagents and conditions: (a) CuI, DIPEA, CH₃CN; (b) POCl₃, DIPEA, CH₃CN; (c) 1M TEAB; (d) I₂, pyridine; (e) TBAF, THF, Ac₂O.


Microwave-assisted organic chemistry has been recently developed for the efficient synthesis of functional compounds in many areas [30,31]. In most previous works, intermolecular reactions for the formation of pyrophosphates of cADPR analogues were performed in super-dilute solutions [8]. Under

these super-dilute solution conditions, the reaction was usually completed by using a syringe-pump over 20 h. In addition, the reaction was very sensitive to traces of water. This strictly dry environment and the long reaction time in a super-dilute solution made the experimental work-up tedious, and thus, the large-scale preparation of cADPR analogues was difficult. Under microwave-assisted conditions, the overall efficiency of intramolecular pyrophosphorylation has been greatly improved [29]. We applied this technology to compound **20** to prepare **21**. Optimization of the reaction conditions was done by changing temperature and reaction time. The reaction of **20** with I₂ in pyridine (75 °C/15 min) gave cyclic compound **21** as its triethylammonium salts in a yield of 81%–87%. Finally, the removal of the TBDMS group of **21** was carried out in a solution of 1M TBAF/THF at room temperature for 1 h to obtain the target compound **5**. Compound **5** was identified by ¹H- and ³¹P-NMR, and HRMS.

2.2. Pharmacology

The Ca²⁺-mobilizing ability of the newly synthesized cADPR analogues **5a–c** were evaluated in Jurket-T cells. For studying the effect of the 3'-OH on activity, the 3'-O-TBDMS substituted precursors **21a–c** were also studied. The results are shown in Figure 2.

Figure 2. Ca²⁺ mobilization activities of compounds **21** and **5** in intact Jurkat T cells.



Conditions: The response to the application of the cIDPRE served as a positive control while addition of buffer was used as blank. Jurkat T-cells were loaded with Fura-2AM and subjected to Ca²⁺ imaging. Changes in mean F/F₀ after the application of compounds or buffer control were averaged for at least three independent experiments for each concentration. **(A)** The peak values of F/F₀ were analyzed as the maximum within 200 s after application. **(B)** Time course of Ca²⁺ mobilization by addition of compounds **21a–c** and **5a–c**. The time point's addition of 750 μM positive control, compound **21a–c** and **5a–c** are indicated by arrows.


In general, the synthesized cADPR analogues are membrane-permeating agonists of the calcium signaling pathway. All compounds showed typical biphasic Ca^{2+} mobilizing kinetics with an initial immediate Ca^{2+} peak and a subsequent plateau phase. The induction of the immediate peak was strongest for compound **5a** at 750 μM , which was almost as active as the positive control cIDPRE, while for the other 2'-deoxy cADPR analogues a less pronounced initial Ca^{2+} peak was observed. The result indicates that the introduction of chlorine (**5b**) or fluorine (**5c**) to the 2'-deoxy position will lead to a decrease of activity.

In comparison to cIDPRE, however, 2'-deoxy cADPR analogues **21a–c** and **5a–c** sustained longer Ca^{2+} mobilization (Figure 2B), which implied their different molecular mechanism. 2'-Deoxy-cADPR was found to be inactive in Jurkat T cells [32]. Potter's finding implied that the 2'-OH group is an important motif for the antagonistic activities of 8-substituted cADPR analogues and 2'-OH deletion may also result in an increase in the agonistic effects of the 8-substituted cADPR analogues [23,24]. This study shows that the agonistic activity is maintained in the case of disappearance of the 2'-OH, which emphasizes the importance of the structure of the nucleobase on the effect of the 2'-OH on calcium mobilization.

The existence of a large hydrophobic *tert*-butyldimethylsilyl (TBDMS) group on the 3'-OH of the southern ribose (compounds **21a–c**) lowers activity, but most of the agonistic properties are still maintained. Potter *et al.* reported that the 3'-hydroxyl group is essential for Ca^{2+} releasing activity of cADPR in sea urchin eggs [23]. In our case, TBDMS attached compound **21** did not much alter the agonistic activity of compound **5**. This result may come from the existence of a hydrophobic group on the 3'-OH that makes compound more membrane permeant and partially compensates the negative effect.

The conformations of the modified cADPR analogues were investigated by molecular modeling. Their optimized conformations were superimposed with cADPR (Figure 3). The result clearly shows that **5a–c** and cADPR have very similar conformations. All riboses are in 2'-endo/3'-exo conformations and the whole backbones are in similar alignments. The consistence of conformations might result in the agonistic activities of cADPR analogues **5a–c**, even if the structures were simplified both in the southern ribose and nucleobase regions. For clearly understanding the structure-activity relationship, further detailed chemistry and/pharmacology investigations are needed.

Figure 3. Superimposition of minimized conformations of cADPR and compounds **5a** (a), **5b** (b), **5c** (c) and **5a–c** (d).



3. Experimental

3.1. Chemistry

3.1.1. General

HR-ESI-MS and ESI-MS were performed with a Bruker BIFLEX III instrument. $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ were recorded with a JEOL AL300 or a Bruker AVANCE III 400; CDCl_3 , D_2O were used as solvents. Chemical shifts are reported in parts per million downfield from TMS (^1H and ^{13}C). $^{31}\text{P-NMR}$ spectra (121.5 MHz) were recorded at room temperature by use of a JEOL AL300 spectrometer. Orthophosphoric acid (85%) was used as external standard. $^{19}\text{F-NMR}$ spectra (470 MHz) were recorded on a Varian VXR-500 spectrometer. Chemical shifts of $^{19}\text{F-NMR}$ are reported in ppm with reference to CF_3COOH as external standard. Compounds were purified on an Alltech preparative C18 reversed-phase columns (2.2×25 cm) with a Gilson HPLC using MeCN/TEAB (pH 7.5) buffer system as eluent. Analytical TLC was performed using commercial glass plates coated to a thickness of 0.25 mm with Kieselgel 60 GF254 silica and visualized under UV light. Flash chromatography was performed using Qingdao (230–400 mesh) silica under a slight positive pressure of air. Microwave-assisted reactions were performed using a Biotage unit (400 w, 2.45 GHz). Solvents and reagents for anhydrous reactions were dried prior to use by conventional methods.

3.1.2. General Procedure for the Synthesis of **18a–c**

To a solution of compound **17** (185 mg, 0.44 mmol) and compound **16a–c** (0.44 mmol) in CH_3CN (15 mL) was added CuI (10 mg, 0.05 mmol) and DIPEA (78 μL , 0.5 mmol), and the mixture was stirred at room temperature for 3 h. The mixture was evaporated *in vacuo* and the residue was partitioned between EtOAc and H_2O . The aqueous phase was extracted again with EtOAc , then the organic layers were combined and dried (Na_2SO_4), filtered and concentrated *in vacuo*. The residue was purified by silica gel column chromatography ($\text{DCM-MeOH} = 40:1-30:1$) to afford a yellow oil.

N-(2-(2-*O*-Bis(phenylthio)phosphorylethoxy)ethyl)-1*H*-1,2,3-triazole-4-carboxamid-1-yl-2'-deoxy-3'-*tert*-butyldimethylsilyl-1'- β -*D*-ribofuranoside (**18a**): Yield: 84%. ESI-TOF⁺: 695.2 [(M+H)⁺]. $^1\text{H-NMR}$ (400 MHz, CDCl_3): δ 8.50 (s, 1H, H-5), 7.54–7.37 (m, 10H, Ar-H), 7.42 (t, 1H, –NH–) 6.44 (dd, 1H, $J = 8$ Hz, H-1'), 4.49–4.47 (m, 1H, H-5'a), 4.25–4.23 (m, 1H, H-5'b), 4.34–4.30 (m, 2H, H-1''), 3.67–3.64 (m, 2H, H-3', H-4'), 3.62–3.56 (m, 4H, H-2'', H-3''), 2.77–2.73 (m, 1H, H-2'a), 2.45 (dd, 1H, $J_{2'a,2'b} = 16$ Hz H-2'b), 0.82 (s, 9H, 3 \times CH₃), 0.06 (s, 6H, 2 \times –Si(CH₃)–). $^{13}\text{C-NMR}$ (75 MHz, CDCl_3): δ 160.3, 143.2, 135.7, 129.4, 126.2, 90.1, 89.5, 76.7, 70.1, 66.8, 62.0, 42.5, 39.1, 25.6, 17.9, –5.0. $^{31}\text{P-NMR}$ (D_2O , decoupled with ^1H): δ 50.91 (s).

N-(2-(2-*O*-Bis(phenylthio)phosphorylethoxy)ethyl)-1*H*-1,2,3-triazole-4-carboxamid-1-yl-2'-chloro-3'-*tert*-butyldimethylsilyl-1'- β -*D*-arabinoside (**18b**): Yield: 85%. ESI-TOF⁺: 729.2 [(M+H)⁺]. $^1\text{H-NMR}$ (400 MHz, CDCl_3): δ 8.34 (s, 1H, H-5), 7.54–7.24 (m, 10H, Ar-H), 6.33 (dd, 1H, $J = 8$ Hz, H-1'), 4.73 (dd, 1H, $J_{5'a,5'b} = 12$ Hz, H-5'a), 4.50 (dd, 1H, $J_{5'a,5'b} = 12$ Hz, H-5'b), 4.36–4.30 (m, 2H, H-4', H-3''), 4.41–3.90 (m, 2H, H-1''), 3.85–3.80 (m, 1H, H-2'), 3.68–3.61 (m, 2H, H-2''), 3.59–3.58 (m, 2H, H-4'', H-3''), 0.89 (s, 9H, 3 \times CH₃), 0.13 (s, 6H, 2 \times –Si(CH₃)–). $^{13}\text{C-NMR}$ (75 MHz, CDCl_3): δ 160.2, 142.5,

135.3, 129.5, 126.3, 88.5, 77.3, 69.7, 66.8, 60.6, 38.6, 25.0, 17.8, −4.5. ^{31}P -NMR (D_2O , decoupled with ^1H): δ 51.27 (s).

N-(2-(2-*O*-Bis(phenylthio)phosphorylethoxy)ethyl)-1*H*-1,2,3-triazole-4-carboxamid-1-yl-2'-fluoro-3'-tert-butyltrimethylsilyl-1'- β -*D*-arabinoside (**18c**): Yield: 83%. ESI-TOF $^+$: 713.2 [(M+H) $^+$]. ^1H -NMR (400 MHz, CDCl_3): δ 8.40 (s, 1H, H-5), 7.50–7.30 (m, 10H, Ar-H), 6.44 (dd, 1H, $J = 8$ Hz, H-1'), 5.08–4.93 (m, 1H, H-5'a) 4.65–4.59 (m, 1H, H-5'b), 4.30–4.28 (m, 2H, H-2', H-4'), 4.01–4.00 (m, 1H, H-3'), 3.78–3.54 (m, 8H, H-1'', H-4'', H-2'', H-3''), 0.86 (s, 9H, 3 \times CH $_3$), 0.08 (s, 6H, 2 \times -Si(CH $_3$) $_2$). ^{13}C -NMR (75 MHz, CDCl_3): δ 160.0, 143.1, 135.3, 129.4, 126.3, 96.2, 66.6, 60.7, 53.5, 38.7, 25.5, 17.8, 5.2. ^{31}P -NMR (D_2O , decoupled with ^1H): δ 51.09 (s).

3.1.3. General Procedure for the Synthesis of **20a–c**

Compound **18a–c** (0.10 mmol) was dissolved in anhydrous CH_3CN (5 mL). DIPEA (0.42 mmol) and POCl_3 (0.35 mmol) were added successively to the solution at -20 °C. The mixture was stirred at 0 °C for 16 h, and then TEAB (5 mL, 1 M, pH 7.5) were added at 0 °C and the stirring was continued for 1 h at room temperature. After evaporation under reduced pressure, the residue was partitioned between H_2O and CHCl_3 , and the aqueous layer was washed with CHCl_3 and evaporated *in vacuo*. The residue was dissolved in TEAB buffer (5 mL, 0.05 M, pH 7.5), and applied to a C18 reversed-phase column (2.2 \times 25 cm). The column was eluted using a linear gradient of 0–80% CH_3CN in TEAB buffer (0.05 M, pH 7.5) to give **19a–c** as its triethylammonium salt. The residue was dissolved in 1 M TEAB (5 mL) and stirred at room temperature for 12 h, and then evaporated *in vacuo*. The residue was partitioned between H_2O and CHCl_3 , and the aqueous layer was washed with CHCl_3 . Evaporated *in vacuo*, dissolved in TEAB buffer (1 mL, 0.05 M, pH 7.5), then applied to a C18 reversed-phase column (2.2 cm \times 25 cm) eluted by a linear gradient of 0–80% CH_3CN in TEAB buffer (0.05 M, pH 7.5) to give **20a–c** as triethylammonium salts.

N-(2-(2-*O*-Phenylthiophosphorylethoxy)ethyl)-1*H*-1,2,3-triazole-4-carboxamid-1-yl-2'-deoxy-3'-tert-butyltrimethylsilyl-5'-phosphoryl-1'- β -*D*-ribofuranoside (**20a**): Yield: 54%. HRMS (ESI-TOF $^-$) Calcd for $\text{C}_{24}\text{H}_{40}\text{N}_4\text{O}_{11}\text{P}_2\text{SSi}$: [(M-H) $^+$] 681.1586; Found: 681.1557. ^1H -NMR (400 MHz, D_2O): δ 8.47 (s, 1H, H-5), 7.47–7.17 (m, 5H, Ar-H) 6.42 (d, 1H, $J = 8$ Hz, H-1'), 4.48 (t, 1H, $J_{5'a,5'b} = 4$ Hz, H-5'a), 4.41 (t, 1H, $J_{5'a,5'b} = 4$ Hz, H-5'b), 3.98–3.96 (m, 2H, H-4', H-3'), 3.57–3.44 (m, 4H, H-2'', H-3''), 3.43–3.40 (m, 2H, H-4''), 3.07 (q, -NCH $_2$ -), 2.70–2.68 (m, 1H, H-2'a), 2.34 (d, 1H, H-2'b), 1.02 (t, -CH $_3$ -), 0.58 (s, 9H, 3 \times CH $_3$), 0.07 (s, 6H, 2 \times -Si(CH $_3$) $_2$). ^{31}P -NMR (D_2O , decoupled with ^1H): δ 15.01 (br, s), 1.70 (br, s).

N-(2-(2-*O*-Phenylthiophosphorylethoxy)ethyl)-1*H*-1,2,3-triazole-4-carboxamid-1-yl-2'-chloro-3'-tert-butyltrimethylsilyl-5'-phosphoryl-1'- β -*D*-arabinoside (**20b**): Yield: 52%. HRMS (ESI-TOF $^-$) Calcd for $\text{C}_{24}\text{H}_{39}\text{ClN}_4\text{O}_{11}\text{P}_2\text{SSi}$: [(M-H) $^+$] 715.1196; Found: 715.1186. ^1H -NMR (400 MHz, D_2O): δ 8.55 (s, 1H, H-5), 7.36–7.06 (m, 5H, Ar-H), 6.47–6.45 (m, 1H, H-1'), 4.48–4.46 (m, 2H, H-5'a, H-5'b), 4.59–4.03 (m, 2H, H-4', H-3'), 4.41–3.90 (m, 2H, H-1''), 3.96–3.95 (m, 1H, H-2'), 3.53–3.49 (m, 4H, H-2'', H-3''), 3.39–3.37 (m, 2H, H-4''), 3.02 (q, -NCH $_2$ -), 1.02 (t, -CH $_3$ -), 0.67 (s, 9H, 3 \times CH $_3$), 0.1 (s, 6H). ^{31}P -NMR (D_2O , decoupled with ^1H): δ 18.24 (br, s), 1.16 (br, s).

N-(2-(2-*O*-Phenylthiophosphorylethoxy)ethyl)-1*H*-1,2,3-triazole-4-carboxamid-1-yl-2'-fluoro-3'-tert-butyl dimethylsilyl-5'-phosphoryl-1'-β-*D*-arabinoside (**20c**): Yield: 50%. HRMS (ESI-TOF[−]) Calcd for C₂₄H₃₉FN₄O₁₁P₂SSi: [(M−H)⁺] 699.1492; Found: 699.1511. ¹H-NMR (400 MHz, D₂O): δ 8.46 (s, 1H, H-5), 7.36–7.09 (m, 5H, Ar-H), 6.50–6.46 (m, 1H, H-1'), 5.30–5.08 (m, 1H, H-4'), 4.11–4.10 (m, 1H, H-5'a), 3.98–3.96 (m, 1H, H-5'b), 3.94–3.91 (m, 3H, H-2', H-1''), 3.57–3.52 (m, 4H, H-2'', H-3''), 3.42–3.39 (m, 2H, H-4''), 3.02 (q, −NCH₂−), 1.09 (t, −CH₃−), 0.74 (s, 9H, 3×CH₃), 0.01 (s, 6H, 2×−Si(CH₃)−). ³¹P-NMR (D₂O, decoupled with ¹H): δ 18.28 (br, s), 1.30 (br, s).

3.1.4. General Procedure for the Synthesis of **21a–c**

The mixture of compound **20a–c** (12.4 mmol) and iodine (73 mg, 0.29 mmol) in pyridine (8 mL) was stirred at 75 °C, assisted by microwave irradiation, for 15 min. Then the pyridine was evaporated, and the residue was partitioned between CHCl₃ and H₂O. The aqueous layer was evaporated and the residue was dissolved in 0.05 M TEAB buffer (5.0 mL), and applied to a C18 reversed-phase column (2.2 × 25 cm). The column was eluted using a linear gradient of 0–80% CH₃CN in TEAB buffer (0.05 M, pH 7.5) to give **21a–c** as triethylammonium salts.

N-(2-(2-*O*-Phosphorylethoxy)ethyl)-1*H*-1,2,3-triazole-4-carboxamid-1-yl-2'-deoxy-3'-tert-butyl dimethylsilyl-5'-phosphoryl-1'-β-*D*-ribofuranoside 2,5'-cyclicpyrophosphate (**21a**): Yield: 87%. HRMS (ESI-TOF[−]) Calcd for C₁₈H₃₄N₄O₁₁P₂Si: [(M−H)⁺] 571.1395; Found: 571.1380. ¹H-NMR (400 MHz, D₂O): δ 8.77 (s, 1H, H-5), 6.39–6.36 (m, 1H, H-1'), 4.47–4.46 (m, 1H, H-5'a), 4.42–4.41 (m, 1H, H-5'b), 4.35–4.34 (m, 2H, H-4', H-3'), 3.96–3.95 (m, 4H, H-2'', H-3''), 3.64–3.59 (m, 2H, H-4''), 3.04 (q, −NCH₂−), 2.57–2.52 (m, 1H, H-2'a), 2.32–2.27 (m, 1H, H-2'b), 1.12 (t, −CH₃−), 0.52 (s, 9H, 3×CH₃), 0.12 (s, 6H). ³¹P-NMR (D₂O, decoupled with ¹H): δ −11.69 (br, s), −12.01 (br, s).

N-(2-(2-*O*-Phosphorylethoxy)ethyl)-1*H*-1,2,3-triazole-4-carboxamid-1-yl-2'-chloro-3'-tert-butyl dimethylsilyl-5'-phosphoryl-1'-β-*D*-arabinoside 2,5'-cyclicpyrophosphate (**21b**): Yield: 82%. HRMS (ESI-TOF[−]) Calcd for C₁₈H₃₃ClN₄O₁₁P₂Si: [(M−H)⁺] 605.1006; Found: 605.1002. ¹H-NMR (400 MHz, D₂O): δ 9.13 (s, 1H, H-5), 6.60 (d, 1H, *J* = 8 Hz, H-1'), 4.76 (d, 1H, *J*_{5'a,5'b} = 8 Hz, H-5'a), 4.55–4.53 (d, 1H, *J*_{5'a,5'b} = 8 Hz, H-5'b), 4.26–4.21 (m, 1H, H-4'), 4.14–4.13 (2H, m, H-1''), 4.05–4.01 (1H, m, H-3'), 3.91–3.88 (1H, m, H-2'), 3.72–3.64 (m, 4H, H-2'', H-3''), 3.58–3.35 (m, 2H, H-4''), 3.02 (q, −NCH₂−), 1.16 (t, −CH₃−), 0.79 (s, 9H, 3×CH₃), 0.09 (s, 6H, 2×−Si(CH₃)−). ³¹P-NMR (D₂O, decoupled with ¹H): δ −12.93 (br, s), −14.43 (br, s).

N-(2-(2-*O*-Phosphorylethoxy)ethyl)-1*H*-1,2,3-triazole-4-carboxamid-1-yl-2'-fluoro-3'-tert-butyl dimethylsilyl-5'-phosphoryl-1'-β-*D*-arabinoside 2,5'-cyclicpyrophosphate (**21c**): Yield: 81%. HRMS (ESI-TOF[−]) Calcd for C₁₈H₃₃FN₄O₁₁P₂Si: [(M−H)⁺] 589.1302; Found: 589.1325. ¹H-NMR (400 MHz, D₂O): δ 9.09 (s, 1H, H-5), 6.63 (dd, 1H, *J* = 8 Hz, H-1'), 5.36–5.21 (m, 1H, H-4'), 4.20–4.16 (m, 1H, H-5'a), 4.16–4.14 (m, 1H, H-5'b), 4.10–3.91 (m, 3H, H-2', H-1''), 3.68–3.52 (m, 4H, H-2'', H-3''), 3.54–3.34 (m, 2H, H-4''), 3.01 (q, −NCH₂−), 1.13 (t, −CH₃−), 0.79 (s, 9H, 3×CH₃), 0.09 (s, 6H, 2×−Si(CH₃)−). ³¹P-NMR (D₂O, decoupled with ¹H): δ −12.90 (br, s), −14.22 (br, s).

3.1.5. General Procedure for the Synthesis of **5a–c**

Compound **21a–c** (33.6 μmol) was dissolved in 1M TBAF/THF (0.5 mL) and the solution was stirred for 2 h, and then was evaporated under reduced pressure. The residue was dissolved in 0.05 M TEAB buffer (2.0 mL), which was loaded to C18 reversed-phase column (2.2×25 cm). The column was eluted using a linear gradient of 0–80% CH_3CN in TEAB buffer (0.05 M, pH 7.5) to afford **5a–c** as triethylammonium salts.

N-(2-(2-*O*-Phosphorylethoxy)ethyl)-1*H*-1,2,3-triazole-4-carboxamid-1-yl-2'-deoxy-5'-phosphoryl-1'- β -*D*-ribofuranoside 2,5'-cyclicpyrophosphate (**5a**): Yield: 36%. HRMS (ESI-TOF⁻) Calcd for $\text{C}_{12}\text{H}_{20}\text{N}_4\text{O}_{11}\text{P}_2$: $[(\text{M}+\text{H})^+]$ 459.1051; Found: 459.1040. ¹H-NMR (400 MHz, D_2O): δ 8.47 (s, 1H, H-5), 6.48–6.42 (m, 1H, H-1'), 4.48–4.47 (m, 1H, H-5a), 4.37–4.30 (m, 1H, H-5'b), 3.99–3.97 (m, 2H, H-1''), 3.87–3.86 (m, 1H, H-3'), 3.61–3.55 (m, 4H, H-3'', H-4''), 3.46–3.43 (m, 2H, H-2''), 3.09 (q, $-\text{NCH}_2-$), 2.89–2.77 (m, 1H, H-2'a), 2.43–2.40 (m, 1H, H-2'b), 1.17 (t, $-\text{CH}_3-$). ³¹P-NMR (D_2O , decoupled with ¹H), δ -9.77 (br, s), -10.05 (br, s).

N-(2-(2-*O*-Phosphorylethoxy)ethyl)-1*H*-1,2,3-triazole-4-carboxamid-1-yl-2'-chloro-5'-phosphoryl-1'- β -*D*-arabinoside 2,5'-cyclicpyrophosphate (**5b**): Yield: 31%. HRMS (ESI-TOF⁻) Calcd for $\text{C}_{12}\text{H}_{19}\text{ClN}_4\text{O}_{11}\text{P}_2$: $[(\text{M}-\text{H})^+]$ 491.0141; Found: 491.0142. ¹H-NMR (400 MHz, D_2O): δ 9.17 (s, 1H, H-5), 6.60 (d, 1H, $J = 8$ Hz, H-1'), 4.77–4.73 (m, 2H, H-5'a, H-5'b), 4.26–4.21 (m, 1H, H-4'), 4.24–4.09 (2H, m, H-1''), 4.00–3.98 (1H, m, H-3'), 3.69–3.66 (1H, m, H-2'), 3.50–3.40 (m, 4H, H-2'', H-3''), 3.10–3.04 (m, 2H, H-4''). 3.04 (q, $-\text{NCH}_2-$), 1.15 (t, $-\text{CH}_3-$). ³¹P-NMR (D_2O , decoupled with ¹H): δ -7.58 (br, s), -9.00 (br, s).

N-(2-(2-*O*-Phosphorylethoxy)ethyl)-1*H*-1,2,3-triazole-4-carboxamid-1-yl-2'-fluoro-5'-phosphoryl-1'- β -*D*-arabinoside 2,5'-cyclicpyrophosphate (**5c**): Yield: 29%. HRMS (ESI-TOF⁻) Calcd for $\text{C}_{12}\text{H}_{19}\text{FN}_4\text{O}_{11}\text{P}_2$: $[(\text{M}-\text{H})^+]$ 475.0437; Found: 475.0449. ¹H-NMR (400 MHz, D_2O): δ 9.14 (s, 1H, H-5), 6.64 (m, 1H, H-1'), 5.46–5.43 (m, 1H, H-5'a), 5.33–5.32 (m, 1H, H-5'b), 4.20–4.15 (m, 2H, H-1''), 4.02–3.99 (m, 2H, H-2', H-4'), 3.69–3.67 (m, 2H, H-2''), 3.52–3.41 (m, 4H, H-3'', H-4''), 3.06 (q, $-\text{NCH}_2-$), 2.96–2.94 (m, 1H, H-3'), 1.13 (t, $-\text{CH}_3-$). ³¹P-NMR (D_2O , decoupled with ¹H): δ -9.57 (br, s), -10.53 (br, s). ¹⁹F-NMR (D_2O , CF_3COOH), δ -202.13–202.32 (m).

3.2. Biological Studies

Cell Culture. The human Jurkat T-lymphocyte cell line was obtained from the Hongkong University Physiology Department Professor Lee Hon Cheung's lab Culture Collection. It was cultured in RPMI Medium 1640 (Invitrogen, Carlsbad, CA, USA) supplemented with 10% fetal bovine serum (FBS), 100 units/mL penicillin, and 2% Hepes Buffer (1 M, pH 7.4) in a humidified (5% CO_2) atmosphere at 37 °C.

Calcium agonistic activity measurement. The cells (3×10^5 cells/well) were plated in 24-well plate with poly L-lysine (100 $\mu\text{g}/\text{mL}$) and incubated in no FBS and penicillin medium at 37 °C overnight for adherence. Before measurement, cells were incubated with Fura-2 AM (2 μM) in HBSS for 20 min in dark at 37 °C. Then cells were washed with calcium measurement buffer HBSS twice and added

200 μL HBSS for measurement. Changes in Fura-2 fluorescence were measured using an Olympus Cell^R Live-cell confocal imaging system, operating in ratio mode (alternating excitation at 340 and 380 nm). Each compound was dissolved at tested concentrations in 50 μL HBSS and added into the cells at a time point 40 s after the start of each measurement. Data (F: ratio between emission at 340 and 380 nm) were recorded for the next 500 s. Changes in mean F/F_0 (F_0 : Ratio between emission at 340 and 380 nm at the start point of measurement) after the application of compounds were averaged for at least three independent experiments for each concentration. The peak values of F/F_0 were analyzed as the maximum within 200 s after compound application, shown in the ordinate of Figure 2. Response to the application of cIDPRE was served as a positive control while addition of buffer was used as blank.

3.3. Calculations

The conformations of studied cADPR analogues were optimized with density functional theory (DFT) quantum chemical method by using Gaussian 09 program package [33]. Equilibrium geometries of all molecules were fully optimized at the B3LYP/6–31G(d) level of theory [34]. Vibrational frequencies, calculated at the same levels, were used to determine the nature of the stationary points and to give the thermodynamic data and zero-point vibrational energies. Then SYBYL-X 1.1 software was used to align the calculated structures of three cADPR analogs with crystal structure of cADPR, respectively.

4. Conclusions

In summary, novel nucleobase-modified and sugar-modified cADPR mimics **5** were synthesized, in which halogen atoms were introduced into the southern ribose for the first time. The synthesis employed a Cu(I)-catalyzed Huisgen [3+2] cycloaddition for the building of the triazole moiety and microwave-assisted reactions were used for the pyrophosphate bond formation. Biological evaluation reveals that these new kinds of cADPR analogues are membrane permeant agonists of the calcium signaling pathway. The introduction of chlorine or fluorine into the 2'-position of the southern ribose led to a decrease of activity. The existence of a hydrophobic group on the 3'-OH of the southern ribose does not alter much the agonistic activity. The result of this study is helpful for the understanding of structure-activity relationships of cADPR analogues and to help design new probes to investigate the cADPR-mediated calcium signaling pathway.

Acknowledgements

This study was supported by the National Natural Sciences Foundation of China (Grant No. 90713005, 20910094) and the Ministry of Education of China (Grant No. 200800010078).

References and Notes

1. Guse, A.H. Regulation of calcium signaling by the second messenger cyclic adenosine diphosphoribose (cADPR). *Curr. Mol. Med.* **2004**, *4*, 239–248.
2. Lee, H.C. *Cyclic ADP-Ribose and NAADP: Structures, Metabolism and Functions*; Kluwer Academic Publisher: Dordrecht, The Netherland, 2002; pp. 217–444.

3. Lee, H.C. Multiplicity of Ca^{2+} messengers and Ca^{2+} stores: A perspective from cyclic ADP-ribose and NAADP. *Curr. Mol. Med.* **2004**, *4*, 227–237.
4. Shuto, S.; Matsuda, A. Chemistry of cyclic ADP-ribose and its analogs. *Curr. Med. Chem.* **2004**, *11*, 827–845.
5. Guse, A.H. Biochemistry, biology, and pharmacology of cyclic adenosine diphosphoribose (cADPR). *Curr. Med. Chem.* **2004**, *11*, 847–855.
6. Potter, B.V.L.; Walseth, T.F. Medicinal chemistry and pharmacology of cyclic ADP-ribose. *Curr. Mol. Med.* **2004**, *4*, 303–311.
7. Kudoh, T.; Fukuoka, M.; Ichikawa, S.; Murayama, T.; Ogawa, Y.; Hashii, M.; Higashida, H.; Kunerth, S.; Weber, K.; Guse, A.H.; *et al.* Synthesis of stable and cell-type selective analogues of cyclic ADP-ribose, a Ca^{2+} -mobilizing second messenger. Structure-activity relationship of the N¹-ribose moiety. *J. Am. Chem. Soc.* **2005**, *127*, 8846–8855.
8. Shuto, S.; Fukuoka, M.; Manikowsky, A.; Ueno, Y.; Nakano, T.; Kuroda, R.; Kuroda, H.; Matsuda, A. A total synthesis of cyclic ADP-carbocyclic-ribose, a stable mimic of Ca^{2+} -mobilizing second messenger cyclic ADP-ribose. *J. Am. Chem. Soc.* **2001**, *123*, 8750–8759.
9. Xu, L.; Walseth, T.; Slama, J. Cyclic ADP-ribose analogues containing the methylenebisphosphonate linkage: Effect of pyrophosphate modifications on Ca^{2+} release activity. *J. Med. Chem.* **2005**, *48*, 4177–4181.
10. Aarhus, R.; Gee, K.; Lee, H.C. Caged cyclic ADP-ribose. Synthesis and use. *J. Biol. Chem.* **1995**, *270*, 7745–7749.
11. Zhang, F.; Yamada, S.; Gu, Q.M.; Jing, P.; Sih, C.J. Synthesis and characterization of cyclic ATP-ribose: A potent mediator of calcium release. *Bioorg. Med. Chem. Lett.* **1996**, *6*, 1203–1208.
12. Qi, N.; Jung, K.; Wang, M.; Na, L.X.; Yang, Z.J.; Zhang, L.R.; Guse, A.H.; Zhang, L.H. A novel membrane-permeant cADPR antagonist modified in the pyrophosphate bridge. *Chem. Commun.* **2011**, *47*, 9462–9464.
13. Moreau, C.; Wagner, G.K.; Weber, K.; Guse, A.H.; Potter, B.V.L. Structural determinants for N1/N7 cyclization of nicotinamide hypoxanthine 5'-dinucleotide (NHD⁺) derivatives by ADP-ribosyl cyclase from *Aplysia californica*: Ca^{2+} -Mobilising activity of 8-substituted cyclic inosine 5-diphosphoribose analogues in T-lymphocytes. *J. Med. Chem.* **2006**, *49*, 5162–5176.
14. Graeff, R.M.; Walseth, T.F.; Hill, T.K.; Lee, H.C. Fluorescent analogs of cyclic ADP-ribose: Synthesis, spectral characterization and use. *Biochemistry* **1996**, *35*, 379–386.
15. Bailey, V.C.; Sethi, J.K.; Fortl, S.M.; Galione, A.; Potter, B.V.L. 7-Deaza cyclic adenosine 5'-diphosphate ribose first example of a Ca^{2+} -mobilizing partial agonist related to cyclic adenosine 5'-diphosphate ribose. *Chem. Biol.* **1997**, *4*, 51–56.
16. Huang, X.C.; Dong, M.; Liu, J.; Zhang, K.H.; Yang, Z.J.; Zhang, L.R.; Zhang, L.H. Concise syntheses of trifluoromethylated cyclic and acyclic analogues of cADPR. *Molecules* **2010**, *15*, 8689–8701.
17. Gu, X.F.; Yang, Z.J.; Zhang, L.R.; Kunerth, S.; Fliegert, R.; Weber, K.; Guse, A.H.; Zhang, L.H. Synthesis and biological evaluation of novel membrane-permeant cyclic ADP-ribose mimics: N¹-[(5"-O-phosphoryl-ethoxy)-methyl]-5'-O-phosphoryl-inosine 5',5"-cyclicpyrophosphate (cIDPRE) and 8-substituted derivatives. *J. Med. Chem.* **2004**, *47*, 5674–5682.

18. Xu, J.F.; Yang, Z.J.; Dammermann, W.; Zhang, L.R.; Guse, A.H.; Zhang, L.H. Synthesis and agonist activity of cyclic ADP-ribose analogues with substitution of the northern ribose by ether or alkane chains. *J. Med. Chem.* **2006**, *49*, 5001–5012.
19. Guse, A.H.; Gu, X.F.; Zhang, L.R.; Weber, K.; Zhang, L.H. A minimal structural analogue of cyclic ADP-ribose. *J. Biol. Chem.* **2005**, *280*, 15952–15959.
20. Huang, L.J.; Zhao, Y.Y.; Yuan, L.; Min, J.M.; Zhang, L.H. Syntheses and calcium mobilizing evaluations of N^1 -glycosyl substituted stable mimics of cyclic ADP-ribose. *J. Med. Chem.* **2002**, *45*, 5340–5352.
21. Kudoh, T.; Fukuoka, M.; Shuto, S.; Matsuda, A. Synthesis and biological activity of cyclic ADP-carbocyclic-ribose analogues: Structure-activity relationship and conformational analysis of N -1-carbocyclic-ribose moiety. *Nucleos. Nucleot. Nucleic Acids* **2005**, *24*, 655–658.
22. Kudoh, T.; Fukuoka, M.; Ichikawa, S.; Murayama, T.; Ogawa, Y.; Hashii, M.; Higashida, H.; Kunerth, S.; Weber, K.; Guse, A.H.; *et al.* Synthesis of stable and cell-type selective analogues of cyclic ADP-ribose, a Ca^{2+} -mobilizing second messenger. Structure-activity relationship of the $N1$ -ribose moiety. *J. Am. Chem. Soc.* **2005**, *127*, 8846–8855.
23. Zhang, B.; Wagner, G.K.; Weber, K.; Garnham, C.; Morgan, A.; Galione, A.; Guse, A.H.; Potter, B.V.L. 2'-Deoxy cyclic adenosine 5'-diphosphate ribose derivatives: Importance of the 2'-hydroxyl motif for the antagonistic activity of 8-substituted cADPR derivatives. *J. Med. Chem.* **2008**, *51*, 1623–1636.
24. Moreau, C.; Ashamu, G.A.; Bailey, V.C.; Galione, A.; Gusec, A.H.; Potter, B.V.L. Synthesis of cyclic adenosine 5'-diphosphate ribose analogues: A $C2'$ *endo/syn* “southern” ribose conformation underlies activity at the sea urchin cADPR receptor. *Org. Biomol. Chem.* **2011**, *9*, 278–290.
25. Li, L.J.; Lin, B.C.; Yang, Z.J.; Zhang, L.R.; Zhang, L.H. A concise route for the preparation of nucleobase-simplified cADPR mimics by click chemistry. *Tetrahedron Lett.* **2008**, *49*, 4491–4493.
26. Kazuhiro, H.M.G. Synthesis and characterization of oligonucleotides containing formadidopyrimidine lesions and nonhydrolyzable analogues. *J. Am. Chem. Soc.* **2001**, *123*, 8638–8637.
27. Du, J.F.; Choi, Y.; Chu, C.K. A practical synthesis of L-FMAU from L-arabinose. *Nucleos. Nucleot.* **1999**, *18*, 187–195.
28. Bruce, G.; Anderson, D.P.L. Isolation, synthesis, and characterization of impurities and degradants from the clofarabine process. *Org. Proc. Res. Dev.* **2008**, *12*, 1229–1237.
29. Li, L.J.; Guse, A.H.; Zhang, L.H. Novel nucleobase-simplified cyclic ADP-ribose analogue: A concise synthesis and Ca^{2+} -mobilizing activity in T-lymphocytes. *Org. Biol. Chem.* **2010**, *8*, 1843–1848.
30. Polshettiwar, V.; Varma, R.S. Aqueous microwave chemistry: A clean and green synthetic tool for rapid drug discovery. *Chem. Soc. Rev.* **2008**, *37*, 1546–1557.
31. Kappe, C.O. Microwave dielectric heating in synthetic organic chemistry. *Chem. Soc. Rev.* **2008**, *37*, 1127–1139.
32. Guse, A.H. Cyclic ADP-ribose. *J. Mol. Med.* **2000**, *78*, 26–35.
33. Frisch, M.J.; Trucks, G.W.; Schlegel, H.B.; Scuseria, G.E.; Robb, M.A.; Cheeseman, J.R.; Montgomery, J.A., Jr.; Vreven, T.; Kudin, K.N.; Burant, J.C.; *et al.* *Gaussian 03*, Revision B.01; Gaussian Inc.: Pittsburgh, PA, USA, 2010.

34. Becke, A.D. Density-functional thermochemistry. III. The role of exact exchange. *J. Chem. Phys.* **1993**, *93*, 5648.

Sample Availability: Samples of the compounds **5a–c** are available from the authors.

© 2012 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).