



US 20030224424A1

(19) **United States**

(12) **Patent Application Publication**

(10) **Pub. No.: US 2003/0224424 A1**

Garcia-Barcelo et al.

(43) **Pub. Date:**

Dec. 4, 2003

(54) **PHOX2B POLYMORPHISMS AS
HIRSCHSPRUNG'S DISEASE DIAGNOSTIC
MARKERS AND METHODS BASED
THEREON**

(52) **U.S. Cl.** **435/6; 435/69.1; 435/189;
435/320.1; 435/325; 536/23.2**

(76) **Inventors: Maria Mercedes Garcia-Barcelo,
Hong Kong (HK); Mai Har Sham,
Hong Kong (HK); Paul Kwong Hang
Tam, Aberdeen (HK); Vincent Chi
Hang Lui, Hong Kong (HK); Benedict
Ling Sze Chen, New Territories (HK)**

(57) **ABSTRACT**

Correspondence Address:
**PENNIE AND EDMONDS
1155 AVENUE OF THE AMERICAS
NEW YORK, NY 100362711**

The invention relates generally to polymorphisms or mutations of the PHOX2B gene. More particularly, the invention relates to polymorphisms or mutations of the PHOX2B gene that are responsible for the disease Hirschsprung's disease (HSCR), which is a neural crest-associated developmental disorder. Specifically, the invention relates to the detection of a single base-pair polymorphism in the PHOX2B gene that is associated with HSCR. The invention also relates to methods and kits for screening for carriers of mutations of the PHOX2B gene and the diagnosis of increased risk of HSCR. The invention further relates to diagnosing predisposition or susceptibility to increased risk of developing HSCR by screening for the presence of a polymorphism associated with HSCR. The invention also relates to compositions for screening for the polymorphism and treatment choices for patients having the polymorphism of the present invention. The invention further relates to providing polymorphisms in the PHOX2B gene for forensic use and in paternity test. The invention also relates to screening assays and therapeutic and prophylactic methods.

(21) **Appl. No.: 10/408,501**

(22) **Filed: Apr. 7, 2003**

Related U.S. Application Data

(60) **Provisional application No. 60/370,968, filed on Apr. 8, 2002.**

Publication Classification

(51) **Int. Cl.⁷ C12Q 1/68; C07H 21/04;
C12N 9/02; C12P 21/02; C12N 5/06**

Figure 1A

1 cctgagtctc ttgaaagcac atttagctgc aagaagaaat gcaatttata agatgctttc
61 tctctgtctc tctttgtttt taaccccctc taaccagggtg tattttttaa gaaatccgct
121 tategttcac ataaaccccc ttggcccact tactctatgt tacagggcgc ctgagctctg
181 ccaatgtccc agtcctttat aacatttcat gcacttcggg gggtaggctt gttgttaa
241 tgagcgtgta acactcttac aaaacagggtt ttctatgaca tcaaggtttc ttctccctaa
301 ccgaggggga aaaaaagaag aggacgaaga agggggaaaa cacacacact atctcaattt
361 atgcctaagg tatatgatca gttaaaaagg cttaaaagct cggggaaatt ggatcagggg
421 gaatcgteac ccaactttca ttatttccaa gtagtgtgat tgaattaaag ggcagggagc
481 tggttagaag ggaggatcag gggctcgggtg cgtaatgggtg tggatttaa ttctaattag
541 agatgcagga atcaatgata gggagggttg acagctcagt tccccagtgc cagcccaata
601 gacggatgag ttattgtcat gtaaaaagcg ccagcaataa gaccaaccgc tttgctattg
661 tccaagtgga aagagccaag tttattatga ggactatag ctctagagac ctgagacaag
721 gcactcaca ggaggctttt tcataaaact aggctctgct ggtagtaagg aggccagttt
781 ggaggcaggc gttgagctgt geacatctcc ccactccagc caccttctcc atatccatc
841 tttatttcat ttttccactt ggctgagcca tccagaacct tttcaatgta taaaatggaa
901 tattcttacc tcaattctc tgccacgag tctgtatgg ctgggatgga cacctcgagc
961 ctggcttcag cctatgctga cttcagttcc tgcagccagg ccagtggctt ccagtataac
1021 ccgataagga ccacttttgg ggccacgtcc ggctgccctt cctcaogcc gggatcctgc
1081 agcctgggca ccctcagga ccaccagagc agtccgtacg ccgcaggtta ggaccttcag
1141 ctttctcagc ggaggaagcc gcctttccgc ccgtatatag gaagccttga ttgcatttga
1201 aatggaaat gtgttttagta tttaccaaac gaaatttget tacacaaatg aaagaattta
1261 tcacgtttga agcgattgca gggaggggta attcacttac agggttacac tatcctagtc
1321 acaccogaac cgcccacaaa attatcttaa gctgccaaaa tgataggeat aatttattta
1381 ctttgcgatg agacgtatag cttagaaaat aattgaatta caaagagtaa agctcattac
1441 tggcagtgtc tcttttttta agaaccgaga gcggctcaca cttcttgggc tggtcatttt
1501 tatgattatt tctttaattt attattattt ttttgcagct ctttccccca acttttgagc
1561 cgggtcaact ttctgagaat tgaaaagttc ccaaagtggg actgtttggg aacttctttc
1621 cctggctccc ctgatattcc gactgatgtt ttgggatttt tttcctctct ggttttttcc
1681 tgctgaaagc actatctcaa gtccgtcaca tcgcgctgtt tcaatccacc caaaggecct
1741 tgtgccagaa aggactccgc caagcccgaa gtttgagccc aggtttccgc agataacaaa

1801 tttcctcggt ttetteccgc agettctctc ggcaactctc tcgcgegggt gtaggtagcg
1861 gctgcegtat gacctgacct tggagtcctc acattctagc tccacggcgc gcgagctgcc
1921 ggetgatattg ctcactttct gtctcctctg tcatactcta gttccttaca aactcttcac
1981 ggaccaeggc ggctcaacg agaagcgcaa gcagcggcgc atccgacca ctttcaccag
2041 tgcccagctc aaagagctgg aaaggtctt cgcggagact cactaccccg acatctacac
2101 tcgggaggag ctggccctga agatcgacct cacagaggcg cgagtccagg tacgcgcgcc
2161 tggaaaccga ccccgctccg ccgcactggt ccggggaggt gtgggggtgag gggcggctgg
2221 tgaaattcga agtcctggag cctcgagtga gaaggaccta gggcccatg gccgatcaga
2281 aatactggat ttgggtgtggc tgtgcgttcg agagaggctt agagcgcacg ctcttggcat
2341 tttatttaca gttgcgaagt gtttcccacc cgagcagaga catggggggc ctggggacgt
2401 ggatgagcga tgcaatttcg gggacaggaa gtgcctgtgg tggaaaggtg gcagactttg
2461 ctcccgtatt ataagttttt ccttctcccc tcccgcccc caaaaaaatg cctcctaact
2521 caagtgtttt taacctggcc ccatggcata taggttcatt ttcccggaaa ctgtgacttg
2581 cattagattht gcaaagggtc tgtgacttca tgaaggtcaa gaacctgac ttactccaac
2641 ctgttaaaca caggtgcgct cacgagttgg ccacagcgcc tttttgggtg agccccgac
2701 cgagaagcgg tgcgcaccat tgcacgcttt tccaggctca aaggccgggg atgggcagcg
2761 gagcaaaccc agagaggatc ccttttctct ttaccaatta gagtttaact ttagaactta
2821 ggcttagggg tgaatggcga gctcggggct tgctcaagaa gccgactgaa cagaggccca
2881 caaaataag gccttccctt ttccgggtct tetgggacct geggettttt aaactctgcc
2941 gcaagccttc atgtccctgg cgtgctcact ccccctaaga aagtttctcc gaaaatgac
3001 agcaataaga agcggtagac ttgggtgatg tgcgcgcggg ggtgatcaca gcgcatgggg
3061 aggagggtgt taaaacaagc cgaagtagaa cttgggcccac cctaaccggg gcttttcttt
3121 ccattttctt tttttctccc cctgcttca cgtctctctt tccgtcttgg gccaggtgtg
3181 gttccagaac cgccgcgcca agtttcgcaa gcaggagcgc gcagcggcag ccgcagcggc
3241 cgcggccaag aacggctcct cgggcaaaaa gtctgactct tccagggacg acgagagcaa
3301 agaggccaag agcactgacc cggacagcac tgggggccc ggtcccacac ccaacccac
3361 cccagctgc ggggcgaatg gaggcggcgg cggcgggccc agcccggctg gagctcggg
3421 ggcggcgggg cccgggggccc cgggaggcga acccggcaag ggcggcgcag cagcagcggc
3481 ggcggcgcgc gcagcggcgg cggcggcagc ggcagcggcg gcagctggag gcctggctgc
3541 ggttgggggc cctggacaag gctgggctcc cggccccggc cccatcact ccatcccga
3601 ttcgcttggg ggtcccttcg gcagcgtcct atcttcgctc caaagacca acggtgccc
3661 agccgcctta gtgaagagca gtatgttctg atctggaatc ctgcggcggc ggcggcggcg
3721 gcgacagcgg gcgagccagg gcccggcgg gcgagtgggc gagcgggtag gcccaaggct
3781 attgtcgtcg ctgctgccat ggctttttca ttgaggccct aaagtaatcg cgctaagaat
3841 aaagggaaaa cggcgtcgc ctcatttcaa ccccactcct acccccttc tcaacccca

3901 aacaaaacaa acaaacttcc ctggcttccg acctgectgg ggcctcgcag cggggccagg
3961 gctccgcttg ctgatcgggg gttgtgagca gcgcggcctg gacgcggggc actctcaggg
4021 ggctgtgtct gcgtgtcagt ttgtgtctgt ctcggggaat gtgtgtctgt ggccaagca
4081 ggtgacagga agagatgggg ggectcaacc aacttagtga cttgtttaga aaaaaagac
4141 aaaaaagtaa aaataaaaac aaaaaagttg gaaggcagaa accattaaaa aacaaaaagc
4201 caacaacca gaaaggttta aaaaacataa ggaaaaaaaa gacaaattaa aggaggggct
4261 aggggagaag ctgcagctgg agctgaaggc tcgatcttgt gaaccctaa atccgctccc
4321 tcctaacagc acggattctc ttggggctct tcttcagggg agagtaggga cgccgtcca
4381 gcecccttc ctatcgtgtc cttgggttcg ggtcaactgc ggcacgactt gctcagactg
4441 tcccggcggc cggagtgact ttctgcacc cccttgectg tcccacctcg ctgaacacca
4501 tcccgccatt agcgcacgg aacccccacac agttgcaact cccaaccccg aatctttgca
4561 gccgttcggc cctgaaagat gccctatcca tgagatgect tttcatctgc aaactctgca
4621 aatgtgtct catgtttcgc aactctttt tccccctcg ctcccgccta ccccgctggc
4681 atttctctt tccaccagct ttaactgaac tttttggcac tgctttggat tggggctaat
4741 tgcagtcac gtaactggct gcagagaaat ctaccgagca aggaaaaggc acacacacac
4801 gtttgcaggg gtgtctcggg ttgcatttct gttggaatga tccgaactgg actcacatcc
4861 tgtatgggtg atggactgta tattgagggt tccattcttc ggcagttta gacatctctg
4921 ttttgattct ttgttgttgt tttatttta aaaggcacia actctagata ttagttgaat
4981 gttgaggctt taacttttcc ggtgtcttcc tacaactgtg ttctgtgact caattgtatc
5041 gtgttaatat cagtgcagac tgtctctctc acgtgaccgt ataatgtttt tctcgtcttg
5101 tagtctctat ggcgtgtctt tatgggtgaa taaggttctc acgggggtcaa tcttttgtgt
5161 ttagagaggg cacggttcag acaatggat atatttttgt tatcagggtc atgtctgtct
5221 gatttctttt ttttctctgt tggactatgt ttgtgaacat aattgtcata agttatgttt
5281 cagatttttg aatttattta tatgtgttat aatgaatgct tctatttaaa agggaaatat
5341 ttctacatgt gcttatagtt ttccaagagt gtaccattaa cttgattgtt gataataaaa
5401 accaaaagca agtctagcaa ttgaactctt cttttctctg attctttttt tttttttttt
5461 tttgggttgg tcattgtttt ttttttttaa gttttttttt aaaaggat

Figure 1B

1 mykneysyln ssayescmag mdtsslasay adfsscsqas gfqynpirtt fgatsgcpsl
61 tpgscslgtl rdhqsspyaa vpyklftdhg glnekrkqrr irttftsaql kelervfaet
121 hypdiytrees lalkidltea rvqvwfqnr akfrkqeraa aaaaaakng ssgkksdssr
181 ddeskeakst dpdstggppp npnptpcga nngggggppsp agapgaagpg qpggepgkge
241 aaaaaaaaaa aaaaaaaaaa gglaaagppg qgwapgppi tsipdslgpp fgsvlsslqr
301 pngakaalvk ssmf

Figure 2

1 cctgagtctc ttgaaagcac atttagctgc aagaagaaat gcaatttata agatgctttc
61 tctctgtctc tctttgtttt taaccccctc taaccagggtg tattttttaa gaaatecget
121 tategttcac ataaaccccc ttggcccact tactctatgt tacagggcgc ctgagtcttg
181 ccaatgtccc agtcctttat aacatttcat gcacttcggg gggtaggctt gttgttaa
241 tgagcgtgta acactcttac aaaacaggtt ttctatgaca tcaaggtttc ttctccctaa
301 ccgaggggga aaaaaagaag aggacgaaga agggggaaaa cacacacact atctcaattt
361 atgectaaag tatatgatca gttaaaaagg cttaaaagct cggggaaatt ggatcaggga
421 gaatcgtcac ccaactttca ttatttccaa gtagtgtgat tgaattaaag ggcaggggagc
481 tggttagaag ggaggatcag gggctcgggtg cgtaatgggtg tggattaaa ttctaattag
541 agatgcagga atcaatgata gggaggttgg acagctcagt tccccagtgc cagcccaata
601 gacggatgag ttattgtcat gtaaaaaagc ccagcaataa gaccaaccgc tttgctattg
661 tccaagtgga aagagccaag tttattatga ggactatatg ctctagagac ctgagacaag
721 gcatctcata ggaggctttt tcataaaact aggctctgct ggtagtaagg aggccagttt
781 ggaggcagge gttgagctgt gcacatctcc ccactccage cacctctcc atatccatct
841 tttatttcat tttccactt ggctgagcca tccagaacct tttcaatgta taaaatggaa
901 tattcttacc tcaattctc tgctacgag tctgtatgg ctgggatgga cacctcgagc
961 ctggcttcag cctatgctga cttcagttcc tgcagccagg ccagtggctt ccagtataac
1021 ccgataagga ccacttttgg ggccacgtcc ggctgccctt cctcaagcc gggatcctgc
1081 agcctgggca cctcagga ccaccagagc agtccgtacg ccgcaggtta ggacctcag
1141 ctttctcage ggaggaagcc gccttccgc ccgtatatag gaagccttga ttgcatttga
1201 aatggaaat gtgtttagta tttacaaac gaaatttgct tacacaaatg aaagaattta
1261 tcacgtttga agcgattgca gggaggggta attcacttac agggttacac tatcctagtc
1321 acaccggaac cgcccacaaa attatcttaa gctgccaaaa tgatagcat aatttattta
1381 ctttgcgatg agacgtatag cttagaaaat aattgaatta caaagagtaa agctcattac
1441 tggcagtgtc tcttttttta agaaccgaga gggctcaca cttcttgggc tggtcatttt
1501 tatgattatt tctttaattt attattattt ttttgcagct ctttccccca acttttgagc

1561 cgggtcaact ttctgagaat tgaaaagttc ccaaagtggg actgtttggt aacttetttc
1621 cctggctccc ctgatattcc gactgatggt ttgggatttt ttctctctct ggttttttcc
1681 tgctgaaagc actatctcaa gtccgtcaca tcgcgctggt tcaatccacc caaaggcgct
1741 tgtgccagaa aggactccgc caagcccgaa gtttgagccc aggtttccgc agataacaaa
1801 ttctctcggg ttcttcccgc agcttctctc ggcaactctc tcgcgcgggt gtaggtagcg
1861 gctgcgctat gacctgacct tggagctctc acattctagc tccacggccg gcgagctgcc
1921 ggctgatttg ctcactttct gtctctctctg tcatacteta gttccttaca aactcttca
1981 ggaccacggc ggctcaacg agaagcgcaa gcagcggcgc atccgcacca ctttaccag
2041 tgcccagctc aaagagctgg aaagggctct cgcggagact cactaccccg acatctacac
2101 tcgggaggag ctggccctga agatcgacct cacagaggcg cgagtccagg tacgcgcgc
2161 tggaaaccga ccccgctccg ccgcactggg cgggggaggt gtggggtagg gggcgctgg
2221 tgaaattcga agtccctggg cctcgagtgc gaaggacct gggccccatg gccgatcaga
2281 aatactggat ttgggtgtgg tgtgcgttcg agagaggctt agagcgcacg ctcttggcat
2341 tttatttaca gttgcgaagt gtttcccacc cgagcagaga catggggggc cttgggaegt
2401 ggatgagcga tgcaatttcg gggacaggaa gtgcctgtgg tggaaaggtg gcagactttg
2461 ctcccgtatt ataagttttt cttctcccc tcccgcctcc caaaaaaatg cctcctaact
2521 caagtgtttt taacctggcc ccatggcata taggttcatt ttcccggaaa ctgtgacttg
2581 cattagattt gcaaagggc tgtgacttca tgaaggtcaa gaacctgac ttactccaac
2641 ctgttaaaca caggtgcgct cagagttgg ccacagcgc tttttgggtg agccccgac
2701 cgagaagcgg tgcgcaccat tgcacgttt tccaggetca aaggccgggg atgggcagcg
2761 gagcaaacc agagaggatc cttttctct ttaccaatta gagttaact ttagaacta
2821 ggcttagggg tgaatggcga gctcggggct tgctcaagaa gccgactgaa cagaggcca
2881 caaaaataag gcttccctt ttccgggtct tctgggacct gggcttttt aaactctgcc
2941 gcaagcctc atgtccctgg cgtgctcact cccctaaga aagtttctcc gaaaatgcac
3001 agcaataaga agcggtagac ttgggtgatg tgcgcgcggg ggtgatcaca gcgcatggg
3061 aggagggtgt taaaacaagc cgaagtagaa cttgggccac cctaaccggt gcttttctt
3121 cccattttct tttttctccc cctgcttca cgtctctct tccgtcttgg gccagggtg
3181 gttccagaac cgccgcgcca agtttcgcaa gcaggagcgc gcagcggcag ccgcagcggc
3241 cgggccaag aacggctcct cgggcaaaaa gtctgactct tccagggacg acgagagcaa
3301 agaggccaag agcactgacc cggacagcac tgggggcca ggtcccaatc ccaacccac
3361 cccagctgc ggggcgaatg gaggcggcgg cggcgggccc agcccggctg gagctccggg
3421 ggcgcgggg cccgggggcc cgggaggcga acccggaag ggcggcgag cagcagcggc
3481 ggcgcgcgcg gcagcggcgg cggcggcagc ggcagcggcg gcagctggag gcctggctgc
3541 ggctgggggc cctggacaag gctgggtctc cggccccggc cccatcact ccatccgga
3601 ttcgcttggg ggteecttcg gcagcgtcct atcttcgctc caaagacca acggtgccaa

3661 agccgcctta gtgaagagca gtatgttctg atctggaate ctgcggcggc ggcggcgccg
 3721 ggcacagcgg ggcagccagg gcccgggcgg ggcagtgggc gagcgggtag gcccaaggct
 3781 attgtcgtcg ctgctgccat ggctttttca ttgaggcct aaagtaateg cgctaagaat
 3841 aaagggaaaa cggcgtcgcc ctcatctca ccccaectct accccttcc tcaaccccca
 3901 aacaaaacaa acaaaacttc ctggcttcgc acctgctgg ggcctcgcag cggggccagg
 3961 gctccgcctg ctgatcgggg gttgtgagca ggcggcctg gacgcggggc actctcaggg
 4021 ggctgtgtct gcgtgtcagt ttgtgtctgt ctcggggaat gtgtgtctgt ggcccaagca
 4081 ggtgacagga agagatgggg ggcctcaacc aacttagtga cttgtttaga aaaaaagac
 4141 aaaaaagtaa aaataaaaac aaaaaagttg gaaggcagaa accattaana acaaaaagc
 4201 caacaaccca gaaaggttta aaaaacataa ggaaaaaaa gacaaattaa aggaggggct
 4261 aggggagaag ctgcagctgg agctgaaggc tcgatcttgt gaaccctaa atccgctccc
 4321 tctaacagc acggattctc ttggggctct tcttcaggga agagtaggga cgcggtcca
 4381 gcccccttc ctatcgtgc cttgggttcg ggtcactgcg ggcagcactt gctcagactg
 4441 tccggcggc cggagtgaact ttctgcacc ccttgctg tcccactcg ctgaacacca
 4501 tcccgccatt agcgcacgg aaccacac agttgcaact cccaacccg aatctttgca
 4561 gccgttcggc cctgaaagat gccctatcca tgagatgcct tttcatctgc aaactctgca
 4621 aatgtgtct catgtttcgc aactctttt tccccctcg ctccgccta ccccgctggc
 4681 attttctct tccaccagct tttactgaac tttttggcag tgctttggat tggggtaat
 4741 tgcagtcac gtaactggct gcagagaaat ctaccgagca aggaaaaggc acacacacac
 4801 gtttgcagg gtgtctcgg ttgcattct gttggaatga tccgaactgg actcacatcc
 4861 tgtatggtg atggactgta tattgagggt tccattctc gcgcagttta gacatctctg
 4921 ttttgattct ttgtttgtt ttttatttta aaaggcaca actctagata ttagttagat
 4981 gttgaggct taacttttcc ggtgtcttcc tacaactgtg ttctgtgact caattgtatc
 5041 gtgttaatat cagtgcagac tgtctctct acgtgaccgt ataatgtttt tctcgtcttg
 5101 tagtctctat ggcgtgtctt tatgggtgaa taagggtctc acgggggcaa tcttttgtgt
 5161 ttagagaggc cacggttcag acaatggat atatttttgt taccaggtgc atgtctgtct
 5221 gatttctttt ttttctctgt tggactatgt ttgtgaacat aattgtcata agttatgttt
 5281 cagatttttg aatttattta tatgtgttat aatgaatgt tctattttaa agggaaatat
 5341 ttctacatgt gcttatagtt ttccaagagt gtaccattaa cttgattgtt gataataaaa
 5401 accaaaagca agtctagcaa ttgaactctt cttttcttg attctttttt ttttttttt
 5461 tttgggttgg tcattgtttt ttttttttaa gttttttttt aaaaggat

Figure 3

1 cctgagtctc ttgaaagcac atttagctgc aagaagaaat gcaatttata agatgcttcc
61 tctctgtctc tctttgtttt taaccccctc taaccagggtg tattttttaa gaaatccgct
121 tatcgttcac ataaaccccc ttggcccact tactctatgt tacagggcgc ctgagtcttg
181 ccaatgtccc agtcctttat aacatttcat gcaacttcggg gggtaggctt gttgttaaat
241 tgagcgtgta aactctttac aaaacagggt ttctatgaca tcaaggtttc ttctccctaa
301 ccgaggggga aaaaaagaag aggacgaaga agggggaaaa cacacacact atctcaattt
361 atgcctaagg tatatgatca gttaaaaagg cttaaaagct cggggaaatt ggatcagggg
421 gaatcgtcac ccaactttca ttatttccaa gtagtgtgat tgaattaaag ggcaggggagc
481 tggttagaag ggaggatcag gggctcgggtg cgtaatgggtg tggattataa ttctaattag
541 agatgcagga atcaatgata gggagggttg acagctcagt tcccagtgcc cagcccaata
601 gacggatgag ttattgtcat gtaaaaagcg ccagcaataa gaccaaccgc tttgctattg
661 tccaagtggg aagagccaag tttattatga ggactatatg ctctagagac ctgagacaag
721 gcattctata ggaggctttt tcataaaact aggcctctgct ggtagtaagg aggccagttt
781 ggaggcagge gttgagctgt gcacatctcc ccaactccagc cacctctccc atatccatct
841 tttatttcat ttttccactt ggctgagcca tccagaacct tttcaatgta taaaatggaa
901 tattcttacc tcaattcttc tgccctacgag tccctgatgg ctgggatgga cacctcagagc
961 ctggcttcag cctatgctga ctccagttcc tgcagccagg ccagtggctt ccagtataac
1021 ccgataagga ccacttttgg ggcccagtc ccgctgcccctt cctcaccgcc gggatcctgc
1081 agcctgggca ccctcagggg ccaccagagc agtccgtacg ccgcagggtt ggaccttcag
1141 ctttctcagc ggaggaagcc gcctttccgc ccgtatatag gaagccttga ttgcatttga
1201 aaatggaaat gtgttttagta tttaccaaac gaaatttgcct tacacaaatg aaagaattta
1261 tcacgtttga agcgattgca gggaggggta attcacttac agggttacac tatectagtc
1321 acaccggaac cccccacaaa attatcttaa gctgcaaaaa tgataggeat aatttattta
1381 ctttgcgatg agacgtatag cttagaaaat aattgaatta caaagagtaa agctcattac
1441 tggcagtgtc tcttttttta agaaccgaga gcggtcaca cttcttgggc tggtcatttt
1501 tatgattatt tctttaattt attattattt ttttgcagct ctttccccca acttttgagc
1561 cgggtcaact ttctgagaat tgaaaagttc ccaaagtggg actgtttggg aacttcttcc
1621 cctggctccc ctgatattcc gactgatggt ttgggatttt tttcctctct ggttttttcc
1681 tgctgaaagc actatctcaa gtcgctcaca tcgctgtgtt tcaatccacc caaaggcgtc
1741 tgtgccagaa aggactccgc caagcccga gtttgagccc aggtttccgc agataacaaa

1801 ttctctcggt ttcttcccgc agcttctctc ggcaactctc tcgcgcggtt gtaggtagcg
1861 getgecgat gacctgacct tggagtcctc acattctagc tccacggcgc gcgagctgcc
1921 ggetgatttg ctcactttct gtctctctcg tcatactcta gttccttaca aactcttcac
1981 ggaccacggc ggccctcaacg agaagcgcaa gcagcggcgc atccgcacca ctttcaccag
2041 tgcccagctc aaagagctgg aaagggctct cgcggagact cactaccccc acatctacac
2101 tcgggaggag ctggccctga agatcgacct cacagaggcg cgagtccagg tacgcgcgcc
2161 tggaaaccga ccccgcctcg ccgcactggt cgggggaggt gtggggtgag gggcggtggt
2221 tgaaattcga agtcctggag cctcgagtga gaaggacctt gggccccatg gccgatcaga
2281 aatactggat ttggtgtggc tgtgcttctg agagaggctt agagcgcacg ctcttgcat
2341 tttatttaca gttgcgaagt gtttcccacc cgagcagaga catggggggc cttgggacgt
2401 ggatgagcga tgcaattctg gggacaggaa gtgcctgtgg tggaaaggtg gcagactttg
2461 ctcccgtatt ataagttttt ccttctcccc tcccgcctcc caaaaaaatg cctcctaact
2521 caagtgcctt taacctggcc ccatggcata taggttcatt ttcccggaaa ctgtgacttg
2581 cattagattt gcaaagggtc tgtgacttca tgaaggtaa gaaccatgac ttactccaac
2641 ctgttaaaca caggtgcgct cacgagttgg ccacagcgc tttttgggtg agccccgcac
2701 cgagaagcgg tgcgcacat tgcacgctt tccaggctca aaggccgggg atgggcagcg
2761 gagcaaacc agagaggatc ccttttctt ttaccaatta gagtttaact ttagaactta
2821 ggcttagggg tgaatggcga gctcggggct tgetcaagaa gccgactgaa cagaggccca
2881 ccaaataag gccttcctt ttccggctt tctgggacct gcgctttttt aaactctgcc
2941 gcaagcctc atgtccctg cgtgctcact ccccctaaga aagtttctcc gaaaatgcac
3001 agcaataaga agcggtagac ttggtggatg tgcgcgcggg ggtgatcaca gcgcatgggg
3061 aggaggggtg taaaacaagc cgaagtagaa cttgggccac cctaaccggt gcttttctt
3121 cccattttct tctttctccc cctgcttcc cgtctctct tccgtcttg gccaggtgtg
3181 gttccagaac cgcgcgcga agtttcgcaa gcaggagcgc gcagcggcag ccgcagcggc
3241 cgcggccaag aacggctcct cgggcaaaaa gtctgactct tccagggacg acgagagcaa
3301 agaggccaag agcactgacc eggacagcac tgggggcccc ggtcccate ccaacccac
3361 ccccagctgc ggggcgaatg gaggcggcgg cggcgggccc agcccggctg gagctccggg
3421 ggcggcgggg cccgggggccc cgggaggcga acccggcaag ggcggcgcag cagcagcggc
3481 ggcggcgcgc gcgcgggcgc cggcggcagc ggcagcggcg gcagctggag gctggtgc
3541 ggctgggggc cctggacaag gctgggctcc cggccccggc cccatcact ccatcccga
3601 ttcgcttggg ggtcccttcg gcagcgtcct atcttcgctc caaagacca acggtgcaa
3661 agccgcctta gtgaagagca gtatgtctg atctggaatc ctgcccgggc ggcggcggcg
3721 gcgacagcgg gcgagccagg gcccgggcgg gcgagtgggc gagcgggtag gcccaaggct
3781 attgtcgtcg ctgctgcat ggtttttca ttgaggcct aaagtaatcg cgctaagaat
3841 aaagggaaaa cggcgtgcgc ctcatttcaa cccactcct accccttcc tcaacccca

3901 aacaaaacaa acaaacttcc ctggcttcgc acctgcctgg ggccctgcag cggggccagg
3961 gctccgcttg ctgatcgggg gttgtgagca gcgcggcctg gacgcggggc actctcaggg
4021 ggctgtgtct gcgtgtcagt ttgtgtctgt ctcggggaat gtgtgtctgt ggcccaagca
4081 ggtgacagga agagatgggg ggccctcaacc aacttagtga cttgtttaga aaaaaagac
4141 aaaaaagtaa aaataaaaac aaaaaagttg gaaggcagaa accattaana aacaaaaage
4201 caacaacca gaaaggttta aaaaacataa ggaaaaaaa gacaaattaa aggaggggct
4261 aggggagaag ctgcagctgg agctgaagge tegatcttgt gaaccctaa atccgctccc
4321 tccaaacagc acggattctc ttggggctct tcttcagggg agagtagggg cgcctgtcca
4381 gcccccttc ctatcgtgtc cttgggttcg ggtcactgcg gcgacgactt gctcagactg
4441 tcccggcgge cggagtgact ttctgcacc ccttgcttg tcccacctcg ctgaacacca
4501 tcccgcatt agcgcacgg aacccccacac agttgcaact cccaaccccg aatctttgca
4561 gcgcttcggc cctgaaagat gccctatcca tgagatgct tttcatctgc aaactctgca
4621 aatgtgtct catgtttcgc aactcttttt tccccctcg ctcccgccta ccccgctggc
4681 atttctctt tccaccagct tttactgaac tttttggcac tgctttggat tggggtaaat
4741 tgcagtcac gtaactggct gcagagaaat ctaccgagca aggaaaagge acacacacac
4801 gtttgcaggg gtgtctcggg ttgcattctt gttggaatga tccgaactgg actcacatcc
4861 tgtatggtgg atggactgta tattgagggg tccattcttc gcgcagttta gacatctctg
4921 ttttgattct ttgttgttgt ttttatttta aaaggcacia actctagata ttagtgaat
4981 gttgaggett taactttttc ggtgtctttc tacaactgtg ttctgtgact caattgtatc
5041 gtgttaatat cagtgcagac tgtctctctc acgtgaccgt ataatgtttt tctcgtcttg
5101 tagtctctat ggcgtgtctt tatgggtgtaa taaggttctc acggggctca tcttttgtgt
5161 ttagagaggg cacggttcag acaatgggat atatttttgt tatcagggtgc atgtctgtct
5221 gatttctttt tttttctgtg tggactatgt ttgtgaacat aattgtcata agttatgttt
5281 cagatttttg aatttattta tatgtgttat aatgaatgct tctatttaaa agggaaatat
5341 ttctacatgt gcttatagtt ttccaagagt gtaccattaa cttgattggt gataataaaa
5401 accaaaagca agtctagcaa ttgaaactct cttttctctg attctttttt tttttttttt
5461 tttgggttgg tcattgtttt ttttttttaa gttttttttt aaaaggat

1 cctgagtctc ttgaaagcac atttagctgc aagaagaaat gcaatttata agatgctttc
 61 tctctgtctc tctttgtttt taacccccctc taaccagggtg tattttttaa gaaatecget
 121 tategttcac ataaaccccc ttggcccact tactctatgt tacagggcgc ctgagtcttg
 181 ccaatgtccc agtcctttat aacatttcat gcacttcggg gggtaggctt gttgttaa
 241 tgagcgtgta acactcttac aaaacagggt ttctatgaca tcaaggtttc ttctccetaa
 301 ccgaggggga aaaaaagaag aggacgaaga agggggaaaa cacacacact atctcaattt
 361 atgcctaagg tatatgatca gttaaaaagg cttaaaagct cggggaaatt ggatcagggg
 421 gaatcgtcac ccaactttca ttatttccaa gtagtgtgat tgaattaaag ggcaggggagc
 481 tggttagaag ggaggatcag gggctcgggtg cgtaatggtg tggtattaaa ttctaattag
 541 agatgcagga atcaatgata gggaggttgg acagctcagt tccccagtgc cagcccaata
 601 gacggatgag ttattgtcat gtaaaaagcg ccagcaataa gaccaaccgc tttgctattg
 661 tccaagtgga aagagccaag tttattatga ggactatatg ctctagagac ctgagacaag
 721 gcatctcata ggaggctttt tcataaaact aggcctctgct ggtagtaagg aggccagttt
 781 ggaggcaggc gttgagctgt gcacatctcc ccactccagc cacttctcc atatccatct
 841 tttatttcat ttttccactt ggctgagcca tccagaacct tttcaatgta taaaatggaa
 901 tattcttacc tcaattctc tgccctacgag tccctgatgg ctgggatgga cacctcagac
 961 ctggcttcag cctatgctga cttcagttcc tgcagccagg ccagtggctt ccagtataac
 1021 ccgataagga ccacttttgg ggccacgtcc ggctgccctt cctcacgcc gggatcctgc
 1081 agcctgggca ccctcaggga ccaccagagc agtccgtacg ccgcaggtta ggacctcag
 1141 ctttctcagc ggaggaagcc gcctttccgc ccgtatatag gaagccttga ttgcatttga
 1201 aaatggaaat gtgttttagta tttaccaaac gaaatttgc tacaacaaatg aaagaattta
 1261 tcacgtttga agcgtttgca gggaggggta attcacttac agggttacac tatectagtc
 1321 acaccgaac cgcccacaaa attatcttaa gctgccaaaa tgataggcat aatttattta
 1381 ctttgcgatg agacgtatag cttagaaaat aattgaatta caaagagtaa agctcattac
 1441 tggcagtgtc tttttttta agaaccgaga gggctcaca cttcttgggc tggtcatttt
 1501 tatgattatt tctttaattt attattattt ttttgcagct ctttccccca acttttgagc
 1561 cgggtcaact ttctgagaat tgaaaagttc ccaaagtggg actgtttggt aacttctttc
 1621 cctggctccc ctgatattcc gactgatggt ttgggatttt tttctctctt ggttttttcc
 1681 tgctgaaagc actatctcaa gtcgctcaca tcgctgctgt tcaatccacc caaaggcget
 1741 tgtgccagaa aggactccgc caagccgaa gtttgagccc aggtttccgc agataacaaa
 1801 tttctcgggt ttcttcccgc agcttctctc ggcaactctc tcgctcgggt gtaggtagcg
 1861 gctgcccgat gacctgacct tggagtcctc acattctagc tccacggccg gcgagctgcc
 1921 ggctgatttg ctcactttct gtctctctg tcatactcta gttccttaca aactcttcac
 1981 ggaccacggc ggcctcaacg agaagcgcaa gcagcggcgc atccgacca ctttcaccag
 2041 tgcccagctc aaagagctgg aaaggtctt cgcggagact cactaccccg acatctacac
 2101 tggggaggag ctggccctga agatcgacct cacagaggcg cgagtccagg tacgcgcgcc
 2161 tggaaaccga ccccgctccg ccgcactggt ccggggagggt gtgggggtgag gggcggctgg

2221 tgaattcga agtcctggag cctcgagtga gaaggaccta gggccccatg gccgatcaga
 2281 aatactggat ttggtgtggc tgtgcgttcg agagaggcctt agagcgcacg ctcttggcat
 2341 tttatttaca gttgcgaagt gtttcccacc cgagcagaga catggggggc cttgggacgt
 2401 ggatgagcga tgcaatttcg gggacaggaa gtgcctgtgg tggaaggtgt gcagactttg
 2461 ctcccgtatt ataagttttt ccttctcccc tcccggcccc caaaaaaatg cctcctaact
 2521 caagtgcctt taacctggcc ccatggcata taggttcatt ttcccggaaa ctgtgacttg
 2581 cattagattt gcaaagggtc tgtgacttca tgaaggtaa gaacctgac ttactccaac
 2641 ctgttaaaca cagggtgcgt cacgagttgg ccacagcgcc tttttgggtg agcccccgac
 2701 cgagaagcgg tgcgcacat tgcacgctt tccaggctca aaggccgggg atgggcagcg
 2761 gagcaaacc agagaggatc ccttttctt ttaccaatta gagtttaact ttagaactta
 2821 ggcttagggg tgaatggcga gctcggggct tgctcaagaa gccgactgaa cagaggccca
 2881 ccaaaataag gccttccctt ttccgggtctt tctgggacct gcgctttttt aaactctgcc
 2941 gcaagccttc atgtccctgg cgtgctcact cccctaaga aagtttctcc gaaaatgcac
 3001 agcaataaga agcggtagac ttggtggatg tgcgcgggg ggtgatcaca gcgcatgggg
 3061 aggagggtgt taaaacaagc cgaagtagaa cttgggccac cctaaccggt gcttttcttt
 3121 cccattttct tctttctccc cctgcttcc cgtctctct tccgtcttgg gccaggtgtg
 3181 gttccagaac cgccgcgcca agtttcgcaa gcaggagcgc gcagcggcag ccgacggc
 3241 cgcggccaag aacggctct cgggcaaaaa gtctgactct tccagggacg acgagagcaa
 3301 agaggccaag agcactgacc cggacagcac tgggggcccc ggtcccaatc ccaacccac
 3361 cccagctgc ggggcgaatg gaggcggcgg cggcggggcc agcccggctg gagctccggg
 3421 ggcggcgggg cccgggggcc cgggagggca acccggcaag ggcggcgcag cagcagcggc
 3481 ggcggcggcg gcagcxxxxx xxxxxxxxxxxx ggcagcggcg gcagctggag gcctggctgc
 3541 ggctgggggc cctggacaag gctgggctcc cggccccggc cccatcacct ccatcccgga
 3601 ttcgcttggg ggtcccttgc gcagcgtcct atcttctctc caaagacca acggtgcca
 3661 agccgcctta gtgaagagca gtatgtctg atctggaatc ctgcggcggc ggcggcggcg
 3721 gcgacagcgg gcgagccagg gcccgggcgg gcgagtgggc gagcgggtag gcccaaggct
 3781 attgtcgtcg ctgctgccat ggttttttca ttgagggcct aaagtaatcg cgctaagaat
 3841 aaagggaaaa cggcgtcgcc ctcatctcaa cccactcct accccttcc tcaacccca
 3901 aacaaaacaa acaaaacttc ctggcttgc acctgctgg gccctcgcag cggggccagg
 3961 gctccgctg ctgatcgggg gttgtgagca gcgcggcctg gacgcggggc actctcaggg
 4021 ggctgtgtct gcgtgtcagt ttgtgtctgt ctgggggaat gtgtgtctgt ggcccaagca
 4081 ggtgacagga agagatgggg gccctcaacc aacttagtga cttgtttaga aaaaaagac
 4141 aaaaaagtaa aaataaaaac aaaaaagttg gaaggcagaa accattaaaa aacaaaaagc
 4201 caacaacca gaaaggttta aaaaacataa gaaaaaaaaa gacaaattaa aggagggct
 4261 aggggagaag ctgcagctgg agctgaaggc tcgatcttgt gaaccctaa atccgctccc

4321 tectaacagc acggattctc ttggggetct ttttcagggg agagtaggga cgccgttcca
4381 gcccccttc ctatcgtgtc cttgggttcg ggtcactgcg gcgacgactt gtcagactg
4441 tcccggcggc cggagtgact ttctcgcacc ccttgcctg tcccacctcg ctgaacacca
4501 tcccgccatt agcgcacggg aacccccacac agttgcaact cccaaccccg aatctttgca
4561 gccgttcggc cctgaaagat gccctatcca tgagatgcct tttcatctgc aaactctgca
4621 aaatgtgtct catgtttcgc aactcttttt tccccctcg ctcccgccta ccccgteggc
4681 attttcttct tccaccagct tttactgaac tttttggcac tgctttggat tggggtcaat
4741 tgcagtccac gtaactggct gcagagaaat ctaccgagca aggaaaaggc acacacacac
4801 gtttgcaggg gtgtctcggg ttgcatttct gttggaatga tccgaactgg actcacatcc
4861 tgtatggtgg atggactgta tattgagggt tccattcttc gcgcagttta gacatctctg
4921 ttttgattct ttgttgttgt ttttatttta aaaggcacia actctagata ttagttgaat
4981 gttgaggctt taacttttcc ggtgtcttcc tacaactgtg ttctgtgact caattgtatc
5041 gtgttaatat cagtgcagac tgtctctctc acgtgaccgt ataatgtttt tctcgtcttg
5101 tagtctctat ggcgtgtctt tatgggtgaa taaggttctc acgggggtcaa tcttttgtgt
5161 ttagagagge cacggttcag acaatgggat atatttttgt tatcagggtgc atgtctgtct
5221 gatttctttt ttttctctgt tggactatgt ttgtgaacat aattgtcata agttatgttt
5281 cagatttttg aatttattta tatgtgttat aatgaatgct tctatttaaa agggaaatat
5341 ttctacatgt gcttatagtt tccaagagt gtaccattaa cttgattggt gataataaaa
5401 accaaaagca agtctagcaa ttgaactctt cttttctctg attctttttt tttttttttt
5461 tttgggttgg tcattgtttt ttttttttaa gttttttttt aaaaggat

Figure 4B

1 mykmeysyln ssayescmag mdtsslasay adfsscsqas gfqynpirtt fgatsgcpsl
61 tpgscslgtl rdhqsspyaa vpyklftdhg glnekrkqrr irttftsaql kelervfaet
121 hypdiytrees lalkidltea rvqvwfqnr akfrkqeraa aaaaaakng ssgkksdssr
181 ddeskeakst dpdstggppp npnptpscga nggggggppsp agapgaagpp gpggepgkkg
241 aaaa~~cccc~~ aaaaaaaaaa gglaaagppg qgwapggppi tsipdslgpp fgsvlsslqr
301 pngakaalvk ssmf

Figure 5




Figure 6





Figure 7






Figure 8



**PHOX2B POLYMORPHISMS AS
HIRSCHSPRUNG'S DISEASE DIAGNOSTIC
MARKERS AND METHODS BASED THEREON**

[0001] This application claims the benefit of U.S. Provisional Application No. 60/370,968, filed Apr. 8, 2002, which is incorporated by reference herein in its entirety.

1. FIELD OF THE INVENTION

[0002] The invention relates generally to polymorphisms or mutations of the PHOX2B gene. More particularly, the invention relates to polymorphisms or mutations of the PHOX2B gene that are responsible for the disease Hirschsprung's disease (HSCR), which is a neural crest-associated developmental disorder. The invention is directed to the identification, isolation, and cloning of the DNA sequence of mutants of the PHOX2B gene, as well as the characterization of their transcripts and gene products. The invention further identifies deletions, insertions, substitutions of nucleotides in the PHOX2B gene. Specifically, the invention is directed to the detection of a single base-pair polymorphism in the PHOX2B gene that is associated with HSCR. The invention also relates to methods and kits for screening for carriers of mutations of the PHOX2B gene and the diagnosis of HSCR. The invention further relates to diagnosing predisposition or susceptibility to increased risk of developing HSCR by screening for the presence of a polymorphism associated with HSCR. The invention also relates to compositions for screening for the polymorphism and treatment choices for patients having the polymorphism of the present invention. The invention further relates to providing polymorphisms in the PHOX2B gene as genetic markers and for forensic use, paternity test and genetic mapping. The invention is also directed to screening assays and therapeutic and prophylactic methods.

2. BACKGROUND OF THE INVENTION

2.1. Hirschsprung's Disease

[0003] Hirschsprung's disease (HSCR), regarded as a multigenic neurocristopathy, is a congenital disorder that involves aganglionosis, in which there is an absence of ganglion cells in the nerve plexuses of the lower digestive tract. The condition presents in the neonatal period as failure to pass meconium, chronic severe constipation, colonic distention, secondary electrolyte disturbances and, sometimes, enterocolitis and bowel perforation. The estimated population incidence is $1/5000$ live birth (Holschneider 1982). Males are 3.5-4.0 times more likely to be affected than females (Bodian and Carter 1963; Badner et al. 1990). HSCR may occur with other neurodevelopmental disorders such as Waardenburg syndrome type 4 (Pingault et al. 1998; WS4), and it may also be associated with a variety of other anomalies including chromosomal abnormalities and syndromes with Mendelian pattern of inheritance. Approximately, 20% of the HSCR cases are familial, with a sibling recurrence risk of 4%.

[0004] Aganglionosis is due to a disorder of the enteric nervous system (ENS) in which ganglion cells fail to innervate the lower gastrointestinal tract during embryonic development. The extent of the aganglionic segment is variable, and is reflected in the severity of the disease. The genetic signals critical for neural crest migration and differentiation

into enteric ganglia include genes encoding neurotrophic factors and their receptors. Substantial evidence has been presented to show that HSCR has a complex genetic etiology requiring the interaction of several unlinked genes, and possibly environmental factors, to produce the phenotype (Angrist et al. 1993; Angrist et al. 1996; Bolk et al. 2000; Doray et al. 1998; Hofstra et al. 1999; Puffenberger et al. 1994; Svensson et al. 1999). It has been postulated that the degree of expression of the receptor tyrosine kinase gene (RET) is critical for the HSCR phenotype (Bolk et al. 2000). However, only around 30% of HSCR cases are heterozygous for mutations in this gene. RET mutations do not account for all HSCR cases and lack genotype-phenotype correlation due to low penetrance, indicating that genes other than RET are implicated in the disease. This has led to the search for other susceptibility and modifying genes that could lead to clinical expression in HSCR patients. Interestingly, HSCR genes identified so far code for protein members of two important signaling pathways involved in the development of enteric ganglia: RET and endothelin receptor B (EDNBR) signaling pathways. Interaction between these two signaling pathways could modify RET expression (Chakravarti 1996), and therefore HSCR phenotype (Bolk et al. 2000; Sakai et al. 2001). Nevertheless, for almost every HSCR gene described, incomplete penetrance and variable expression of the HSCR phenotype has also been observed implying that modifying locus are needed for the expression of the HSCR phenotype (Salomon et al. 1996; Hofstra et al. 1997). Therefore, other genes implicated in RET expression and/or intestinal neurodevelopment are to be considered as candidate genes for HSCR. A number of transgenic mouse models have been established which demonstrated the involvement of these genes in enteric neuron system development. Mice that lack the function of Ret are devoid of enteric neurons and similar phenotypes are seen in mice with mutations in other components of the RET (Schuchardt et al. 1994; Shen et al. 2002) as well as EDNBR pathways (Baynash et al. 1994). In addition, mutations in the gene encoding the transcription factor SOX10 account for the Dom phenotype which is regarded as a model for HSCR (Herbarth et al. 1998). Interestingly, Sox10 is needed to modulate Ret and Ednrb expression (Lang et al. 2000; Southard-Smith et al. 1999). In humans, SOX10 mutations have been identified in patients with HSCR and pigmentary anomalies consistent with WS4 (Pingault et al. 1998).

[0005] HSCR non-Mendelian pattern suggests a complex disorder with major contributions from predisposing genetic factors, which interact with the overall genetic background and/or environmental insults to determine the phenotype. The ability to identify the genetic factors that increase the risk for HSCR would be a breakthrough for the understanding of HSCR, and other polygenic complex disorders and congenital malformations considered multifactorial in origin.

2.2. The PHOX2B Gene

[0006] The paired mesoderm homeobox 2b gene (PHOX2B) encodes a transcription factor (homeodomain protein), which is involved in the development of several noradrenergic neuron populations. In mice, PHOX2B expression starts as soon as the enteroblasts invade the foregut mesenchyme and it is maintained throughout the development into enteric neurons. In PHOX2B homozygous mutants, enteric-neuron precursors arrive at the foregut but

migration towards the mid- and hindgut is arrested (Dubreuil et al., 2000, *Development* 127:5191-5201). Consequently, the homozygous disruption of the PHOX2B gene results in the absence of enteric ganglia, feature which is reminiscent of HSCR. Furthermore, there is no Ret expression in PHOX2B mutant embryos. This indicates that regulation of Ret by PHOX2B could account for the failure of the ENS to develop (Pattyn et al., 1999, *Nature* 399:366-370).

[0007] PHOX2B transcription factor (Goridis, C., and Brunet, J. -F., *Curr. Opin. Neurobiol.* 9:47-53, 1999) is expressed in all central and peripheral noradrenergic neurons just as these neurons are acquiring their differentiated phenotype (Tiveron, M. -C., et al., *J. Neurosci.* 16:7649-7660, 1996; Pattyn, A., et al., *Development* 124:4065-4075, 1997). PHOX2B binding site contribute to dopamine- β -hydroxylase (DBH) promoter activity and forced expression of the factor can activate the promoter in DBH-negative cell lines (Swanson, D. J., et al., *J. Biol. Chem.* 272:27382-27392, 1997; Yang, C., et al., *J. Neurochem.* 71:1813-1826, 1998; Kim, H. -S., et al., *J. Neurosci.* 18:8247-8260, 1998). Moreover, loss of function experiments demonstrate that PHOX2B is an essential determinant of noradrenergic phenotype (Morin, X., et al., *Neuron* 18:411-423, 1997;).

[0008] The DNA and amino acid sequences for PHOX2B have previously been reported (Yokoyama et al. 1996). Its coding region consists of 945 base pairs (bp) allocated in three exons yielding a homeodomain protein of 314 amino acid residues (Yokoyama et al. 1999)(see FIG. 2). PHOX2B DNA and amino acid sequences are hereby incorporated by reference SEQ ID No. 1 and SEQ ID No. 2 respectively). Exon 1 stretches from base 885 to base 1126. Exon 2 stretches from base 1962 to 2149. Exon 3 stretches from 3176 to 3691. (See FIG. 8). The wildtype gene sequence for PHOX2B is provided in SEQ ID No. 1 in FIG. 1.

2.3. Polymorphisms

[0009] The genomes of all organisms undergo spontaneous mutation in the course of their continuing evolution generating variant forms of progenitor sequences (Gusella, *Ann. Rev. Biochem.* 55, 831-854 (1986)). The variant form may confer an evolutionary advantage or disadvantage relative to a progenitor form or may be neutral. In some instances, a variant form confers a lethal disadvantage and is not transmitted to subsequent generations of the organism. In other instances, a variant form confers an evolutionary advantage to the species and is eventually incorporated into the DNA of many or most members of the species and effectively becomes the progenitor form. In many instances, both progenitor and variant form(s) survive and co-exist in a species population. The coexistence of multiple forms of a sequence gives rise to polymorphisms.

[0010] Several different types of polymorphism have been reported. A restriction fragment length polymorphism (RFLP) means a variation in DNA sequence that alters the length of a restriction fragment as described in Botstein et al., *Am. J. Hum. Genet.* 32, 314-331 (1980). The restriction fragment length polymorphism may create or delete a restriction site, thus changing the length of the restriction fragment. RFLPs have been widely used in human and animal genetic analyses (see WO 90/13668; WO90/11369; Donis-Keller, *Cell* 51, 319-337 (1987); Lander et al., *Genetics* 121, 85-99 (1989)). When a heritable trait can be linked

to a particular RFLP, the presence of the RFLP in an individual can be used to predict the likelihood that the animal will also exhibit the trait.

[0011] Other polymorphisms take the form of short tandem repeats (STRs) that include tandem di-, tri- and tetra-nucleotide repeated motifs. These tandem repeats are also referred to as variable number tandem repeat (VNTR) polymorphisms. VNTRs have been used in identity and paternity analysis (U.S. Pat. No. 5,075,217; Armour et al., 1992, *FEBS Lett.* 307, 113-115; Horn et al., WO 91/14003; Jeffreys, EP 370,719), and in a large number of genetic mapping studies.

[0012] Other polymorphisms take the form of single nucleotide variations between individuals of the same species. Approximately 1 in every 1000 nucleotides differs between any two copies of the human genome (Cooper, 1996, *Hum. Genet.* 69:201-205). Some of these genetic variations, or SNPs, lead to differences in the proteins encoded by such genes. Others are "silent", residing in non-protein coding regions of the genome. Some of these polymorphisms may also result in defective protein expression (e.g., as a result of defective splicing). Other single nucleotide polymorphisms have no phenotypic effects. Such SNPs are now being used, for example, to diagnose genetic disorders, determine a predisposition to genetic disease, identify or determine the ancestry of a genetic sample, or correlate genetic sequences with phenotypic conditions, such as complex disorders or drug response and toxicity (Risch and Merikangas, 1996, *Science* 273:1516-1517). This powerful combination of genetic and molecular biological approaches is changing the face of drug development. SNPs have been correlated with Huntington's disease, Alzheimer's disease, and various forms of breast cancer. In the emerging field of pharmacogenomics, specific SNPs are being used to determine and predict a patient's susceptibility to diseases as well as drug toxicity and response. Pharmacogenomics can also provide tools to identify new targets for designing drugs and to optimize the use of existing drugs. The hope is that this understanding will ultimately lead to the early diagnosis, prevention, and treatment of genetic diseases. Single nucleotide polymorphisms are far more frequent than RFLPs, STRs and VNTRs. Some single nucleotide polymorphisms occur in protein-coding sequences, in which case, one of the polymorphic forms may give rise to the expression of a defective or other variant protein and, potentially, a genetic disease. Examples of genes, in which polymorphisms within coding sequences give rise to genetic disease include β -globin (sickle cell anemia) and CFTR (cystic fibrosis).

[0013] Single nucleotide polymorphisms can be used in the same manner as RFLPs, and VNTRs but offer several advantages. Single nucleotide polymorphisms occur with greater frequency and are spaced more uniformly throughout the genome than other forms of polymorphism. The greater frequency and uniformity of single nucleotide polymorphisms means that there is a greater probability that such a polymorphism will be found in close proximity to a genetic locus of interest than would be the case for other polymorphisms. Also, the different forms of characterized single nucleotide polymorphisms are often easier to distinguish than other types of polymorphism (e.g., by use of assays employing allele-specific hybridization probes or primers).

[0014] Despite the increased amount of nucleotide sequence data being generated in recent years, only a minute proportion of the total repository of polymorphisms in humans and other organisms has so far been identified. The paucity of polymorphisms hitherto identified is due to the large amount of work required for their detection by conventional methods. For example, a conventional approach to identifying polymorphisms might be to sequence the same stretch of oligonucleotides in a population of individuals by didoxy sequencing.

3. SUMMARY OF THE INVENTION

[0015] The present invention is based upon the observation of the present inventors that a distinct polymorphism in the PHOX2B gene is associated with HSCR. The present invention provides diagnostic screening of a population for predisposition or susceptibility to increased risk for HSCR by detecting a polymorphism in the PHOX2B gene. In accordance with the present invention, mutated form of the PHOX2B gene, which is associated with HSCR, have been identified, isolated, cloned and sequenced. Other mutations identified in the present invention in the PHOX2B gene comprise polymorphisms which include addition, deletion, rearrangement, or replacement of one or more nucleotides. The present invention, therefore, represents the first opportunity to accurately and non-invasively screen and diagnose HSCR in a substantial portion of the population. The present also provides compositions for screening for the polymorphism and improved treatment choices for patients having the polymorphism of the present invention. In addition, the present invention provides method of treatment, prophylaxis, management or amelioration of one or more symptoms associated with HSCR. Still further the PHOX2B gene may be used as genetic markers, including but not limited to, for identifying or determining the source of a genetic sample, or correlate a test sample with other genetic sample, such as for example, in forensic testing and paternity testing.

[0016] In a preferred embodiment, the mutation of the PHOX2B gene which is associated with HSCR comprises a single nucleotide polymorphism. In another preferred embodiment, the mutation of the PHOX2B gene comprises deletion of one or more nucleotides.

[0017] In specific embodiments, the present invention relates to an isolated nucleic acid molecule comprising a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 3, 4, and 6.

[0018] In a specific embodiment, the present invention relates to an isolated nucleic acid molecule comprising nucleic acid sequence of SEQ ID NO: 3 (which corresponds to the genomic sequence of the PHOX2B gene containing a mutation in which a single base A is substituted by a G at position 2250, as shown in FIG. 1 (“A→G (2250)”). In an embodiment, the mutation is a heterozygous change, in which only one allele of the PHOX2B gene harbors the A to G change, as shown in FIG. 1B. In another embodiment, the mutation is a homozygous change, in which both alleles of the PHOX2B gene harbor the A to G change, as shown in FIG. 1C.

[0019] In a specific embodiment, the present invention relates to an isolated nucleic acid molecule comprising nucleic acid sequence of SEQ ID NO: 4 (which corresponds to the genomic sequence of the PHOX2B gene containing a

mutation in which a single base A is substituted by a C at position 3493, as shown in FIG. 2 (“A→C (3493)”). In an embodiment, the mutation is a heterozygous change, in which only one allele of the PHOX2B gene harbors the A to C change. In another embodiment, the mutation is a homozygous change, in which both alleles of the PHOX2B gene harbor the A to C change.

[0020] In a specific embodiment, the present invention relates to an isolated nucleic acid molecule comprising nucleic acid sequence of SEQ ID NO: 5 (which corresponds to the genomic sequence of the PHOX2B gene comprising a fifteen bases deletion from position 3495-3499, as shown in FIG. 3 (“DEL (3495)”). In an embodiment, the mutation is a heterozygous change, in which only one allele of the PHOX2B gene harbors the deletion, as shown in FIG. 3. In another embodiment, the mutation is a homozygous change, in which both alleles of the PHOX2B gene harbor the deletion.

[0021] In one embodiment, the nucleic acid is DNA. In another embodiment, the DNA is cDNA. In another embodiment, the nucleic acid is RNA.

[0022] The present invention provides nucleic acid molecules comprising fragments having a portion of the PHOX2B gene of the invention. The nucleic acid fragments comprise nucleotide sequences corresponding to a portion of a nucleic acid molecule of the invention.

[0023] The present invention provides a cloning vector comprising a nucleic acid molecule of the invention in a host cell for the vector. The invention further provides an expression vector comprising the nucleic acid molecule of the invention operably linked to a promoter sequence capable of directing expression of the nucleic acid molecule in host cells for the vector.

[0024] The present invention provides a method of producing a mutant PHOX2B polypeptide comprising the steps of transforming host cells with a vector of the invention; culturing the cells; and isolating the mutant PHOX2B polypeptide.

[0025] The present invention provides a mutant PHOX2B polypeptide which comprises the amino acid sequence of SEQ ID NO: 6 (which corresponds to the wild-type amino acid sequence of the PHOX2B gene containing a mutation in which 5 alanine residues at positions 245-249, as shown in FIG. 3). In one embodiment, the peptide is labeled. In another embodiment, the peptide is a fusion protein. In a specific embodiment, the fusion protein is a fusion protein comprising PHOX2B polypeptide and a heterologous polypeptide. As used herein, the term “fusion protein” refers to a polypeptide that comprises an amino acid sequence of a first protein or functional fragment, analog or derivative thereof, and an amino acid sequence of a heterologous protein (i.e., a second protein or functional fragment, analog or derivative thereof different than the first protein or functional fragment, analog or derivative thereof).

[0026] The present invention provides a use of a peptide of the invention as an immunogen for the production of antibodies.

[0027] The present invention provides a method to detect the presence of a PHOX2B gene mutation that is associated with HSCR in an individual. The method comprises the

steps of: providing a biological sample comprising DNA or RNA from the individual; and assessing the DNA or RNA for the presence of the mutant PHOX2B allele of a nucleotide mutation A→G(2250), wherein the presence of the PHOX2B mutation in the genome of the individual indicates that the individual have an increased risk of predisposition or susceptibility to HSCR.

[0028] In an embodiment, the invention is directed to a method of determining the source of a first biological sample from a subject compared to a second biological sample. The method comprises: (a) providing a first biological sample from the subject; (b) testing the first biological sample for the presence of a mutation of the PHOX2B gene at nucleotide positions 2250, 3493, and/or 3495-3499 of SEQ ID NO: 1; (c) testing the second biological sample for the presence of a mutation of the PHOX2B gene at nucleotide positions 2250, 3493, and/or 3495-3499 of SEQ ID NO: 1; and (d) comparing the mutations present in step (b) and (c); wherein the presence of identical mutation in the first biological sample compared to the second biological sample indicates that the first and second biological samples are obtained from an identical source.

[0029] The present invention provides a method to detect the presence of a PHOX2B gene mutation in an individual. The method comprises the steps of: providing a biological sample comprising DNA or RNA from the individual; and assessing the DNA or RNA for the presence of the mutant PHOX2B allele of a nucleotide mutation A→C (3493).

[0030] The present invention provides a method to detect the presence of a PHOX2B gene mutation in an individual. The method comprises the steps of: providing DNA or RNA from the individual; and assessing the DNA or RNA for the presence of the mutant PHOX2B allele of a deletion, DEL (3459).

[0031] The present invention provides a method to diagnose or detect a predisposition, or increased risk to HSCR by detecting the presence of a PHOX2B gene mutation, wherein the detection can be made in combination with one or more polymorphisms in the PHOX2B gene or other genes including but not limited to, RET, EDNBR, SOX10, EDN3, and GDNF. Thus, in an embodiment of the invention, the method further comprises assessing the DNA or RNA for the presence of any HSCR-associated polymorphisms, wherein the presence of the any HSCR-associated polymorphisms in combination with the presence of at least one other polymorphism indicates that the individual has an increased risk of predisposition or susceptibility to HSCR.

[0032] The present invention provides a method to detect the presence or absence of a PHOX2B gene mutation that is associated with HSCR in an individual. The method comprises the steps of subjecting the DNA or RNA of an individual to amplification using oligonucleotide primers flanking the nucleotide polymorphism A→G(2250).

[0033] The present invention provides a method to detect the presence of a PHOX2B gene mutation in an individual. The method comprises the steps of subjecting the DNA or RNA of an individual to amplification using oligonucleotide primers flanking the nucleotide polymorphism A→C (3493), or flanking a portion or the entire deletion, DEL (3495).

[0034] The present invention provides oligonucleotides for use in an assay to detect the presence of PHOX2B gene

mutation in an individual. In specific embodiments, the oligonucleotides of the invention comprise the sequences of SEQ ID NOS: 7, 8, 9, 10, 11, and 12. The present invention provides a kit for the detection of the presence of a PHOX2B gene mutation. The kit of the invention comprises the oligonucleotides of the invention.

[0035] Another aspect of the invention is a kit for the detection of a polymorphism in the PHOX2B gene which is associated with HSCR in a subject. The sample kit comprises a first oligonucleotide of at least 8 nucleotides in length of a portion of a nucleic acid sequence of SEQ ID NO: 3, wherein the oligonucleotide is used to amplify a region comprising two or more nucleotides at position 2250 of the PHOX2B gene in the DNA or RNA from a biological sample of the subject. In another aspect of the invention the kit further comprises at least a second oligonucleotide of 8 nucleotides in length which is a reverse complement of a portion of a nucleic acid sequence of SEQ ID NO: 3, wherein the first and second oligonucleotides comprise a primer pair.

[0036] A further aspect of the invention is a method for treating a subject diagnosed as having HSCR, comprising modulating the PHOX2B level or activity. The subject can be homozygous for A→G (2250) or heterozygous for A→G (2250).

[0037] A further aspect of the invention is a method for forensic testing or paternity testing, comprising determining the genotype of a test individual. The test individual can be homozygous for A→G (2250), A→C (3493) or DEL (3495); or heterozygous for A→G (2250), A→C (3493) or DEL (3495); or compound heterozygous or compound homozygous for A→G (2250) and A→C (3493); A→G (2250) and DEL (3495); or A→C (3493) and DEL (3495).

[0038] The present invention provides an animal model for HSCR which comprises a mammal comprises a mutant nucleic acid molecule of the invention.

[0039] A further aspect of the invention is a method to screen mammals for susceptibility to HSCR, comprising, screening such mammals for a mutation in the PHOX2B gene and wherein those mammals identified as having a mutation are more susceptible to HSCR than mammals not identified as having a mutation. In a preferred embodiment, the mammal is human.

[0040] The present invention provides therapeutics for treatment, mitigation of symptoms associated with HSCR. The therapeutics modulates the level or activity of the PHOX2B gene.

4. BRIEF DESCRIPTION OF THE FIGURES

[0041] **FIGS. 1A & B.** (A) A portion of the sequence of wild-type PHOX2B (Genbank accession number AB015671)(SEQ ID NO: 1) which encode a wild-type PHOX2B protein (SEQ ID NO: 2). (B) The amino acid sequence of SEQ ID NO: 2 is encoded by the nucleic acid molecule of SEQ ID NO: 1.

[0042] **FIG. 2.** A novel polymorphism from patients with HSCR having a single base substitution at position 2250 of the wild-type sequence of the PHOX2B gene, where an A is changed to a G (SEQ ID NO: 3).

[0043] **FIG. 3.** The sequence of a novel polymorphism identified in the third exon of PHOX2B having a single base substitution at position 3493 of the PHOX2B gene sequence, where an A is changed to a C (A/C) (SEQ ID NO: 4).

[0044] **FIGS. 4A & B.** (A) The sequence of a novel polymorphism identified in the third exon of PHOX2B (SEQ ID NO: 5). (B) In this case the mutation is a fifteen-base deletion resulting in the loss of 5 amino-acid (alanine-alanine-alanine-alanine-alanine) starting at position 3495 of the PHOX2B gene sequence (SEQ ID NO: 6).

[0045] **FIGS. 5A, B, & C.** (A) A portion of the sequence of wild-type PHOX2B (Genbank accession number AB015671)(SEQ ID NO: 1) which encode a wildtype PHOX2B protein (SEQ ID NO: 2). (B) A novel polymorphism from patients with HSCR having a single base substitution at position 2250 of the wild-type sequence of the PHOX2B gene, where an A is changed to a G (SEQ ID NO: 3). The figure shows a heterozygous change, in which only one allele of the PHOX2B gene harbors the A to G change. (C) The change A to G (A/G) are shown in both alleles of the gene.

[0046] **FIGS. 6A & B.** (A) A portion of the sequence of wild-type PHOX2B. (B) A novel polymorphism identified in the third exon of PHOX2B having a single base substitution at position 3493 of the PHOX2B gene sequence, where an A is changed to a C (A/C) (SEQ ID NO: 4). It shows a heterozygous change, in which only one allele of the PHOX2B gene harbors the A to C change.

[0047] **FIGS. 7A & B.** (A) A portion of the wild-type sequence of this part of the PHOX2B gene. (B) A novel polymorphism identified in the third exon of PHOX2B (SEQ ID NO: 5). In this case the mutation is a fifteen-base deletion resulting in the loss of 5 amino-acid (alanine-alanine-alanine-alanine-alanine) starting at position 3495 of the PHOX2B gene sequence (SEQ ID NO: 6).

[0048] **FIG. 8.** A schematic diagram of Polymorphisms identified in the PHOX2B gene. The three PHOX2B exons are represented by rectangles. Black boxes indicate the homeodomain. Striped boxes indicate poly-alanine stretches (from codon 162 to 169 and from codon 245 to 264). Downward arrows show polymorphisms with frequencies that differ between patients with HSCR and controls. Upward arrows show polymorphisms found in every sample analyzed.

5. DETAILED DESCRIPTION OF THE INVENTION

[0049] The present inventors discovered that a polymorphism at the PHOX2B gene is associated with HSCR. Accordingly, the present invention provides a method for screening subjects for HSCR using the mutation as a genetic marker in the PHOX2B gene. The present invention further provides other genetic markers comprising mutations of the PHOX2B gene including various sequence polymorphisms, and/or sequence variants wherein nucleotide substitutions in the gene sequence that may or may not affect the function of the gene product. The PHOX2B gene as described herein refers to wild-type, variants, analogs, mutants, and polymorphic forms of the PHOX2B gene. The PHOX2B gene comprises not only coding sequences but also regulatory regions such as promoter, enhancer, and terminator regions.

The PHOX2B gene further includes all introns and other DNA sequences spliced from the final PHOX2B gene RNA transcript. More specifically, the invention relates to diagnostic methods of HSCR based upon a particular polymorphism in the PHOX2B gene. In certain embodiments, the present invention is directed to an isolated nucleic acid molecule comprising the nucleic acid sequence of SEQ ID NO: 1, wherein the nucleic acid molecule comprises mutations from nucleotide positions 1-2250. In specific embodiments, the mutation is selected from the group consisting of substitution, deletion, inversion, insertion, of one or more nucleotides selected from the nucleotide positions 1-2250 of SEQ ID NO: 1. In least preferred embodiments, the mutation is a substitution at nucleotide positions 3004, 5343, or 5403 of SEQ ID NO: 1.

[0050] In a specific embodiment, the polymorphism involves an adenine (A) to a guanine (G) transition at the nucleic acid position 2250 of SEQ ID NO: 1. The nucleotide sequence of this polymorphism is shown in SEQ ID NO: 3 and is known as A→G (2250).

[0051] The present invention also provides a method for diagnosing a subject for HSCR or screening a subject predisposed or susceptible to HSCR associated with a genetic polymorphism in the PHOX2B gene, said method comprising: (a) providing a biological sample from the subject; and (b) testing the sample for the presence mutations from nucleotide positions 1-2250 of SEQ ID NO: 1, wherein, the presence of the mutation indicates that the subject is genetically predisposed to HSCR.

[0052] The present invention further provides other polymorphisms, which involves an adenine (A) to cytosine (C) transition at the nucleic acid position 3493 of SEQ ID NO: 1. The nucleotide sequence of this polymorphism is shown in SEQ ID NO: 4 and is known as A→C (3493). In another embodiment, the polymorphism involves a deletion of fifteen nucleotides at nucleic acid position 3495-3499 of SEQ ID NO: 1. The nucleotide sequence of this polymorphism is shown in SEQ ID NO: 5 and is known as DEL (3495). In other embodiments, the polymorphism involves a substitution of a nucleotide at nucleotide position at 1129 or 2304. These polymorphisms may be used as genetic markers for genomic mapping or in identifying or determining the source of a genetic sample, or correlate a test sample with other biological samples, such as in forensic testing or paternity testing.

[0053] The invention provides a method of diagnosing a disease condition associated with the mutation of the PHOX2B gene of the invention. The invention provides a method of identifying a mammal, including a human, predisposed or susceptible to a risk associated with a particular genotype in the PHOX2B gene. The method comprises determining the genotype of the PHOX2B gene in the mammal. In an embodiment, the method is to screen for an individual at risk of a condition or disease such as increased risk of HSCR by identifying the polymorphisms of the invention in the PHOX2B gene. Further, the present invention provides mutant PHOX2B gene which include the coding sequences as well as the non-coding sequences found in non-human species. All DNA sequences provided herein are understood to include complementary strands unless otherwise noted. It is understood that an oligonucleotide may be selected from either strand of the PHOX2B genomic

or cDNA sequences. Furthermore, RNA equivalents can be prepared by substituting uracil for thymine, and are included in the scope of this invention, along with RNA copies of the DNA sequences of the invention isolated from cells. The oligonucleotide of the invention can be modified by the addition of peptides, labels, and other chemical moieties and are within the scope of the invention.

[0054] The present invention also provides PHOX2B proteins which includes wild-type and mutant forms of the protein, functionally active fragments, derivatives and analogs thereof. The present invention includes protein as isolated from human and animal sources, produced by enzymatic or chemical means, or through recombinant expression in an organism.

[0055] In addition, the present invention relates to the creation of appropriate animal models for the common, nonsyndromic HSCR since they do not yet exist. The present invention also provides tests for genetic factors would also serve as biomarkers, valuable for diagnosis, and useful in research on all aspects of the neural crest derived congenital malformations. A biomarker could be used prognostically for predicting whether a non-symptomatic subject is likely to transmit to his/her offsprings a predilection to develop Hirschsprung's disease. In particular there is need for a blood test for polymorphisms causing neural crest derived spectrum disorders. Families with affected members need to know whether they carry a mutation which could affect future pregnancies. The present invention further provides clinicians a test as an aid in diagnosis, and to classify subjects according to the etiology of their disease.

5.1. Mutations in the PHOX2B Gene

[0056] Specific diseases or disorders, e.g., genetic diseases or disorders, are associated with an aberrant expression of PHOX2B gene. Specifically, they may be associated with specific allelic variants of polymorphic regions of certain genes, which do not necessarily encode a mutated protein. Thus, the presence of a specific allelic variant of a polymorphic region of a gene, such as a single nucleotide polymorphism ("SNP"), in a subject can render the subject susceptible to developing a specific disease or disorder. Polymorphic regions in genes, e.g., PHOX2B gene, can be identified, by determining the nucleotide sequence of genes in populations of individuals. If a polymorphic region, e.g., SNP is identified, then the link with a specific disease can be determined by studying specific populations of individuals, e.g., individuals which developed a specific disease, such as HSCR. A polymorphic region can be located in any region of a gene, e.g., exons, in coding or non coding regions of exons, introns, and promoter region.

[0057] Gene mutations may be a single base substitution mutation resulting in an amino acid substitution, a single base substitution mutation resulting in a translational stop, an insertion mutation, a deletion mutation, or a gene rearrangement. The mutation may be located in an intron, an exon of the gene, or a promotor or other regulatory region which affects the expression of the gene or can serve as a markers in linkage disequilibrium analysis. DNA markers are used to track the inheritance of the genes through families. DNA markers are fragments of DNA with a defined physical location on a chromosome, whose inheritance can be monitored. The closer a DNA marker is to a susceptibility

gene, the greater the probability that the marker and the susceptibility gene will be passed together from parent to child. This phenomenon is called genetic linkage. Once linkage to a specific chromosomal region has been obtained, the size of the region is narrowed down using a combination of physical and genetic mapping until the region is small enough to be sequenced and the susceptibility gene can be identified. After identification of the susceptibility gene, any polymorphisms in this gene can be determined and an analysis performed to see whether these mutations occur with greater prevalence in HSCR patients compared to non-HSCR individuals. A single nucleotide polymorphism (SNP) is shown in SEQ ID No: 3, which have been identified in HSCR patient's samples. This polymorphism and other mutations are shown hereinafter in the examples in Section 6.

[0058] PHOX2B comprises polymorphic regions, we have identified a specific allele which is associated with HSCR or with an increased likelihood of developing HSCR. Thus, the invention provides methods for determining the identity of the allele or allelic variant of a polymorphic region of the PHOX2B gene in a subject, to thereby determine whether the subject is at risk of transmitting and/or developing a disease or disorder that is associated with HSCR. Knowledge of the particular alteration or alterations, resulting in defective or deficient PHOX2B gene or protein in an individual (the PHOX2B genetic profile), alone or in conjunction with information on other genetic defects contributing to the same disease (the genetic profile of the particular disease) allows, understanding of disease pathogenesis for the dominant HSCR. Tests for genetic factors would also serve as biomarkers, valuable for diagnosis, and useful in research on all aspects of the neural crest derived congenital malformations. A biomarker could be used prognostically for predicting whether a non-symptomatic subject is likely to transmit to his/her offsprings a predilection to develop HSCR. For example, a PHOX2B population profile can be performed, by determining the PHOX2B gene profile, e.g., the identity of PHOX2B genes, in a patient population having a disease, which is caused by or contributed to by a defective or deficient PHOX2B gene.

[0059] The present methods of the invention provide means for determining if a subject is likely to transmit a disease, condition or disorder that is associated with the PHOX2B gene activity. The methods for determining whether a subject is likely to transmit a disease, which might be caused by or contributed to by an aberrant PHOX2B activity is characterized as comprising detecting, in a sample of cells from the subject, the presence or absence of a genetic alteration characterized by at least one of: (i) an alteration affecting the integrity of a gene encoding an PHOX2B polypeptide, or (ii) the mis-expression of the PHOX2B gene. For example, such genetic alterations can be detected by ascertaining the existence of at least one of: (i) a deletion of one or more nucleotides from a PHOX2B gene, (ii) an addition of one or more nucleotides to an PHOX2B gene, (iii) a substitution of one or more nucleotides of an PHOX2B gene, (iv) a gross chromosomal rearrangement of an PHOX2B gene, (v) a gross alteration in the level of a messenger RNA transcript of an PHOX2B gene, (vi) aberrant modification of an PHOX2B gene, such as of the methylation pattern of the genomic DNA, (vii) the presence of a non-wild type splicing pattern of a messenger RNA transcript of PHOX2B gene, (viii) a non-wild type level of an PHOX2B polypeptide, (ix) allelic loss of an PHOX2B

gene, and/or (x) inappropriate post-translational modification of PHOX2B polypeptide.

[0060] The present invention provides several variants of polynucleotide sequence of SEQ ID No. 1. These variants comprise nucleic acid sequence of SEQ ID No: 3, SEQ ID No: 4 and SEQ ID No: 5. Only nucleic acid sequence of SEQ ID NO: 3 contains a sequence polymorphism that is correlated with HSCR.

[0061] In a specific embodiment, the present invention is directed to an isolated nucleic acid molecule comprising the nucleic acid sequence of SEQ ID NO: 1, wherein a nucleotide at position 2250 is substituted by a nucleotide base that is not adenine.

[0062] In another specific embodiment, the present invention is directed to an isolated nucleic acid molecule comprising the nucleic acid sequence of SEQ ID NO: 1, wherein a nucleotide at position 3494 is substituted by a nucleotide base that is not adenine.

[0063] In another specific embodiment, the present invention is directed to an isolated nucleic acid molecule comprising the nucleic acid sequence of SEQ ID NO: 1, wherein nucleotides from nucleotide positions 3495 to 3499 are deleted.

[0064] In another specific embodiment, the present invention is directed to an isolated nucleic acid molecule comprising at least 6 contiguous nucleotides of SEQ ID NO: 1 including a nucleotide substitution at position 2250 by a nucleotide base that is not adenine.

[0065] In another specific embodiment, the present invention is directed to an isolated nucleic acid molecule comprising at least 6 contiguous nucleotides of SEQ ID NO: 1 including a nucleotide substitution at position 3494 by a nucleotide base that is not adenine.

[0066] In another specific embodiment, the present invention is directed to an isolated nucleic acid molecule comprising at least 6 contiguous nucleotides of SEQ ID NO: 1 including one or more nucleotides at positions 3495 to 3499.

[0067] In a specific embodiment, the present invention is directed to an isolated nucleic acid molecule comprising at least 6 contiguous nucleotides of SEQ ID NO: 1 including a nucleotide substitution at position 2250 by a nucleotide base that is not adenine and that hybridizes to a nucleic acid molecule consisting of: (a) the nucleic acid sequence of SEQ ID NO: 1; or (b) a nucleotide sequence that encodes the amino acid sequence of SEQ ID NO: 2, under moderately stringent conditions, wherein the moderately stringent conditions comprise hybridization in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate, 1 mM EDTA at 65° C., and washing in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate, 1 mM EDTA at 65° C., and washing in 0.2×SSC/0.1% sodium dodecyl sulfate at 42° C., or the complement thereof.

[0068] In a specific embodiment, the present invention is directed to an isolated nucleic acid molecule comprising at least 6 contiguous nucleotides of SEQ ID NO: 1 including a nucleotide substitution at position 3493 by a nucleotide base that is not adenine and that hybridizes to a nucleic acid molecule consisting of: (a) the nucleic acid sequence of SEQ ID NO: 1; or (b) a nucleotide sequence that encodes the amino acid sequence of SEQ ID NO: 2, under moderately stringent conditions, wherein the moderately stringent con-

ditions comprise hybridization in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate, 1 mM EDTA at 65° C., and washing in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate, 1 mM EDTA at 65° C., and washing in 0.2×SSC/0.1% sodium dodecyl sulfate at 42° C., or the complement thereof.

[0069] In a specific embodiment, the present invention is directed to an isolated nucleic acid molecule comprising at least 6 contiguous nucleotides of SEQ ID NO: 1 including one or more nucleotides at positions 3495 to 3499 and that hybridizes to a nucleic acid molecule consisting of: (a) the nucleic acid sequence of SEQ ID NO: 1; or (b) a nucleotide sequence that encodes the amino acid sequence of SEQ ID NO: 2, under moderately stringent conditions, wherein the moderately stringent conditions comprise hybridization in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate, 1 mM EDTA at 65° C., and washing in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate, 1 mM EDTA at 65° C., and washing in 0.2×SSC/0.1% sodium dodecyl sulfate at 42° C., or the complement thereof.

[0070] The present invention further provides a method of determining predisposition of a subject to HSCR comprising determining the presence of a polymorphism of the invention which is associated with HSCR. In another aspect, the present invention provides a method of determining predisposition of a subject to transmit HSCR. The method comprises identifying in a biological sample from a subject, a polymorphism or haplotype (a set of polymorphisms which is inherited together as a group) in the PHOX2B gene, as compared with a wild-type DNA from a non-HSCR subject. Certain polymorphisms or haplotypes may correlate with the severity and/or nature of the HSCR phenotype, e.g. with mild, moderate or severe aganglionosis as defined by established clinical parameters. Identification of polymorphisms may therefore be useful for prognosis of various degrees of the HSCR.

[0071] The PHOX2B gene is a homeobox gene and since the Homeobox genes are highly conserved among species, the present invention provides screening for related polymorphisms in all animals. The screening method can be utilized to identify animals carrying defects in genes like those which give rise to HSCR and neurodevelopmental disorders in humans. These animals may help to study disease progression and efficacies of treatment methods. These animals include, but are not limited to, non-primate (e.g., cows, pigs, horses, chickens, cats, dogs, rats, etc.), and primate (e.g. monkey such as acynomolgous monkey). The preferred mammal to be screened is human. In specific embodiment, the individuals screened are developmentally disabled children or adults in order to determine whether they carry mutations in the PHOX2B gene that are associated with HSCR. Similarly, the parents or relatives of disabled children may be screened to determine whether they are carriers of the mutated gene, hence may be diagnosed with increased risk of a predisposition or susceptibility associated with HSCR.

[0072] As used herein, "PHOX2B gene" refers to both the wild-type and mutants of the PHOX2B gene, which include: (a) a gene containing the DNA sequence shown in FIGS. 1A, 2, 3, or 4A; (b) any DNA sequence that encodes the amino acid sequence shown in FIGS. 1B or 4B; (c) any DNA sequence that hybridizes to the complement of the DNA sequences that encode the amino acid sequence shown

in FIGS. 1A, 2, 3, or 4A, under highly stringent conditions, e.g., hybridization to filter-bound DNA in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65° C., and washing in 0.1×SSC/0.1% SDS at 68° C. (Ausubel F. M. et al., eds., 1989, Current Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & sons, Inc., New York, at page 2.10.3); or (d) any DNA sequence that hybridizes to the complement of the DNA sequences that encode the amino acid sequence shown in FIGS. 1A, 2, 3, or 4A, under moderately stringent conditions, e.g., hybridization in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate, 1 mM EDTA at 65° C., and washing in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate, 1 mM EDTA at 65° C., and washing in 0.2×SSC/0.1% sodium dodecyl sulfate at 42° C., (Ausubel et al., 1989, supra). In certain embodiments, the PHOX2B gene encodes a gene product functionally equivalent to a PHOX2B gene product encoded by sequences shown in FIGS. 1B or 4B.

[0073] In one embodiment of the invention, PHOX2B gene may also encompass fragments and degenerate variants of DNA sequences of (a) through (d), including naturally occurring variants thereof. The PHOX2B gene fragment may be a complementary DNA (cDNA) molecule or a genomic DNA molecule that may comprise one or more intervening sequences or introns, as well as regulating regions located beyond the 5' and 3' ends of the coding region or within an intron.

[0074] As used herein, the term "fragment" refers to a peptide or polypeptide comprising an amino acid sequence of at least 5 contiguous amino acid residues, at least 10 contiguous amino acid residues, at least 15 contiguous amino acid residues, at least 20 contiguous amino acid residues, at least 25 contiguous amino acid residues, at least 40 contiguous amino acid residues, at least 50 contiguous amino acid residues, at least 60 contiguous amino residues, at least 70 contiguous amino acid residues, at least contiguous 80 amino acid residues, at least contiguous 90 amino acid residues, at least contiguous 100 amino acid residues, at least contiguous 125 amino acid residues, at least 150 contiguous amino acid residues, at least contiguous 175 amino acid residues, at least contiguous 200 amino acid residues, or at least contiguous 250 amino acid residues of the amino acid sequence of another polypeptide. In a specific embodiment, a fragment of a polypeptide retains at least one function of the polypeptide.

[0075] A PHOX2B gene sequence preferably exhibits at least about 80% overall similarity at the nucleotide level to the nucleic acid sequence depicted in FIGS. 1A, 2, 3, or 4A, more preferably exhibits at least about 85-90% overall similarity to the nucleic acid sequence in FIGS. 1A, 2, 3, or 4A and most preferably exhibits at least about 95% overall similarity to the nucleic acid sequence in FIGS. 1A, 2, 3, or 4A.

[0076] To determine the percent identity of two amino acid sequences or of two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino acid or nucleic acid sequence). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same

amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % identity = number of identical overlapping positions/total number of positions × 100%). In one embodiment, the two sequences are the same length.

[0077] The determination of percent identity between two sequences can also be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul, 1990, Proc. Natl. Acad. Sci. U.S.A. 87:2264-2268, modified as in Karlin and Altschul, 1993, Proc. Natl. Acad. Sci. U.S.A. 90:5873-5877. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul et al., 1990, J. Mol. Biol. 215:403. BLAST nucleotide searches can be performed with the NBLAST nucleotide program parameters set, e.g., for score=100, wordlength=12 to obtain nucleotide sequences homologous to the nucleic acid molecules of the present invention. BLAST protein searches can be performed with the XBLAST program parameters set, e.g., to score=50, wordlength=3 to obtain amino acid sequences homologous to a protein molecule of the present invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., 1997, Nucleic Acids Res. 25:3389-3402. Alternatively, PSI-BLAST can be used to perform an iterated search which detects distant relationships between molecules (Id.). When utilizing BLAST, Gapped BLAST, and PSI-Blast programs, the default parameters of the respective programs (e.g., of XBLAST and NBLAST) can be used (see, e.g., the NCBI website). Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, 1988, CABIOS 4:11-17. Such an algorithm is incorporated in the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used.

[0078] The percent identity between two sequences can be determined using techniques similar to those described above, with or without allowing gaps. In calculating percent identity, typically only exact matches are counted. The sequence alignment techniques as used above may also be used to predict corresponding polymorphisms in all mammals. Polymorphisms in any mammal, other than human, that correspond to the polymorphisms identified in the present invention may be used to correlate a disease phenotype, such as HSCR, with the corresponding polymorphism in the mammal.

[0079] The PHOX2B gene sequences of the invention are preferably of mammalian origin, and most preferably human. Mammals, include but are not limited to, mice, rats, cats, dogs, cattle, pigs, sheep, guinea pigs and rabbits.

[0080] The invention also encompasses nucleic acid molecules encoding mutant PHOX2B, peptide fragments of PHOX2B, truncated PHOX2B, and PHOX2B fusion proteins.

[0081] The PHOX2B gene sequences of the invention further include isolated nucleic acid molecules which

hybridize under highly stringent or moderate stringent conditions to at least about 6, preferably about 12, more preferably about 18, consecutive nucleotides of the PHOX2B gene sequences of (a) through (d).

[0082] The invention also includes nucleic acid molecules, preferably DNA molecules, that hybridize to, and are therefore the complements of, the DNA sequences (a) through (d), in the preceding paragraph. Such hybridization conditions may be highly stringent or moderately stringent, as described above. In instances wherein the nucleic acid molecules are deoxyoligonucleotides ("oligos"), highly stringent conditions may refer, e.g., to washing in 6xSSC/0.05% sodium pyrophosphate at 37° C. (for 14-base oligos), 48° C. (for 17-base oligos), 55° C. (for 20-base oligos), and 60° C. (for 23-base oligos). These nucleic acid molecules may encode or act as PHOX2B gene antisense molecules useful, for example, in PHOX2B gene regulation. With respect to PHOX2B gene regulation, such techniques can be used to modulate, for example, the phenotype of HSCR. Further, such sequences may be used as part of ribozyme and/or triple helix sequences, also useful for PHOX2B gene regulation.

5.2. Screening for Mutations in the PHOX2B Gene

[0083] The present invention provides diagnostic screening of a population for pre-disposition or susceptibility to increased risk for HSCR resulting from aberrant expression of PHOX2B gene. In particular, the screening relates to detecting a polymorphism in the PHOX2B gene. With knowledge of the mutation of the PHOX2B gene as disclosed herein, diagnostic information from the screening may be used in various ways. For example, the screening may be useful in assisting the medical diagnosis of a symptomatic patient or screening for presymptomatic individuals, including prenatal diagnosis. In this application, a patient with a high probability for being affected with HSCR could be tested with the gene-based diagnostic. A positive result would show that the individual may be a carrier of HSCR or that the individual is susceptible to HSCR. Also, first degree relatives of newly diagnosed presymptomatic or symptomatic HSCR individuals may be screened and determine whether these individuals are at a higher risk for disease and, if identified, could benefit from therapeutic intervention. Once identified, such individuals could be targeted for preventative treatment with the therapeutic compositions of the invention. It may also result in substantially different management, especially prevention and treatment, with subsequent substantial improvement in mortality and morbidity from HSCR, especially in at risk populations.

[0084] The diagnostic methods of the present invention are particularly useful in combination with a subject's family history of HSCR. For example, a parent suffering from HSCR and who has a family history of HSCR, may well have a child who is susceptible to HSCR, but who has not yet shown symptoms. To alleviate the concern of the parent, as well as the child, and to take any preventive measures which might delay or prevent onset, methods of the present invention can be used to determine whether the child is a carrier of a PHOX2B mutation associated with HSCR. If so, further testing can be performed, for example, neurological tests, clinical diagnosis such as diagnosing the symptoms of HSCR including but not limited to intestinal obstruction, constipation, vomiting, abdominal distension,

enterocolitis, diarrhea, fever, septic shock, to provide further support for the predicted susceptibility.

[0085] Since marker by marker approach ignores the polygenic nature of HSCR and does not take into account possible interactions among susceptibility genes, sets of SNPs in the different susceptibility loci described for HSCR may be used to screen for HSCR in a population. Accordingly, one embodiment of the present invention is directed to assembling an extensive catalogue of SNPs in candidate genes for HSCR, including but not limited to RET, EDNRB, EDN3 and GDNF genes, for use in diagnosing, screening, and/or staging the population for different degrees of HSCR manifestation by considering mutations in all the candidate genes together.

[0086] Accordingly, in one embodiment, the present invention provides a method of screening or diagnosing, or staging HSCR in a subject. The method comprises (a) providing a biological sample from the subject; and (b) testing the sample for the presence of mutation in one or more genes selected from the group consisting of PHOX2B, RET, EDNRB, EDN3 and GDNF, wherein mutations in the PHOX2B gene and one or more genes indicate that the subject is genetically predisposed to HSCR.

[0087] Polymorphisms in the PHOX2B gene may be used as genetic markers for genetic mapping. The polymorphisms of the invention may be used in a number of methods including but not limited to, identifying or determining the source of a genetic sample, or correlate a test sample with a biological sample, including but not limited to, uses in forensic testing and paternity testing, use in the study of population genetics, such as evolution, origin of a given race, and population migration.

[0088] Accordingly, in an embodiment, the invention is directed to a method of determining the source of a first biological sample from a subject compared to a second biological sample. The method comprises: (a) providing a first biological sample from the subject; (b) testing the first biological sample for the presence of a mutation of the PHOX2B gene at nucleotide positions 2250, 3493, and/or 3495-3499 of SEQ ID NO: 1; (c) testing the second biological sample for the presence of a mutation of the PHOX2B gene at nucleotide positions 2250, 3493, and/or 3495-3499 of SEQ ID NO: 1; and (d) comparing the mutations present in step (b) and (c); wherein the presence of identical mutation in the first biological sample compared to the second biological sample indicates that the first and second biological samples are obtained from an identical source.

[0089] As discussed in Section 2.3, polymorphisms exist in different individuals of a population. These polymorphisms are referred to as polymorphic forms. Determination of which polymorphic forms occupy a set of polymorphic sites in an individual identifies a set of polymorphic forms that distinguishes the individual. See generally National Research Council, *The Evaluation of Forensic DNA Evidence* (Eds. Pollard et al., National Academy Press, DC, 1996). Since polymorphic sites are generally within a 50,000 bp region in the human genome, the probability of recombination between these polymorphic sites is low. That low probability means the haplotype (a set of polymorphic sites) should be inherited without change for at least several generations. The more sites that are analyzed, the lower the

probability that the set of polymorphic forms in one individual is the same as that in an unrelated individual. Preferably, if multiple sites are analyzed, the sites are unlinked.

[0090] The capacity to identify a distinguishing or unique set of forensic markers in an individual is useful for forensic analysis. For example, one can determine whether a blood sample from a suspect matches a blood or other tissue sample from a crime scene by determining whether the set of polymorphic forms occupying selected polymorphic sites is the same in the suspect and the sample. If the set of polymorphic markers does not match between a suspect and a sample, it can be concluded (barring experimental error) that the suspect was not the source of the sample. If the set of markers does match, one can conclude that the DNA from the suspect is consistent with that found at the crime scene. If frequencies of the polymorphic forms at the loci tested have been determined (e.g., by analysis of a suitable population of individuals), one can perform a statistical analysis to determine the probability that a match of suspect and crime scene sample would occur by chance. For calculation of the probability that two random individuals have the same polymorphic or allelic form at a given polymorphic site, see WO 95/12607. If several polymorphic loci are tested, the cumulative probability of non-identity for random individuals becomes very high (e.g., one billion to one). Such probabilities can be taken into account together with other evidence in determining the guilt or innocence of the suspect.

[0091] Polymorphisms in the PHOX2B gene may be used in paternity testing. The object of paternity testing is usually to determine whether a male is the father of a child. In most cases, the mother of the child is known and thus, the mother's contribution to the child's genotype can be traced. Paternity testing investigates whether the part of the child's genotype not attributable to the mother is consistent with that of the putative father. Paternity testing can be performed by analyzing sets of polymorphisms in the putative father and the child. If the set of polymorphisms in the child attributable to the father does not match the putative father, it can be concluded, barring experimental error, that the putative father is not the biological father. If the set of polymorphisms in the child attributable to the father does match the set of polymorphisms of the putative father, a statistical calculation can be performed to determine the probability of coincidental match. For calculation of the probability of parentage exclusion, (representing the probability that a random male will have a polymorphic form at a given polymorphic site that makes him incompatible as the father), see WO 95/12607. If several polymorphic loci are included in the analysis, the cumulative probability of exclusion of a random male is very high. This probability can be taken into account in assessing the liability of a putative father whose polymorphic marker set matches the child's polymorphic marker set attributable to his/her father.

[0092] Individuals carrying mutations in the PHOX2B gene may be detected at either the DNA, the RNA, or the protein level using a variety of techniques, such as, e.g., Southern, Northern, in situ hybridization, and PCR. A biological sample is isolated from a mammal and then tested for the presence of a mutated gene or a product thereof. In a specific embodiment, the sample is tested for the presence of a polymorphism which is associated with HSCR. In another embodiment, the sample is tested for a polymorphism which

matches that of another sample. The genomic DNA used for the diagnosis may be obtained from body cells, such as those present in peripheral blood, urine, saliva, bucca, surgical specimen, and autopsy specimens. In preferred embodiments, biological samples suitable for testing include blood, saliva, amniotic fluid, and tissue. The most preferred biological sample is blood. However, any biological sample from which genetic material or the products of the marker genes can be isolated is suitable.

5.2.1. Nucleic Acid Based Screening

[0093] The DNA may be used directly or may be amplified enzymatically in vitro through use of PCR (Saiki et al. *Science* 239:487-491 (1988)) or other in vitro amplification methods such as the ligase chain reaction (LCR) (Wu and Wallace *Genomics* 4:560-569 (1989)), strand displacement amplification (SDA) (Walker et al. *Proc. Natl. Acad. Sci. U.S.A.* 89:392-396 (1992)), self-sustained sequence replication (3SR) (Fahy et al. *PCR Methods Appl.* 1:25-33 (1992)), prior to mutation analysis. The methodology for preparing nucleic acids in a 30 form that is suitable for mutation detection is well known in the art.

[0094] The detection of mutations in specific DNA sequences, such as the PHOX2B gene, can be accomplished by a variety of methods including, but not limited to, restriction-fragment-length-polymorphism detection based on allele-specific restriction-endonuclease cleavage (Kan and Dozy, 1978, *Lancet* ii:910-912), hybridization with allele-specific oligonucleotide probes (Wallace et al., 1978 *Nucl Acids Res* 6:3543-3557), including immobilized oligonucleotides (Saiki et al., 1989 *Proc. Natl. Acad. Sci. U.S.A.* 86:6230-6234) or oligonucleotide arrays (Maskos and Southern, 1993, *Nucl Acids Res* 21:2269-2270), allele-specific PCR (Newton et al., 1989, *Nucl Acids Res* 17:2503-2516), mismatch-repair detection (MRD) (Faham and Cox, 1995, *Genome Res* 5:474-482), binding of MutS protein (Wagner et al., 1995, *Nucl Acids Res* 23:3944-3948), denaturing-gradient gel electrophoresis (DGGE) (Fisher and Lerman et al., 1983, *Proc. Natl. Acad. Sci. U.S.A.* 80:1579-1583), single-strand-conformation-polymorphism detection (Orita et al., 1983, *Genomics* 5:874-879), RNAase cleavage at mismatched base-pairs (Myers et al., 1985, *Science* 230:1242), chemical (Cotton et al. *Proc. Natl. Acad. Sci. U.S.A.* 85:4397-4401 (1988)) or enzymatic (Youil et al. *Proc. Natl. Acad. Sci. U.S.A.* 92:87-91 (1995)) cleavage of heteroduplex DNA, methods based on allele specific primer extension (Syvänen et al. *Genomics* 8:684-692 (1990)), genetic bit analysis (GBA) (Nikiforov et al. *Nucl Acids Res* 22:4167-4175 (1994)), the oligonucleotide-ligation assay (OLA) (Landegren et al. *Science* 241:1077 (1988)), the allele-specific ligation chain reaction (LCR) (Barrany *Proc. Natl. Acad. Sci. U.S.A.* 88:189-193 (1991)), gap-LCR (Abravaya et al. *Nucl Acids Res* 23:675-682 (1995)), and radioactive and/or fluorescent DNA sequencing using standard procedures well known in the art.

[0095] In preferred embodiments, screening for mutated nucleic acids can be accomplished by direct sequencing of nucleic acids. In fact, putative mutants identified by other methods may be sequenced to determine the exact nature of the mutation. Nucleic acid sequences can be determined through a number of different techniques which are well known to those skilled in the art, such as but not limited to, RNA or DNA chip hybridization. Fodor et al., 1993, *Nature*

364:555-556; Pease, 1994, Proc. Natl. Acad. Sci. USA 91(11):5022-5026. In order to sequence the nucleic acid, sufficient copies of the material must first be amplified. In preferred embodiments, amplification of a selected, or target, nucleic acid sequence may be carried out by any suitable means. See generally Kwoh, D. and Kwoh, T., Am Biotechnol Lab, 8, 14 (1990). Examples of suitable amplification techniques include, but are not limited to, polymerase chain reaction, ligase chain reaction (see Barany, Proc Natl Acad Sci USA 88, 189 (1991), strand displacement amplification (see generally Walker, G. et al., Nucleic Acids Res. 20, 1691 (1992); Walker, G. et al., Proc Natl Acad Sci USA 89, 392 (1992), transcription-based amplification (see Kwoh, D. et al., Proc Natl Acad Sci USA, 86, 1173 (1989), self-sustained sequence replication (or "3SR") (see Guatelli, J. et al., Proc Natl Acad Sci USA, 87, 1874 (1990), the replicase system (see Lizardi, P. et al., Biotechnology, 6, 1197 (1988), nucleic acid sequence-based amplification (or "NASBA") (see Lewis, R., Genetic Engineering News, 12(9), 1 (1992), the repair chain reaction (or "RCR") (see Lewis, R., Genetic Engineering News, 12(9), 1 (1992), and boomerang DNA amplification (or "BDA") (see Lewis, R., Genetic Engineering News, 12(9), 1 (1992). In a most preferred embodiment, polymerase chain reaction is used.

[0096] Single strand polymorphism assay ("SSPA") analysis and the closely related heteroduplex analysis methods may be used for screening for single-base polymorphisms (Orita, M. et al., Proc Natl Acad Sci USA, 86, 2766 (1989). In these methods, the mobility of PCR-amplified test DNA from clinical specimens is compared with the mobility of DNA amplified from normal sources by direct electrophoresis of samples in adjacent lanes of native polyacrylamide or other types of matrix gels. Single-base changes often alter the secondary structure of the molecule sufficiently to cause slight mobility differences between the normal and mutant PCR products after prolonged electrophoresis. The presence of polymorphisms, including mutations, in nucleic acids by using mass spectrometry may be used as discussed in U.S. Pat. No: 5,869,242.

[0097] Ligase chain reaction is yet another method of screening for mutated nucleic acids. Ligase chain reaction (LCR) is also carried out in accordance with known techniques. LCR is especially useful to amplify, and thereby detect, single nucleotide differences between two DNA samples. In general, the reaction involves two pairs of oligonucleotide probes: one pair binds to one strand of the sequence to be detected; the other pair binds to the other strand of the sequence to be detected. The reaction is carried out by, first, denaturing (e.g., separating) the strands of the sequence to be detected, then reacting the strands with the two pairs of oligonucleotide probes in the presence of a heat stable ligase so that each pair of oligonucleotide probes hybridize to target DNA and, if there is perfect complementarity at their junction, adjacent probes are ligated together. The hybridized molecules are then separated under denaturation conditions. The process is cyclically repeated until the sequence has been amplified to the desired degree. Detection may then be carried out in a manner like that described above with respect to PCR.

[0098] Southern hybridization is also an effective method of identifying differences in sequences. Hybridization conditions, such as salt concentration and temperature can be adjusted for the sequence to be screened. Southern blotting

and hybridizations protocols are described in Current Protocols in Molecular Biology (Greene Publishing Associates and Wiley-Interscience), pages 2.9.1-2.9.10. Probes can be labelled for hybridization with random oligomers (primarily 9-mers) and the Klenow fragment of DNA polymerase. Very high specific activity probe can be obtained using commercially available kits such as the Ready-To-Go DNA Labeling Beads (Pharmacia Biotech), following the manufacturer's protocol. Briefly, 25 ng of DNA (probe) is labelled with ³²P-dCTP in a 15 minute incubation at 37° C. Labelled probe is then purified over a ChromaSpin (Clontech) nucleic acid purification column. Possible competition of probes having high repeat sequence content, and stringency of hybridization and washdown will be determined individually for each probe used. Alternatively, fragments of a candidate gene may be generated by PCR, the specificity may be verified using a rodent-human somatic cell hybrid panel, and subcloning the fragment. This allows for a large prep for sequencing and use as a probe. Once a given gene fragment has been characterized, small probe preps can be done by gel or column purifying the PCR product.

[0099] These mismatch detection protocols use samples generated by PCR and thus require use of very little genomic template. All of these methods can provide clues regarding the location of the sequence change which leads to the appearance of anomalous bands, hence facilitating subsequent cloning and sequencing strategies.

[0100] Methods of screening for mutated nucleic acids can be carried out using either deoxyribonucleic acids ("DNA") or messenger ribonucleic acids ("mRNA") isolated from the biological sample.

[0101] In specific embodiments, DNA amplification techniques that may be used involve the use of a probe, a pair of probes, or two pairs of probes which specifically bind to DNA encoding the gene of interest, but do not bind to DNA which does not encode the gene, under the same hybridization conditions, and which serve as the primer or primers for the amplification of the gene of interest or a portion thereof in the amplification reaction.

[0102] Identification of a sequence polymorphism may be effected by conventional sequencing and sequence analysis techniques, for example as described in Cotton, R. G. H., Mutation Detection, Oxford University Press, 1997; Landegren, U., Laboratory Protocols for Mutation Detection, Oxford University Press; and R. G. H. Cotton et al, Mutation Detection, Oxford, University Press, 1998. In preferred embodiments, nucleic acid sequencing can be performed by chemical or enzymatic method. The enzymatic method relies on the ability of DNA polymerase to extend a primer, hybridized to the template to be sequenced, until a chain-terminating nucleotide is incorporated. In a preferred embodiment, the methods utilize didoexynucleotides. Primers may be labelled with radioactive or fluorescent labels. Various DNA polymerases are available including Klenow fragment, AMV reverse transcriptase, Thermus aquaticus DNA polymerase, and modified T7 polymerase.

[0103] Oligonucleotides useful in diagnostic assays are typically at least 6 consecutive nucleotides in length, and may range at least 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or 18 nucleotides in length. Such oligonucleotides can be derived from either the PHOX2B genomic or cDNA sequences.

[0104] A further aspect of the invention is an oligonucleotide of at least 8 nucleotides in length selected from nucleotides 1-46, 48-123; 120-369; 365-394; 390-540; 538-646; 643-1004; 1001-1080; 1083-1109; 1106-1304; 1301-1366; 1363-1386; 1389-1514; 1516-1778; 1773-1917; 1921-2010; 2051-2146; 2154-2209; 2234-2368; 2367-2422; 2420-2464; 2465-2491; 2488-2568; 2872-2901; 2902-2934; 2936-2954; 2449-3001; 3000-3042; 3420-3435; 3451-3708; 3703-3754; 3750-3770; 3774-3840; 3840-3962; 3964-3978; 3974-3992; 3990-4157; 4153-4251; 4257-4282; 4284-4321; 4316-4333; 4337-4391; 4386-4400; 4398-4436; 4444-4547; 4572-4714; 4709-4777; 5165-5397; and 5394-5508 of SEQ ID NO: 1, 3, or 4; or the reverse complement thereof.

[0105] A further aspect of the invention is an oligonucleotide of at least 9 nucleotides in length selected from nucleotides 1-47; 47-124; 119-370; 364-395; 389-541; 537-647; 642-1005; 1000-1081; 1082-1110; 1105-1305; 1300-1367; 1362-1387; 1388-1515; 1515-1918; 1920-2011; 2050-2147; 2153-2210; 2233-2369; 2366-2423; 2419-2465; 2464-2492; 2487-2569; 2871-2935; 2935-3002; 2999-3043; 3419-3436; 3450-3755; 3749-3771; 3773-3841; 3839-3963; 3963-3979; 3973-3993; 3989-4158; 4152-4252; 4256-4283; 4283-4334; 4336-4401; 4397-4437; 4443-4548; 4571-4778; 5164-5398; and 5393-5508 of SEQ ID NO: 1, 3, or 4; or the reverse complement thereof.

[0106] A further aspect of the invention is an oligonucleotide of at least 10 nucleotides in length selected from nucleotides 1-48; 46-125; 118-1006; 999-1082; 1081-1111; 1104-1306; 1299-1368; 1361-1388; 1387-1516; 1514-1919; 1919-2012; 2049-2148; 2152-2211; 2232-2370 2365-2424; 2418-2466; 2463-2493; 2486-2570; 2870-2936; 2934-3003; 2998-3044; 3418-3437; 3449-3772; 3772-3842; 3838-3964; 3962-3994; 3988-4284; 4282-4335; 4335-4402; 4396-4438; 4442-4549; 4570-4779; and 5163-5508 of SEQ ID NO: 1, 3, or 4; or the reverse complement thereof.

[0107] A further aspect of the invention is an oligonucleotide of at least 11 nucleotides in length selected from nucleotides 1-49; 45-1389; 1386-1517, 1513-1920; 1918-2013; 2048-2149; 2151-2212; 2231-2371; 2364-2425; 2417-2467; 2462-2571; 2869-2937; 2933-3004; 2997-3045; 3417-3438; 3448-3773; 3771-3843; 3837-3965; 3961-3995; 3987-4285; 4281-4336; 4334-4403; 4395-4439; 4441-4550; 4569-4780; and 5162-5508 of SEQ ID NO: 1, 3, or 4; or the reverse complement thereof.

[0108] A further aspect of the invention is an oligonucleotide of at least 12 nucleotides in length selected from nucleotides 1-50, 44-1390; 1385-1518; 1512-1921; 1917-2014; 2047-2150; 2150-2213; 2230-2372; 2363-2468; 2461-2572; 2868-2938; 2932-3005; 2996-3046; 3416-3439; 3447-3774; 3770-3844; 3836-3966; 3960-4286; 4280-4337; 4333-4440; 4440-4551; 4568-4781; and 5161-5508 of SEQ ID NO: 1, 3, or 4; or the reverse complement thereof.

[0109] A further aspect of the invention is an oligonucleotide of at least 13 nucleotides in length selected from nucleotides 1-51; 43-1391; 1384-1519; 1511-1922; 1916-2015; 2046-2151; 2149-2214; 2229-2469; 2460-2573; 2867-2939; 2931-3047; 3415-3440; 3446-3775; 3769-3845; 3835-3967; 3959-4287; 4279-4338; 4332-4441; 4439-4552; 4567-4782; and 5160-5508 of SEQ ID NO: 1, 3, or 4; or the reverse complement thereof.

[0110] A further aspect of the invention is an oligonucleotide of at least 14 nucleotides in length selected from

nucleotides 1-52; 42-1392; 1383-1520; 1510-1923; 1915-2016; 2045-2152; 2148-2215; 2228-2574; 2866-2940; 2930-3048; 3414-3441; 3445-3776; 3768-3968; 3959-4288; 4278-4339; 4331-4442; 4438-4553; 4566-4783; and 5159-5508 of SEQ ID NO: 1, 3, or 4; or the reverse complement thereof.

[0111] A further aspect of the invention is an oligonucleotide of at least 15 nucleotides in length selected from nucleotides 1-53; 41-1393; 1382-1521; 1509-1924; 1914-2017; 2044-2153; 2147-2216; 2227-2575; 2865-2942; 2929-3049; 3413-3442; 3444-3777; 3767-3969; 3958-4289; 4277-4340; 4330-4443; 4437-4554; 4565-4784; and 5158-5508 of SEQ ID NO: 1, 3, or 4; or the reverse complement thereof.

[0112] A further aspect of the invention is an oligonucleotide of at least 16 nucleotides in length selected from nucleotides 1-1394; 1381-1925; 1913-2018; 2043-2154; 2146-2217; 2226-2576; 2864-3050; 3412-3443; 3443-3778; 3766-4341; 4329-4444; 4436-4555; 4564-4785; and 5157-5508 of SEQ ID NO: 1, 3, or 4; or the reverse complement thereof.

[0113] A further aspect of the invention is an oligonucleotide of at least 17 nucleotides in length selected from nucleotides 1-1926; 1912-2019; 2042-2155; 2145-2218; 2225-2577; 2863-3051; 3411-3779; 3765-4342; 4329-4445; 4435-4556; 4563-4786; and 5156-5508 of SEQ ID NO: 1, 3, or 4; or the reverse complement thereof.

[0114] A further aspect of the invention is an oligonucleotide of at least 18 nucleotides in length selected from nucleotides 1-2020; 2041-2156; 2144-2219; 2224-2578; 2862-3052; 3410-3780; 3764-4446; 4434-4557; 4562-4787; and 5155-5508 of SEQ ID NO: 1, 3, or 4; or the reverse complement thereof.

[0115] A further aspect of the invention is an oligonucleotide of at least 8 nucleotides in length selected from nucleotides 1-46, 48-123; 120-369; 365-394; 390-540; 538-646; 643-1004; 1001-1080; 1083-1109; 1106-1304; 1301-1366; 1363-1386; 1389-1514; 1516-1778; 1773-1917; 1921-2010; 2051-2146; 2154-2209; 2234-2368; 2367-2422; 2420-2464; 2465-2491; 2488-2568; 2872-2901; 2902-2934; 2936-2954; 2449-3001; 3000-3042; 3420-3435; 3451-3708; 3703-3754; 3750-3770; 3774-3840; 3840-3962; 3964-3978; 3974-3992; 3990-4157; 4153-4251; 4257-4282; 4284-4321; 4316-4333; 4337-4391; 4386-4400; 4398-4436; 4444-4547; 4572-4714; 4709-4777; 5165-5397; and 5394-5493 of SEQ ID NO: 5; or the reverse complement thereof.

[0116] A further aspect of the invention is an oligonucleotide of at least 9 nucleotides in length selected from nucleotides 1-47; 47-124; 119-370; 364-395; 389-541; 537-647; 642-1005; 1000-1081; 1082-1110; 1105-1305; 1300-1367; 1362-1387; 1388-1515; 1515-1918; 1920-2011; 2050-2147; 2153-2210; 2233-2369; 2366-2423; 2419-2465; 2464-2492; 2487-2569; 2871-2935; 2935-3002; 2999-3043; 3419-3436; 3450-3755; 3749-3771; 3773-3841; 3839-3963; 3963-3979; 3973-3993; 3989-4158; 4152-4252; 4256-4283; 4283-4334; 4336-4401; 4397-4437; 4443-4548; 4571-4778; 5164-5398; and 5393-5493 of SEQ ID NO: 5; or the reverse complement thereof.

[0117] A further aspect of the invention is an oligonucleotide of at least 10 nucleotides in length selected from nucleotides 1-48; 46-125; 118-1006; 999-1082; 1081-1111;

1104-1306; 1299-1368; 1361-1388; 1387-1516; 1514-1919; 1919-2012; 2049-2148; 2152-2211; 2232-2370 2365-2424; 2418-2466; 2463-2493; 2486-2570; 2870-2936; 2934-3003; 2998-3044; 3418-3437; 3449-3772; 3772-3842; 3838-3964; 3962-3994; 3988-4284; 4282-4335; 4335-4402; 4396-4438; 4442-4549; 4570-4779; and 5163-5493 of SEQ ID NO: 5; or the reverse complement thereof.

[0118] A further aspect of the invention is an oligonucleotide of at least 11 nucleotides in length selected from nucleotides 1-49; 45-1389; 1386-1517, 1513-1920; 1918-2013; 2048-2149; 2151-2212; 2231-2371; 2364-2425; 2417-2467; 2462-2571; 2869-2937; 2933-3004; 2997-3045; 3417-3438; 3448-3773; 3771-3843; 3837-3965; 3961-3995; 3987-4285; 4281-4336; 4334-4403; 4395-4439; 4441-4550; 4569-4780; and 5162-5493 of SEQ ID NO: 5; or the reverse complement thereof.

[0119] A further aspect of the invention is an oligonucleotide of at least 12 nucleotides in length selected from nucleotides 1-50, 44-1390; 1385-1518; 1512-1921; 1917-2014; 2047-2150; 2150-2213; 2230-2372; 2363-2468; 2461-2572; 2868-2938; 2932-3005; 2996-3046; 3416-3439; 3447-3774; 3770-3844; 3836-3966; 3960-4286; 4280-4337; 4333-4440; 4440-4551; 4568-4781; and 5161-5493 of SEQ ID NO: 5; or the reverse complement thereof.

[0120] A further aspect of the invention is an oligonucleotide of at least 13 nucleotides in length selected from nucleotides 1-51; 43-1391; 1384-1519; 1511-1922; 1916-2015; 2046-2151; 2149-2214; 2229-2469; 2460-2573; 2867-2939; 2931-3047; 3415-3440; 3446-3775; 3769-3845; 3835-3967; 3959-4287; 4279-4338; 4332-4441; 4439-4552; 4567-4782; and 5160-5493 of SEQ ID NO: 5; or the reverse complement thereof.

[0121] A further aspect of the invention is an oligonucleotide of at least 14 nucleotides in length selected from nucleotides 1-52; 42-1392; 1383-1520; 1510-1923; 1915-2016; 2045-2152; 2148-2215; 2228-2574; 2866-2940; 2930-3048; 3414-3441; 3445-3776; 3768-3968; 3959-4288; 4278-4339; 4331-4442; 4438-4553; 4566-4783; and 5159-5493 of SEQ ID NO: 5; or the reverse complement thereof.

[0122] A further aspect of the invention is an oligonucleotide of at least 15 nucleotides in length selected from nucleotides 1-53; 41-1393; 1382-1521; 1509-1924; 1914-2017; 2044-2153; 2147-2216; 2227-2575; 2865-2942; 2929-3049; 3413-3442; 3444-3777; 3767-3969; 3958-4289; 4277-4340; 4330-4443; 4437-4554; 4565-4784; and 5158-5493 of SEQ ID NO: 5; or the reverse complement thereof.

[0123] A further aspect of the invention is an oligonucleotide of at least 16 nucleotides in length selected from nucleotides 1-1394; 1381-1925; 1913-2018; 2043-2154; 2146-2217; 2226-2576; 2864-3050; 3412-3443; 3443-3778; 3766-4341; 4329-4444; 4436-4555; 4564-4785; and 5157-5493 of SEQ ID NO: 5; or the reverse complement thereof.

[0124] A further aspect of the invention is an oligonucleotide of at least 17 nucleotides in length selected from nucleotides 1-1926; 1912-2019; 2042-2155; 2145-2218; 2225-2577; 2863-3051; 3411-3779; 3765-4342; 4329-4445; 4435-4556; 4563-4786; and 5156-5493 of SEQ ID NO: 5; or the reverse complement thereof.

[0125] A further aspect of the invention is an oligonucleotide of at least 18 nucleotides in length selected from

nucleotides 1-2020; 2041-2156; 2144-2219; 2224-2578; 2862-3052; 3410-3780; 3764-4446; 4434-4557; 4562-4787; and 5155-5493 of SEQ ID NO: 5; or the reverse complement thereof.

[0126] It will be appreciated that such preferred oligonucleotides can be a part of a kit for detecting polymorphisms in the PHOX2B gene, especially for the detection of polymorphisms in PHOX2B DNA or RNA in a patient sample. As will be appreciated, the mutation analysis may also be performed on samples of RNA by reverse transcription into cDNA therefrom. The kit of the present invention may be used for the identification, analysis and typing of polymorphic DNA according to the present methods. Kits according to the present invention may comprise a carrying means being compartmentalized to receive in close confinement therein one or more containers such as vials, tubes, bottles and the like. Each of such containers may comprise components or a mixture of components needed to perform genotyping analysis. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or diagnostic products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

5.2.2. Protein Based Screening

[0127] Mutations may also be detected at the protein level using antibodies specific for the mutant PHOX2B protein. It may also be possible to base a PHOX2B mutation assay on altered cellular or subcellular localization of the mutant form of the PHOX2B protein. Alternatively, the detection of a mutated gene associated with HSCR can be carried out by collecting a biological sample and testing for the presence or form of the protein produced by the gene. The mutation in the gene may result in the production of a mutated form of the peptide or the lack of production of the gene product.

[0128] In an embodiment, the determination of the presence of the mutant PHOX2B protein can be carried out, for example, by isoelectric focusing, protein sizing, or immunoassay. In an immunoassay, an antibody that selectively binds to the mutated protein can be utilized (for example, an antibody that selectively binds to the mutated form of PHOX2B encoded protein). Such methods for isoelectric focusing and immunoassay are well known in the art, and are discussed in further detail below.

[0129] Changes in the size or charge of the polypeptide can be identified by isoelectric focusing or protein sizing techniques. Changes resulting in amino acid substitutions, where the substituted amino acid has a different charge than the original amino acid, can be detected by isoelectric focusing. Isoelectric focusing of the polypeptide through a gel having an ampholine gradient at high voltages separates proteins by their pI. The pH gradient gel can be compared to a simultaneously run gel containing the wild-type protein. Protein sizing techniques such as protein electrophoresis and sizing chromatography can also be used to detect changes in the size of the product.

[0130] As an alternative to isoelectric focusing or protein sizing, the step of determining the presence of the mutated polypeptides in a sample may be carried out by an antibody assay with an antibody which selectively binds to the mutated polypeptides (i.e., an antibody which binds to the

mutated polypeptides but exhibits essentially no binding to the wild-type polypeptide without the polymorphism in the same binding conditions).

[0131] Antibody assays may, in general, be homogeneous assays or heterogeneous. In a homogeneous assay, the immunological reaction usually involves the specific antibody, a labeled analyte, and the sample of interest. The signal arising from the label is modified, directly or indirectly, upon the binding of the antibody to the labeled analyte. Both the immunological reaction and detection of the extent thereof are carried out in a homogeneous solution. Immunochemical labels which may be employed include, but are not limited to, free radicals, radioisotopes, fluorescent dyes, enzymes, bacteriophages, and coenzymes.

[0132] In a heterogeneous assay approach, the reagents are usually the specimen, the antibody of the invention and means for producing a detectable signal. Similar specimens as described above may be used. The antibody is generally immobilized on a support, such as a bead, plate, or slide, and contacted with the specimen suspected of containing the antigen in a liquid phase. The support is then separated from the liquid phase and either the support phase or the liquid phase is examined for a detectable signal employing means for producing such signal. The signal is related to the presence of the analyte in the specimen. Means for producing a detectable signal include, but are not limited to, the use of radioactive labels, fluorescent labels, enzyme labels, and so forth. For example, if the antigen to be detected contains a second binding site, an antibody which binds to that site can be conjugated to a detectable group and added to the liquid phase reaction solution before the separation step. The presence of the detectable group on the solid support indicates the presence of the antigen in the test sample. Examples of suitable immunoassays are the radioimmunoassay, immunofluorescence methods, enzyme-linked immunoassays, and the like.

[0133] Those skilled in the art will be familiar with numerous specific immunoassay formats and variations thereof which may be useful for carrying out the method disclosed herein. See U.S. Pat. Nos. 4,727,022, 4,659,678, 4,376,110, 4,275,149, 4,233,402, and 4,230,767.

[0134] Antibodies to distinct isoforms of the PHOX2B gene (i.e., wild-type or mutant-specific antibodies) can be used to detect the presence or absence of the wild-type or mutant gene products by immunoblotting (Western blotting) or other immunostaining methods. Thus, in accordance with another aspect of the present invention, a kit is provided that is suitable for use in screening and assaying for the presence of the PHOX2B gene by an immunoassay through the use of an antibody which specifically binds to a gene product of the PHOX2B gene in combination with a reagent for detecting the binding of the antibody to the gene product.

5.3. Correlation of Polymorphisms with Phenotypic Traits

[0135] The polymorphisms described in the present invention may contribute to the phenotype of an organism in different ways. Some polymorphisms occur within a protein coding sequence and contribute to phenotype by affecting protein structure. The effect may be neutral, beneficial or detrimental, or both beneficial and detrimental, depending on the circumstances. For example, a heterozygous sickle

cell mutation confers resistance to malaria, but a homozygous sickle cell mutation is usually lethal. Other polymorphisms occur in noncoding regions but may exert phenotypic effects indirectly via influence on replication, transcription, and translation. A single polymorphism may affect more than one phenotypic trait. Likewise, a single phenotypic trait may be affected by polymorphisms in different genes. Further, some polymorphisms predispose an individual to a distinct mutation that is causally related to a certain phenotype.

[0136] Phenotypic traits include diseases that have known but unmapped genetic components. Phenotypic traits also include symptoms of, or susceptibility to, multifactorial diseases of which a component is or may be genetic, such as autoimmune diseases, inflammation, cancer, diseases of the nervous system, and infection by pathogenic microorganisms. Some examples of autoimmune diseases include rheumatoid arthritis, multiple sclerosis, diabetes (insulin-dependent and non-independent), systemic lupus erythematosus and Graves disease. Some examples of cancers include cancers of the bladder, brain, breast, colon, esophagus, kidney, leukemia, liver, lung, oral cavity, ovary, pancreas, prostate, skin, stomach and uterus. Phenotypic traits also include characteristics such as longevity, appearance (e.g., baldness, obesity), strength, speed, endurance, fertility, and susceptibility or receptivity to particular drugs or therapeutic treatments. In the present invention, a particular polymorphism is related to predisposition or susceptibility to HSCR. More specifically, certain polymorphisms in the PHOX2B gene is related to neural crest derived congenital malformation.

[0137] Correlation is performed for a population of individuals who have been tested for the presence or absence of a phenotypic trait of interest and for polymorphic marker sets. To perform such analysis, the presence or absence of a set of polymorphisms (i.e. a polymorphic set) is determined for a set of individuals, some of whom exhibit a particular trait, and some of which exhibit lack of the trait. The alleles of each polymorphism of the set are then reviewed to determine whether the presence or absence of a particular allele is associated with the trait of interest. Correlation can be performed by standard statistical methods such as a X^2 test and statistically significant correlations between polymorphic form(s) and phenotypic characteristics are noted. For example, it might be found that the presence of allele A1 at polymorphism A correlates with heart disease. As a further example, it might be found that the combined presence of allele A1 at polymorphism A and allele B1 at polymorphism B correlates with increased milk production of a farm animal.

[0138] Such correlations can be exploited in several ways. In the case of a strong correlation between a set of one or more polymorphic forms and a disease for which treatment is available, detection of the polymorphic form set in a human or animal patient may justify immediate administration of treatment, or at least the institution of regular monitoring of the patient. Detection of a polymorphic form correlated with serious disease in a couple contemplating a family may also be valuable to the couple in their reproductive decisions. For example, the female partner might elect to undergo in vitro fertilization to avoid the possibility of transmitting such a polymorphism from her husband to her offspring. In the case of a weaker, but still statistically

significant correlation between a polymorphic set and human disease, immediate therapeutic intervention or monitoring may not be justified. Nevertheless, the patient can be motivated to begin simple life-style changes (e.g., diet, exercise) that can be accomplished at little cost to the patient but confer potential benefits in reducing the risk of conditions to which the patient may have increased susceptibility by virtue of variant alleles. Identification of a polymorphic set in a patient correlated with enhanced receptiveness to one of several treatment regimes for a disease indicates that this treatment regime should be followed.

[0139] For animals and plants, correlations between characteristics and phenotype are useful for breeding for desired characteristics. For example, Beitz et al., U.S. Pat. No. 5,292,639 discuss use of bovine mitochondrial polymorphisms in a breeding program to improve milk production in cows.

5.4. Genetic Mapping of Phenotypic Traits

[0140] The present section describes identification of a physical linkage between a genetic locus associated with a trait of interest and polymorphic markers that are not associated with the trait, but are in physical proximity with the genetic locus responsible for the trait and co-segregate with it. Such analysis is useful for mapping a genetic locus associated with a phenotypic trait to a chromosomal position, and thereby cloning gene(s) responsible for the trait. See Lander et al., Proc. Natl. Acad. Sci. (USA) 83, 7353-7357 (1986); Lander et al., Proc. Natl. Acad. Sci. (USA) 84, 2363-2367 (1987); Donis-Keller et al., Cell 51, 319-337 (1987); Lander et al., Genetics 121, 185-199 (1989). Genes localized by linkage can be cloned by a process known as directional cloning. See Wainwright, Med. J. Australia 159, 170-174 (1993); Collins, Nature Genetics 1, 3-6 (1992).

[0141] Linkage studies are typically performed on members of a family. Available members of the family are characterized for the presence or absence of a phenotypic trait and for a set of polymorphic markers. The distribution of polymorphic markers in an informative meiosis is then analyzed to determine which polymorphic markers co-segregate with a phenotypic trait. See, e.g., Kerem et al., Science 245, 1073-1080 (1989); Monaco et al., Nature 316, 842 (1985); Yamoka et al., Neurology 40, 222-226 (1990); Rossiter et al., FASEB Journal 5, 21-27 (1991).

[0142] Linkage is analyzed by calculation of LOD (log of the odds) values. A lod value is the relative likelihood of obtaining observed segregation data for a marker and a genetic locus when the two are located at a recombination fraction θ , versus the situation in which the two are not linked, and thus segregating independently (Thompson & Thompson, Genetics in Medicine (5th ed, W. B. Saunders Company, Philadelphia, 1991); Strachan, "Mapping the human genome" in The Human Genome (BIOS Scientific Publishers Ltd, Oxford), Chapter 4). A series of likelihood ratios are calculated at various recombination fractions (θ), ranging from $\theta=0.0$ (coincident loci) to $\theta=0.50$ (unlinked). Thus, the likelihood at a given value of θ is: probability of data if loci linked at θ to probability of data if loci unlinked. The computed likelihoods are usually expressed as the \log_{10} of this ratio (i.e., a lod score). For example, a lod score of 3 indicates 1000:1 odds against an apparent observed linkage being a coincidence. The use of

logarithms allows data collected from different families to be combined by simple addition. Computer programs are available for the calculation of lod scores for differing values of θ (e.g., LIPED, MLINK (Lathrop, Proc. Nat. Acad. Sci. (USA) 81, 3443-3446 (1984)). For any particular lod score, a recombination fraction may be determined from mathematical tables. See Smith et al., Mathematical tables for research workers in human genetics (Churchill, London, 1961); Smith, Ann. Hum. Genet. 32, 127-150 (1968). The value of θ at which the lod score is the highest is considered to be the best estimate of the recombination fraction.

[0143] Positive lod score values suggest that the two loci are linked, whereas negative values suggest that linkage is less likely (at that value of θ) than the possibility that the two loci are unlinked. By convention, a combined lod score of +3 or greater (equivalent to greater than 1000:1 odds in favor of linkage) is considered definitive evidence that two loci are linked. Similarly, by convention, a negative lod score of -2 or less is taken as definitive evidence against linkage of the two loci being compared. Negative linkage data are useful in excluding a chromosome or a segment thereof from consideration. The search focuses on the remaining non-excluded chromosomal locations.

5.5. Production of Mutant PHOX2B Gene Product

[0144] In various embodiments of the invention, sequences encoding wild-type or mutant PHOX2B gene product are inserted into an expression vector for propagation and expression in recombinant cells. In a specific embodiment, a mutant PHOX2B gene product comprises the amino acid sequence of SEQ ID NO: 6, which is encoded by a nucleic acid molecule comprising the nucleic acid sequence of SEQ ID NO: 5. The mutant PHOX2B gene product differs from the wild-type PHOX2B protein comprising the amino acid sequence of SEQ ID NO: 2 in that the mutant PHOX2B protein comprises a deletion of five alanine residues at amino acid position 245-249 of SEQ ID NO: 2.

[0145] In a specific embodiment, a lentiviral vector can be used to express PHOX2B or modified gene product (Case et al., 1999, Proc. Natl. Acad. Sci. 96: 2988-2993; Miyoshi et al., 1998, J. Virology 72: 8150-8157).

[0146] An expression construct, as used herein, refers to a nucleotide sequence encoding a PHOX2B gene operably associated with one or more regulatory regions which enables expression of the PHOX2B gene in an appropriate host cell. "Operably-associated" refers to an association in which the regulatory regions and the PHOX2B sequence to be expressed are joined and positioned in such a way as to permit transcription, and ultimately, translation.

[0147] Such expression construct can also be used to produce antisense PHOX2B RNA, provided that the nucleotide sequence encoding PHOX2B is inserted in the reverse orientation.

[0148] The regulatory regions necessary for transcription of the PHOX2B can be provided by the expression vector. A translation initiation codon (ATG) may also be provided if the PHOX2B gene sequence lacking its cognate initiation codon is to be expressed. In a compatible host-expression construct system, cellular transcriptional factors, such as RNA polymerase, will bind to the regulatory regions on the

expression construct to effect transcription of the PHOX2B sequence in the host cell. The precise nature of the regulatory regions needed for gene expression may vary from host cell to host cell. Generally, a promoter is required which is capable of binding RNA polymerase and promoting the transcription of an operably-associated nucleic acid sequence. Such regulatory regions may include those 5'-non-coding sequences involved with initiation of transcription and translation, such as the TATA box, capping sequence, CAAT sequence, and the like. The non-coding region 3' to the coding sequence may contain transcriptional termination regulatory sequences, such as terminators and polyadenylation sites.

[0149] Both constitutive and inducible regulatory regions may be used for expression of the PHOX2B gene. It may be desirable to use inducible promoters to control the high level expression of the PHOX2B gene once the expression construct is introduced into patient's cells in vivo. Examples of useful regulatory regions are provided below.

[0150] In order to attach DNA sequences with regulatory functions, such as promoters, to the PHOX2B gene sequence or to insert the PHOX2B gene sequence into the cloning site of a vector, linkers or adapters providing the appropriate compatible restriction sites may be ligated to the ends of the cDNAs by techniques well known in the art (Wu et al., 1987, *Methods in Enzymol* 152:343-349). Cleavage with a restriction enzyme can be followed by modification to create blunt ends by digesting back or filling in single-stranded DNA termini before ligation. Alternatively, a desired restriction enzyme site can be introduced into a fragment of DNA by amplification of the DNA by use of PCR with primers containing the desired restriction enzyme site.

[0151] An expression construct comprising a PHOX2B sequence operably associated with regulatory regions can be directly introduced into appropriate host cells for expression and production of PHOX2B protein without further cloning. See, for example, U.S. Pat. No. 5,580,859. The expression constructs can also contain DNA sequences that facilitate integration of the PHOX2B sequence into the genome of the host cell, e.g., via homologous recombination. In this instance, it is not necessary to employ an expression vector comprising a replication origin suitable for appropriate host cells in order to propagate and express the PHOX2B in the host cells.

5.6. Host-Vector Systems

[0152] Described herein are systems of vectors and host cells that can be used for the expression of PHOX2B. A variety of expression vectors may be used in the present invention which include, but are not limited to, plasmids, cosmids, phage, phagemids, or modified viruses. Typically, such expression vectors comprise a functional origin of replication for propagation of the vector in an appropriate host cell, one or more restriction endonuclease sites for insertion of the PHOX2B gene sequence, and one or more selection markers. The expression vector must be used with a compatible host cell which may be derived from a prokaryotic or an eukaryotic organism including but not limited to bacteria, yeasts, insects, mammals, and humans.

[0153] In one aspect, expression constructs and vectors are introduced into host cells for the purpose of producing the PHOX2B. Any cell type that can produce mammalian pro-

teins and is compatible with the expression vector may be used, including those that have been cultured in vitro or genetically engineered. Host cells may be obtained from normal or affected subjects, including healthy humans, private laboratory deposits, public culture collections such as the American Type Culture Collection, or from commercial suppliers.

[0154] Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins. A host cell may be chosen which modifies and processes the expressed gene products in a specific fashion similar to the way the recipient processes its PHOX2B. For the purpose of producing large amounts of PHOX2B for administration to a subject, it is preferable that the type of host cell used in the present invention has been used for expression of heterologous genes, and is reasonably well characterized and developed for large-scale production processes.

[0155] Vectors based on *E. coli* are the most popular and versatile systems for high level expression of foreign proteins (Makrides, 1996, *Microbiol Rev*, 60:512-538). Non-limiting examples of regulatory regions that can be used for expression in *E. coli* may include but not limited to lac, trp, lpp, phoA, recA, tac, T3, T7 and λP_L (Makrides, 1996, *Microbiol Rev*, 60:512-538). Non-limiting examples of prokaryotic expression vectors may include the λ gt vector series such as λ gt11 (Huynh et al., 1984 in "DNA Cloning Techniques", Vol. I: A Practical Approach (D. Glover, ed.), pp. 49-78, IRL Press, Oxford), and the pET vector series (Studier et al., 1990, *Methods Enzymol.*, 185:60-89). However, a potential drawback of a prokaryotic host-vector system is the inability to perform many of the post-translational processing of mammalian cells. Thus, a eukaryotic host-vector system is preferred, a mammalian host-vector system is more preferred, and a human host-vector system is the most preferred.

[0156] For expression of PHOX2B in mammalian host cells, a variety of regulatory regions can be used, for example, the SV40 early and late promoters, the cytomegalovirus (CMV) immediate early promoter, and the Rous sarcoma virus long terminal repeat (RSV-LTR) promoter. Inducible promoters that may be useful in mammalian cells include but are not limited to those associated with the metallothionein gene, mouse mammary tumor virus glucocorticoid responsive long terminal repeats (MMTV-LTR), β -interferon gene, and hsp70 gene (Williams et al., 1989, *Cancer Res.* 49:2735-42 ; Taylor et al., 1990, *Mol. Cell Biol.*, 10:165-75).

[0157] The following animal regulatory regions, which exhibit tissue specificity and have been utilized in transgenic animals, can also be used in tumor cells of a particular tissue type: elastase I gene control region which is active in pancreatic acinar cells (Swift et al., 1984, *Cell* 38:639-646; Ornitz et al., 1986, *Cold Spring Harbor Symp. Quant. Biol.* 50:399-409; MacDonald, 1987, *Hepatology* 7:425-515); insulin gene control region which is active in pancreatic beta cells (Hanahan, 1985, *Nature* 315:115-122), immunoglobulin gene control region which is active in lymphoid cells (Grosschedl et al., 1984, *Cell* 38:647-658; Adames et al., 1985, *Nature* 318:533-538; Alexander et al., 1987, *Mol. Cell. Biol.* 7:1436-1444), mouse mammary tumor virus control region which is active in testicular, breast, lymphoid

and mast cells (Leder et al., 1986, Cell 45:485-495), albumin gene control region which is active in liver (Pinkert et al., 1987, Genes and Devel. 1:268-276), alpha-fetoprotein gene control region which is active in liver (Krumlauf et al., 1985, Mol. Cell. Biol. 5:1639-1648; Hammer et al., 1987, Science 235:53-58; alpha 1-antitrypsin gene control region which is active in the liver (Kelsey et al., 1987, Genes and Devel. 1:161-171), beta-globin gene control region which is active in myeloid cells (Mogram et al., 1985, Nature 315:338-340; Kollias et al., 1986, Cell 46:89-94; myelin basic protein gene control region which is active in oligodendrocyte cells in the brain (Readhead et al., 1987, Cell 48:703-712); myosin light chain-2 gene control region which is active in skeletal muscle (Sani, 1985, Nature 314:283-286); gonadotropic releasing hormone gene control region which is active in the hypothalamus (Mason et al., 1986, Science 234:1372-1378); probasin and prostate-specific antigen (PSA) gene control region which is active in the prostate (Brookes et al., 1998, The Prostate 35:18-26).

[0158] The efficiency of expression of the PHOX2B in a host cell may be enhanced by the inclusion of appropriate transcription enhancer elements in the expression vector, such as those found in SV40 virus, Hepatitis B virus, cytomegalovirus, immunoglobulin genes, metallothionein, β -actin (see Bittner et al., 1987, Methods in Enzymol. 153:516-544; Gorman, 1990, Curr. Op. in Biotechnol. 1:36-47).

[0159] The expression vector may also contain sequences that permit maintenance and replication of the vector in more than one type of host cell, or integration of the vector into the host chromosome. Such sequences may include but are not limited to replication origins, autonomously replicating sequences (ARS), centromere DNA, and telomere DNA. It may also be advantageous to use shuttle vectors which can be replicated and maintained in at least two types of host cells.

[0160] In addition, the expression vector may contain selectable or screenable marker genes for initially isolating, identifying or tracking host cells that contain DNA encoding a PHOX2B. For long term, high yield production of PHOX2B-peptide complexes, stable expression in mammalian cells is preferred. A number of selection systems may be used for mammalian cells, including but not limited to the Herpes simplex virus thymidine kinase (Wigler et al., 1977, Cell 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalski and Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and adenine phosphoribosyltransferase (Lowy et al., 1980, Cell 22:817) genes can be employed in tk⁻, hgp^rt⁻ or apr^t- cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for dihydrofolate reductase (dhfr), which confers resistance to methotrexate (Wigler et al., 1980, Natl. Acad. Sci. USA 77:3567; O'Hare et al., 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, Proc. Natl. Acad. Sci. USA 78:2072); neomycin phosphotransferase (neo), which confers resistance to the aminoglycoside G-418 (Colberre-Garapin et al., 1981, J. Mol. Biol. 150:1); and hygromycin phosphotransferase (hyg), which confers resistance to hygromycin (Santerre et al., 1984, Gene 30:147). Other selectable markers, such as but not limited to histidinol and ZeocinTM can also be used.

[0161] Preferred mammalian host cells include but are not limited to those derived from humans, monkeys and rodents, (see, for example, Kriegler M. in "Gene Transfer and Expression: A Laboratory Manual", New York, Freeman & Co. 1990).

[0162] A number of viral-based expression systems may also be utilized with mammalian cells to produce PHOX2B. Vectors using DNA virus backbones have been derived from simian virus 40 (SV40) (Hamer et al., 1979, Cell 17:725), adenovirus (Van Doren et al., 1984, Mol Cell Biol 4:1653), adeno-associated virus (McLaughlin et al., 1988, J Virol 62:1963), and bovine papillomas virus (Zinn et al., 1982, Proc Natl Acad Sci 79:4897). In cases where an adenovirus is used as an expression vector, the donor DNA sequence may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing heterologous products in infected hosts. (See e.g., Logan and Shenk, 1984, Proc. Natl. Acad. Sci. (USA) 81:3655-3659).

[0163] Bovine papillomavirus (BPV) can infect many higher vertebrates, including man, and its DNA replicates as an episome. A number of shuttle vectors have been developed for recombinant gene expression which exist as stable, multicopy (20-300 copies/cell) extrachromosomal elements in mammalian cells. Typically, these vectors contain a segment of BPV DNA (the entire genome or a 69% transforming fragment), a promoter with a broad host range, a polyadenylation signal, splice signals, a selectable marker, and "poisonless" plasmid sequences that allow the vector to be propagated in *E. coli*. Following construction and amplification in bacteria, the expression gene construct are transfected into cultured mammalian cells by, for example, the calcium phosphate coprecipitation technique. For those host cells that do not manifest a transformed phenotype, selection of transformants is achieved by use of a dominant selectable marker, such as histidinol and G418 resistance. For example, a PHOX2B gene sequence can be inserted into BPV vectors, such as pBCMGSNeo and pBCMHis (Karasuyama et al., Eur. J. Immunol. 18:97-104; Ohe et al., Human Gene Therapy, 6:325-33) which can then be transfected into a diverse range of cell types for expression of the PHOX2B.

[0164] Alternatively, the vaccinia 7.5K promoter may be used. (See, e.g., Mackett et al., 1982, Proc. Natl. Acad. Sci. (USA) 79:7415-7419; Mackett et al., 1984, J. Virol. 49:857-864; Panicali et al., 1982, Proc. Natl. Acad. Sci. 79:4927-4931.) In cases where a human host cell is used, vectors based on the Epstein-Barr virus (EBV) origin (OriP) and EBV nuclear antigen 1 (EBNA-1; a trans-acting replication factor) can be used. Such vectors can be used with a broad range of human host cells, e.g., EBO-pCD (Spickofsky et al., 1990, DNA Prot Eng Tech 2:14-18); pDR2 and λ DR2 (available from Clontech Laboratories).

[0165] PHOX2B may also be made with a retrovirus-based expression system. Retroviruses, such as Moloney murine leukemia virus, can be used since most of the viral gene sequence can be removed and replaced with PHOX2B gene sequence while the missing viral functions can be supplied in trans. In contrast to transfection, retroviruses can

efficiently infect and transfer genes to a wide range of cell types including, for example, primary hematopoietic cells. Moreover, the host range for infection by a retroviral vector can be manipulated by the choice of envelope used for vector packaging.

[0166] For example, a retroviral vector can comprise a 5' long terminal repeat (LTR), a 3' LTR, a packaging signal, a bacterial origin of replication, and a selectable marker. The PHOX2B DNA is inserted into a position between the 5' LTR and 3' LTR, such that transcription from the 5' LTR promoter transcribes the cloned DNA. The 5' LTR comprises a promoter, including but not limited to an LTR promoter, an R region, a U5 region and a primer binding site, in that order. Nucleotide sequences of these LTR elements are well known in the art. A heterologous promoter as well as multiple drug selection markers may also be included in the expression vector to facilitate selection of infected cells. See, McLaughlin et al., 1990, *Prog Nucleic Acid Res and Molec Biol* 38:91-135; Morgenstern et al., 1990, *Nucleic Acid Res* 18:3587-3596; Choulika et al., 1996, *J Virol* 70:1792-1798; Boesen et al., 1994, *Biotherapy* 6:291-302; Salmons and Gunzberg, 1993, *Human Gene Therapy* 4:129-141; and Grossman and Wilson, 1993, *Curr. Opin. in Genetics and Devel.* 3:110-114.

[0167] Other useful eukaryotic host-vector system may include yeast and insect systems. In yeast, a number of vectors containing constitutive or inducible promoters may be used with *Saccharomyces cerevisiae* (baker's yeast), *Schizosaccharomyces pombe* (fission yeast), *Pichia pastoris*, and *Hansenula polymorpha* (methylotropic yeasts). For a review see, *Current Protocols in Molecular Biology*, Vol. 2, 1988, Ed. Ausubel et al., Greene Publish. Assoc. & Wiley Interscience, Ch. 13; Grant et al., 1987, *Expression and Secretion Vectors for Yeast*, in *Methods in Enzymology*, Eds. Wu & Grossman, 1987, Acad. Press, N.Y., Vol. 153, pp. 516-544; Glover, 1986, *DNA Cloning*, Vol. II, IRL Press, Wash., D.C., Ch. 3; and Bitter, 1987, *Heterologous Gene Expression in Yeast*, *Methods in Enzymology*, Eds. Berger & Kimmel, Acad. Press, N.Y., Vol. 152, pp. 673-684; and *The Molecular Biology of the Yeast Saccharomyces*, 1982, Eds. Strathern et al., Cold Spring Harbor Press, Vols. I and II.

[0168] In an insect system, *Autographa californica* nuclear polyhidrosis virus (AcNPV) a baculovirus, can be used as a vector to express PHOX2B in *Spodoptera frugiperda* cells. The PHOX2B gene sequences may be cloned into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). These recombinant viruses are then used to infect host cells in which the inserted DNA is expressed. (See e.g., Smith et al., 1983, *J Virol* 46:584; Smith, U.S. Pat. No. 4,215,051.)

[0169] Any of the cloning and expression vectors described herein may be synthesized and assembled from known DNA sequences by well known techniques in the art. The regulatory regions and enhancer elements can be of a variety of origins, both natural and synthetic. Some vectors and host cells may be obtained commercially. Non-limiting examples of useful vectors are described in Appendix 5 of *Current Protocols in Molecular Biology*, 1988, ed. Ausubel et al., Greene Publish. Assoc. & Wiley Interscience, which is incorporated herein by reference; and the catalogs of

commercial suppliers such as Clontech Laboratories, Stratagene Inc., and Invitrogen, Inc.

[0170] The resulting recombinant PHOX2B can be delivered to the cells in a patient by various methods known in the art.

5.7. Expression of PHOX2B

[0171] Expression constructs containing cloned nucleotide sequence encoding PHOX2B can be introduced into the host cell by a variety of techniques known in the art, including but not limited to, for prokaryotic cells, bacterial transformation (Hanahan, 1985, in *DNA Cloning, A Practical Approach*, 1:109-136), and for eukaryotic cells, calcium phosphate mediated transfection (Wigler et al., 1977, *Cell* 11:223-232), liposome-mediated transfection (Schaefer-Ridder et al., 1982, *Science* 215:166-168), electroporation (Wolff et al., 1987, *Proc Natl Acad Sci* 84:3344), and microinjection (Capechi, 1980, *Cell* 22:479-488).

[0172] For long term, high yield production of properly processed PHOX2B, expression in mammalian cells is preferred. Cell lines that stably express PHOX2B may be engineered by using a vector that contains a selectable marker. By way of example but not limitation, following the introduction of the expression constructs, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the expression construct confers resistance to the selection and optimally allows cells to stably integrate the expression construct into their chromosomes and to grow in culture and to be expanded into cell lines. Such cells can be cultured for a long period of time while PHOX2B is expressed continuously.

[0173] The recombinant cells may be cultured under standard conditions of temperature, incubation time, optical density, and media composition.

[0174] In an embodiment where the expression construct comprising sequences encoding a PHOX2B is used in gene therapy, the PHOX2B gene sequence is introduced into cancer cells in vivo. Such introduction can be carried out by any method known in the art, such as but not limited to transfection, transduction, microinjection, infection with a viral or bacteriophage vector containing the PHOX2B gene sequences, liposome-mediated gene transfer, microcell-mediated gene transfer, etc. Numerous techniques are known in the art for the introduction of foreign genes into cells (see e.g., Loeffler and Behr, 1993, *Meth. Enzymol.* 217:599-618; Cohen et al., 1993, *Meth. Enzymol.* 217:618-644; Cline, 1985, *Pharmac. Ther.* 29:69-92) may be used in accordance with the present invention, provided that the physiological functions of the recipient are not disrupted. The technique should provide for the stable transfer of the PHOX2B gene sequence to the cell, so that the sequence is expressible by the cell and preferably heritable and expressible by its cell progeny.

[0175] In another embodiment, the expression characteristics of an endogenous gene (e.g., PHOX2B gene) within a cell, cell line or microorganism may be modified by inserting a DNA regulatory element heterologous to the endogenous gene of interest into the genome of a cell, stable cell line or cloned microorganism such that the inserted regulatory element is operatively linked with the endogenous gene

and controls, modulates or activates. For example, endogenous genes which are normally "transcriptionally silent", i.e., a PHOX2B gene which is normally not expressed, or are expressed only at very low levels in a cell line or microorganism, may be activated by inserting a regulatory element which is capable of promoting the expression of a normally expressed gene product in that cell line or microorganism. Alternatively, transcriptionally silent, endogenous PHOX2B gene may be activated by insertion of a promiscuous regulatory element that works across cell types.

[0176] A heterologous regulatory element may be inserted into a stable cell line or cloned microorganism, such that it is operatively linked with and activates expression of endogenous PHOX2B gene, using techniques, such as targeted homologous recombination, which are well known to those of skill in the art, and described e.g., in Chappel, U.S. Pat. No. 5,272,071; PCT publication No. WO 91/06667, published May 16, 1991; Skoultchi U.S. Pat. No. 5,981,214; Treco et al U.S. Pat. No. 5,968,502 and PCT publication No. WO 94/12650, published Jun. 9, 1994. Alternatively, non-targeted e.g., non-homologous recombination techniques which are well-known to those of skill in the art and described, e.g., in PCT publication No. WO 99/15650, published Apr. 1, 1999, may be used.

5.8. Antibodies to PHOX2B Gene

[0177] In another embodiment, the present invention relates to the uses of antibodies or fragments thereof capable of specifically recognizing one or more epitopes of the PHOX2B gene, mutant PHOX2B gene, PHOX2B gene products, mutant PHOX2B gene products, epitopes of conserved variants of the PHOX2B gene products, epitopes of mutant PHOX2B gene products, or peptide fragments of the PHOX2B gene products. In addition, antibodies are useful in a variety of ways in accordance with the present invention. These antibodies can be utilized for the diagnosis of HSCR and in certain applications, targeting of affected tissues. Such antibodies may include, but are not limited to, polyclonal antibodies, monoclonal antibodies (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, Fv fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above.

[0178] Such antibodies may be used, for example, in the detection of a PHOX2B gene product in a biological sample and may, therefore, be utilized as part of a diagnostic or prognostic technique whereby patients may be tested for abnormal levels of PHOX2B gene products, and/or for the presence of abnormal forms of the such gene products. Such antibodies may also be included as a reagent in a kit for use in a diagnostic or prognostic technique. Such antibodies may also be utilized in conjunction with, for example, compound screening schemes, as described, below, for the evaluation of the effect of test compounds on PHOX2B gene product levels and/or activity. Antibodies to PHOX2B gene product may be used in a method for the inhibition of abnormal PHOX2B gene product activity. Thus, such antibodies may, therefore, be utilized as part of treatment methods for HSCR. Additionally, such antibodies can be used in conjunction with the gene therapy techniques described, below, to, for example, evaluate the normal and/or engineered PHOX2B-expressing cells prior to their introduction into the patient.

[0179] Described herein are methods for the production of antibodies of such antibodies or fragments thereof. Any of such antibodies or fragments thereof may be produced by standard immunological methods or by recombinant expression of nucleic acid molecules encoding the antibody or fragments thereof in an appropriate host organism.

[0180] Potentially immunogenic domains of the PHOX2B protein or mutant PHOX2B protein are predicted from hydropathy and surface probability profiles. Then oligopeptides which span the predicted immunogenic sites are chemically synthesized. These oligopeptides can be designed to contain the specific mutant amino acids to allow the detection of and discrimination between the mutant versus wild-type gene products. In specific embodiment, mutant PHOX2B protein comprising the amino acid sequence of SEQ ID NO: 6 may be used to raise antibodies for the particular portion of the protein that comprises the deletion of five alanine residues.

[0181] For the production of antibodies against a PHOX2B gene product, various host animals may be immunized by injection with a PHOX2B gene product, or a fragment thereof. Fragments of PHOX2B can be synthesized as antigenic peptides in accordance with the known amino acid sequence of PHOX2B. Such host animals may include but are not limited to rabbits, mice, and rats, to name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including but not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*.

[0182] Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of animals immunized with an antigen, such as a PHOX2B gene product, or an antigenic functional derivative thereof. For the production of polyclonal antibodies, host animals such as those described above, may be immunized by injection with PHOX2B gene product supplemented with adjuvants as also described above.

[0183] Monoclonal antibodies, which are homogeneous populations of antibodies to a particular antigen, may be obtained by any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique of Kohler and Milstein, (1975, Nature 256:495-497; and U.S. Pat. No. 4,376,110), the human B-cell hybridoma technique (Kosbor et al., 1983, Immunology Today 4:72; Cole et al., 1983, Proc. Natl. Acad. Sci. USA 80:2026-2030), and the EBV-hybridoma technique (Cole et al., 1985, Monoclonal Antibodies And Cancer Therapy, Alan R. Liss, Inc., pp. 77-96). Such antibodies may be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof. The hybridoma producing the mAb of this invention may be cultivated in vitro or in vivo. Production of high titers of mAbs in vivo makes this the presently preferred method of production.

[0184] In addition, techniques developed for the production of "chimeric antibodies" (Morrison et al., 1984, Proc. Natl. Acad. Sci., 81:6851-6855; Neuberger et al., 1984,

Nature, 312:604-608; Takeda et al., 1985, Nature, 314:452-454) by splicing the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region, e.g., humanized antibodies.

[0185] Alternatively, techniques described for the production of single chain antibodies (U.S. Pat. No. 4,946,778; Bird, 1988, Science 242:423-426; Huston et al., 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; and Ward et al., 1989, Nature 334:544-546) can be adapted to produce single chain antibodies against PHOX2B gene products. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide. Techniques for the assembly of functional Fv fragments in *E. coli* may also be used (Skerra et al., 1988, Science 242:1038-1041).

[0186] Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, such fragments include but are not limited to: the F(ab')₂ fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed (Huse et al., 1989, Science, 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

15 5.9. Methods of Treatment, Prophylaxis, Management or amelioration of Symptoms Associated with HSCR

[0187] The invention provides a method of diagnosis and therapy comprising diagnosing patients at increased risk of HSCR and treating, prophylactic, management, or ameliorating HSCR in an individual having such increased risk by methods of the invention. Management refers to the beneficial effects that a subject derives from a prophylactic or therapeutic agent, which does not result in a cure of the disease. In certain embodiments, a subject is administered one or more prophylactic or therapeutic agents to "manage" a disorder so as to prevent the progression or worsening of the disorder. Known therapies for HSCR can be effective in halting advancement of the disease, or at least slowing the advancement. It is thus an advantage of the invention for early identification of patients at increased risk of HSCR so that preventive therapy can be started as soon as possible, optimizing any intervention potential for affecting outcome. The decision of a physician on how and whether to initiate therapy in anticipation of the disease can be taken with increased confidence. Diseases or disorders that are related to decreased level or activity of PHOX2B may be treated by administering a therapeutic of the invention that intend to replace or act in place of the functionally deficient endogenous PHOX2B gene. An increased level or activity of PHOX2B may be treated by administering a therapeutic of the invention that inhibit the level or activity of endogenous PHOX2B. A variety of suitable treatments of patients at increased risk of HSCR are described in the art. These treatments include, but are not limited to, surgical removal of aganglionic bowel, anastomosis of ano-rectum with gan-

glionic bowel, colostomy, gene therapy, regeneration of enteric nerves by administration of growth factors, and intestinal or neural cell transplant. Other treatments will be known to persons of skill in the art.

5.10. Therapeutics of the Invention

[0188] The invention provides for treatment, prophylaxis, management or amelioration of one or more symptoms associated with the disease, disorder, or infection by administration of therapeutic compound (termed herein "therapeutics") of the invention. Compounds such as those identified through assays described infra, in Section 5.11, which modulate PHOX2B activity can be used in accordance with the invention to treat HSCR. Such molecules can include, but are not limited to peptides, including soluble peptides, and small organic or inorganic molecules, and can be referred to as PHOX2B agonists or antagonists. Dominant negative mutants of PHOX2B and PHOX2B-repressor fusion proteins are also compounds that interfere with the interaction of PHOX2B with intracellular macromolecules may also be used. Techniques for the determination of effective doses and administration of such compounds are described, in Sections 5.13 and 5.14.

[0189] Any technique which serves to selectively administer nucleic acid molecules to a cell population of interest can be used, for example, by using a delivery complex. Such a delivery complex can comprise an appropriate nucleic acid molecule and a targeting means. Such targeting means can comprise, for example, sterols, lipids, viruses or target cell specific binding agents. Viral vectors that can be used with recombinant viruses include, but are not limited to adenovirus, adeno-associated virus, herpes simplex virus, vaccinia virus, and retrovirus vectors, in addition to other particles that introduce DNA into cells, such as liposomes.

[0190] The therapeutics can be utilized for the prevention of a variety of disorder or diseases related to neuron development, in particular, HSCR, e.g., in individuals who are predisposed as a result of familial history or in individuals with an enhanced risk due to environmental factors.

[0191] Described below are methods and compositions for treating HSCR using the PHOX2B gene or gene product as a therapeutic target. The outcome of a treatment is to at least produce in a treated subject a healthful benefit.

[0192] All such methods comprise methods which modulate PHOX2b gene activity and/or expression which in turn modulate the phenotype of the treated subject.

[0193] The invention also encompasses methods to increase the transcriptional activity of PHOX2B gene, and methods to increase the expression of genes that are under the transcriptional control of PHOX2B proteins. The invention also encompasses methods to increase the translation of PHOX2B transcripts and increasing the stability of the PHOX2B protein. Furthermore, the invention encompasses decreasing the expression of genes that are negatively regulated by PHOX2B. The increase of the transcriptional activity of the PHOX2B gene may provide a novel therapeutic approach for HSCR and other disease conditions that are associated with reduced transcriptional activity of the PHOX2B protein, or underexpression of the protein.

[0194] PHOX2B are responsible for various processes related to neural development, including but not limited to,

promoting neuron growth and development, development of autonomic crest derivatives, formation, migration and differentiation of neural crest cells into autonomic neurons, coordination of quantitative and qualitative aspects of neurogenesis, generating neurons of the correct phenotype and in proper numbers at the appropriate times and locations, promoting cell cycle exit, cell migration, and initiates cell differentiation. Hence, PHOX2b gene product may be used as therapeutics for the treatment, prophylaxis, management, or amelioration of symptoms associated with any disorders or diseases that are related to PHOX2b aberrations. In a specific embodiment, the therapeutics of the present invention may be used to treat HSCR. Accordingly, the present invention provides therapeutic compositions comprising the nucleotide sequences of PHOX2B nucleic acid molecules as described in Sections 5.1 and 5.5. In specific embodiments, the PHOX2B nucleic acid molecules comprise the open reading frame depicted in FIG. 1 (SEQ ID NO: 1), or nucleotide sequences encoding a PHOX2B protein. The invention provides therapeutic compositions comprising purified nucleic acids consisting of at least 10 contiguous nucleotides (i.e., a hybridizable portion) of a PHOX2B sequence; in other embodiments, the nucleic acids consist of at least 15 (continuous) nucleotides, 25 nucleotides, 50 nucleotides, 100 nucleotides, 150 nucleotides, 200 nucleotides, or 250 nucleotides of a PHOX2B sequence. In another embodiment, the nucleic acids are smaller than 35, 200 or 250 nucleotides in length. Nucleic acids can be single or double stranded. The invention also provides therapeutic compositions comprising nucleic acids hybridizable to or complementary to the foregoing sequences. In specific aspects, the nucleic acids comprise a sequence complementary to at least 10, 25, 50, 100, 200, or 250 nucleotides of a PHOX2B gene. In a specific embodiment, a nucleic acid which is hybridizable to a PHOX2B nucleic acid (e.g., having sequence SEQ ID NO: 1, 3, 4, or 5), or to a nucleic acid encoding a PHOX2B derivative, under conditions of moderate stringency is provided.

[0195] Therapeutic compositions comprising nucleic acids encoding derivatives and analogs of PHOX2B proteins, and PHOX2B antisense nucleic acids are additionally provided. As is readily apparent, as used herein, a "nucleic acid encoding a fragment or portion of a PHOX2B protein" shall be construed as referring to a nucleic acid encoding only the recited fragment or portion of the PHOX2B protein and not the other contiguous portions of the PHOX2B protein as a continuous sequence.

[0196] Therapeutic compositions comprising fragments of PHOX2B nucleic acids comprising regions conserved between other PHOX2B nucleic acids, of the same or different species, are also provided. Therapeutic compositions comprising nucleic acids encoding one or more domains of PHOX2B are provided.

[0197] Successful treatment of HSCR may be brought about by techniques which serve to decrease PHOX2B activity. Activity can be decreased by, for example, directly decreasing PHOX2B gene product activity and/or by decreasing the level of PHOX2B gene expression.

[0198] For example, compounds such as those identified through assays described, above, which decrease PHOX2B activity can be used in accordance with the invention to treat HSCR.

[0199] Further, antisense and ribozyme molecules which inhibit PHOX2B gene expression can also be used in accordance with the invention to reduce the level of PHOX2B gene expression, thus effectively reducing the level of PHOX2B gene product present, thereby decreasing the level of PHOX2B activity. Still further, triple helix molecules can be utilized in reducing the level of PHOX2B gene activity. Such molecules can be designed to reduce or inhibit either wild type, or if appropriate, mutant target gene activity. Techniques for the production and use of such molecules are well known to those of skill in the art.

5.10.1. Antisense Molecules

[0200] The use of antisense molecules as inhibitors of gene expression is a specific, genetically based therapeutic approach (for a review, see Stein, in Ch. 69, Section 5 "Cancer: Principle and Practice of Oncology", 4th ed., ed. by De Vita et al., J. B. Lippincott, Philadelphia 1993). The present invention provides the therapeutic or prophylactic use of nucleic acids of at least six nucleotides that are antisense to a gene or cDNA encoding PHOX2B or a portion thereof. An "antisense" PHOX2B nucleic acid as used herein refers to a nucleic acid capable of hybridizing to a portion of a PHOX2B RNA (preferably mRNA) by virtue of some sequence complementarity. The invention further provides pharmaceutical compositions comprising an effective amount of the PHOX2B antisense nucleic acids of the invention in a pharmaceutically acceptable carrier, as described infra.

[0201] In another embodiment, the invention is directed to methods for inhibiting the expression of a PHOX2B nucleic acid sequence in a mammalian cell in vitro or in vivo comprising providing the cell with an effective amount of a composition comprising an PHOX2B antisense nucleic acid of the invention.

[0202] The antisense nucleic acid of the invention may be complementary to a coding and/or noncoding region of a PHOX2B mRNA. The antisense molecules will bind to the complementary PHOX2B gene mRNA transcripts and reduce or prevent translation. Absolute complementarity, although preferred, is not required. A sequence "complementary" to a portion of an RNA, as referred to herein, means a sequence having sufficient complementarity to be able to hybridize with the RNA, forming a stable duplex; in the case of double-stranded antisense nucleic acids, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid. Generally, the longer the hybridizing nucleic acid, the more base mismatches with an RNA it may contain and still form a stable duplex (or triplex, as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

[0203] Nucleic acid molecules that are complementary to the 5' end of the message, e.g., the 5' untranslated sequence up to and including the AUG initiation codon, should work most efficiently at inhibiting translation. However, sequences complementary to the 3' untranslated sequences of mRNAs have recently shown to be effective at inhibiting translation of mRNAs as well. See generally, Wagner, R., 1994, Nature 372:333-335. Thus, oligonucleotides comple-

mentary to either the 5'- or 3'-non-translated, non-coding regions of the PHOX2B gene, as shown, for example, in FIG. 1, could be used in an antisense approach to inhibit translation of endogenous PHOX2B gene mRNA.

[0204] Nucleic acid molecules complementary to the 5' untranslated region of the mRNA should include the complement of the AUG start codon. Antisense oligonucleotides complementary to mRNA coding regions are less efficient inhibitors of translation but could be used in accordance with the invention. Whether designed to hybridize to the 5', 3'- or coding region of target or pathway gene mRNA, antisense nucleic acids should be at least six nucleotides in length, and are preferably oligonucleotides ranging from 6 to about 50 nucleotides in length. In specific aspects the oligonucleotide is at least 10 nucleotides, at least 17 nucleotides, at least 25 nucleotides or at least 50 nucleotides.

[0205] Regardless of the choice of target sequence, it is preferred that in vitro studies are first performed to quantify the ability of the antisense oligonucleotide to inhibit gene expression. It is preferred that these studies utilize controls that distinguish between antisense gene inhibition and nonspecific biological effects of oligonucleotides. It is also preferred that these studies compare levels of the target RNA or protein with that of an internal control RNA or protein. Additionally, it is envisioned that results obtained using the antisense oligonucleotide are compared with those obtained using a control oligonucleotide. It is preferred that the control oligonucleotide is of approximately the same length as the test oligonucleotide and that the nucleotide sequence of the oligonucleotide differs from the antisense sequence no more than is necessary to prevent specific hybridization to the target sequence.

[0206] The oligonucleotides can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, hybridization, etc. The oligonucleotide may include other appended groups such as peptides (e.g., for targeting host cell receptors in vivo), or agents facilitating transport across the cell membrane (see, e.g., Letsinger et al., 1989, Proc. Natl. Acad. Sci. U.S.A. 86:6553-6556; Lemaitre et al., 1987, Proc. Natl. Acad. Sci. 84:648-652; PCT Publication No. WO88/09810, published Dec. 15, 1988) or the blood-brain barrier (see, e.g., PCT Publication No. WO89/10134, published Apr. 25, 1988), hybridization-triggered cleavage agents. (See, e.g., Krol et al., 1988, BioTechniques 6:958-976) or intercalating agents. (See, e.g., Zon, 1988, Pharm. Res. 5:539-549). To this end, the oligonucleotide may be conjugated to another molecule, e.g., a peptide, hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

[0207] The antisense oligonucleotide may comprise at least one modified base moiety which is selected from the group including but not limited to 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine,

N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarbonylmethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

[0208] The antisense oligonucleotide may also comprise at least one modified sugar moiety selected from the group including but not limited to arabinose, 2-fluoroarabinose, xylulose, and hexose.

[0209] In yet another embodiment, the antisense oligonucleotide comprises at least one modified phosphate backbone selected from the group consisting of a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

[0210] In yet another embodiment, the antisense oligonucleotide is an α -anomeric oligonucleotide. An α -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gautier et al., 1987, Nucl. Acids Res. 15:6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue et al., 1987, Nucl. Acids Res. 15:6131-6148), or a chimeric RNA-DNA analogue (Inoue et al., 1987, FEBS Lett. 215:327-330).

[0211] Oligonucleotides of the invention may be synthesized by standard methods known in the art, e.g. by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides may be synthesized by the method of Stein et al. (1988, Nucl. Acids Res. 16:3209), methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., 1988, Proc. Natl. Acad. Sci. U.S.A. 85:7448-7451), etc.

[0212] While antisense nucleotides complementary to the PHOX2B coding region could be used, those complementary to the transcribed untranslated region are most preferred.

[0213] The antisense molecules should be delivered to cells which express the PHOX2B gene in vivo. A number of methods have been developed for delivering antisense DNA or RNA to cells; eg., antisense molecules can be injected directly into the tissue site, or modified antisense molecules, designed to target the desired cells (e.g., antisense linked to peptides or antibodies that specifically bind receptors or antigens expressed on the target cell surface) can be administered systemically.

[0214] However, it is often difficult to achieve intracellular concentrations of the antisense sufficient to suppress translation of endogenous mRNAs. Therefore a preferred approach utilizes a recombinant DNA construct in which the antisense oligonucleotide is placed under the control of a strong pol III or pol II promoter. The use of such a construct to transfect target cells in the patient will result in the transcription of sufficient amounts of single stranded RNAs that will form complementary base pairs with the endog-

enous PHOX2B gene transcripts and thereby prevent translation of the PHOX2B gene mRNA. For example, a vector can be introduced in vivo such that it is taken up by a cell and directs the transcription of an antisense RNA. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA. Such vectors can be constructed by recombinant DNA technology methods standard in the art, and described in Section 5.6. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in mammalian cells. Expression of the sequence encoding the antisense RNA can be by any promoter known in the art to act in mammalian, preferably human cells. Such promoters can be inducible or constitutive. Such promoters include but are not limited to: the SV40 early promoter region (Bernoist and Chambon, 1981, *Nature* 290:304-310), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., 1980, *Cell* 22:787-797), the herpes thymidine kinase promoter (Wagner et al., 1981, *Proc. Natl. Acad. Sci. U.S.A.* 78:1441-1445), the regulatory sequences of the metallothionein gene (Brinster et al., 1982, *Nature* 296:39-42), etc. Any type of plasmid, cosmid, YAC or viral vector can be used to prepare the recombinant DNA construct which can be introduced directly into the tissue site. Alternatively, viral vectors can be used which selectively infect the desired tissue.

5.10.2. Ribozyme Molecules

[0215] Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA (For a review see, for example Rossi, J., 1994, *Current Biology* 4:469-471). The mechanism of ribozyme action involves sequence specific hybridization of the ribozyme molecule to complementary target RNA, followed by an endonucleolytic cleavage. The composition of ribozyme molecules must include one or more sequences complementary to the target gene mRNA, and must include the well known catalytic sequence responsible for mRNA cleavage. For this sequence, see U.S. Pat. No. 5,093,246, which is incorporated by reference herein in its entirety. As such, within the scope of the invention are engineered hammerhead motif ribozyme molecules that specifically and efficiently catalyze endonucleolytic cleavage of RNA sequences encoding target gene proteins.

[0216] Ribozyme molecules designed to catalytically cleave PHOX2B gene mRNA transcripts can also be used to prevent translation of PHOX2B gene mRNA and expression of target or pathway gene. (See, eg., PCT International Publication WO90/11364, published Oct. 4, 1990; Sarver et al., 1990, *Science* 247:1222-1225). While ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy PHOX2B gene mRNAs, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole requirement is that the target mRNA have the following sequence of two bases: 5'-UG-3'. The construction and production of hammerhead ribozymes is well known in the art and is described more fully in Haseloff and Gerlach, 1988, *Nature*, 334:585-591. Preferably the ribozyme is engineered so that the cleavage recognition site is located near the 5' end of the PHOX2B gene mRNA; i.e., to increase efficiency and minimize the intracellular accumulation of non-functional mRNA transcripts.

[0217] The ribozymes of the present invention also include RNA endoribonucleases (hereinafter "Cech-type ribozymes") such as the one which occurs naturally in *Tetrahymena Thermophila* (known as the IVS, or L-19 IVS RNA) and which has been extensively described by Thomas Cech and collaborators (Zaug, et al., 1984, *Science*, 224:574-578; Zaug and Cech, 1986, *Science*, 231:470-475; Zaug, et al., 1986, *Nature*, 324:429-433; published International patent application No. WO 88/04300 by University Patents Inc.; Been and Cech, 1986, *Cell*, 47:207-216). The Cech-type ribozymes have an eight base pair active site which hybridizes to a target RNA sequence whereafter cleavage of the target RNA takes place. The invention encompasses those Cech-type ribozymes which target eight base-pair active site sequences that are present in an PHOX2B gene.

[0218] As in the antisense approach, the ribozymes can be composed of modified oligonucleotides (eg. for improved stability, targeting, etc.) and should be delivered to cells which express the PHOX2B gene in vivo. A preferred method of delivery involves using a DNA construct "encoding" the ribozyme under the control of a strong constitutive pol III or pol II promoter, so that transfected cells will produce sufficient quantities of the ribozyme to destroy endogenous PHOX2B gene messages and inhibit translation. Because ribozymes unlike antisense molecules, are catalytic, a lower intracellular concentration is required for efficiency.

[0219] Anti-sense RNA and DNA, ribozyme, and triple helix molecules of the invention can be prepared by any method known in the art for the synthesis of DNA and RNA molecules. These include techniques for chemically synthesizing oligodeoxyribonucleotides and oligoribonucleotides well known in the art such as for example solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules can be generated by in vitro and in vivo transcription of DNA sequences encoding the antisense RNA molecule. Such DNA sequences can be incorporated into a wide variety of vectors which incorporate suitable RNA polymerase promoters such as the T7 or SP6 polymerase promoters. Alternatively, antisense cDNA constructs that synthesize antisense RNA constitutively or inducibly, depending on the promoter used, can be introduced stably into cell lines.

[0220] Various well-known modifications to the DNA molecules can be introduced as a means of increasing intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences of ribo- or deoxy-nucleotides to the 5' and/or 3' ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the oligodeoxyribonucleotide backbone.

5.10.3. Therapeutic Antibodies

[0221] Antibodies exhibiting capability to downregulate PHOX2B gene product activity can be utilized to treat HSCR. Such antibodies can be generated using standard techniques described in Section 5.8, above, against full length wild type or mutant PHOX2B proteins, or against peptides corresponding to portions of the proteins. The antibodies include but are not limited to polyclonal, monoclonal, Fab fragments, single chain antibodies, chimeric antibodies, and the like.

[0222] Because PHOX2B is an intracellular protein, it is preferred that internalizing antibodies be used. However, lipofectin or liposomes can be used to deliver the antibody or a fragment of the Fab region which binds to the PHOX2B gene product epitope into cells. Where fragments of the antibody are used, the smallest inhibitory fragment which binds to the PHOX2B protein's binding domain is preferred. For example, peptides having an amino acid sequence corresponding to the domain of the variable region of the antibody that binds to the PHOX2B protein can be used. Such peptides can be synthesized chemically or produced via recombinant DNA technology using methods well known in the art (e.g., see Creighton, 1983; and Sambrook et al., 1989). Alternatively, single chain antibodies, such as neutralizing antibodies, which bind to intracellular epitopes can also be administered. Such single chain antibodies can be administered, for example, by expressing nucleotide sequences encoding single-chain antibodies within the target cell population by utilizing, for example, techniques such as those described in Marasco et al. (Marasco, W. et al., 1993, Proc. Natl. Acad. Sci. USA 90:7889-7893).

5.11. Screening Assays for Compounds that Modulate PHOX2B Activity

[0223] The present invention provides a drug target, PHOX2B gene, for the treatment, prophylaxis, amelioration, and management of HSCR. Hence, a therapeutics of the invention may be a compound that modulates the PHOX2B gene. Such compound may bind or interact with the PHOX2B gene or gene product. Accordingly, provided herein are screening assays for compounds that may, through its interaction with the PHOX2B gene or PHOX2B gene product, affect the onset, or progression of HSCR.

[0224] The following assays are designed to identify: (i) compounds that bind to PHOX2B gene products, including mammalian and non-mammalian homologs of PHOX2B; (ii) compounds that bind to other intracellular proteins and/or segments of nucleic acid that interact with a PHOX2B gene product, including mammalian and non-mammalian homologs of PHOX2B; (iii) compounds that interfere with the interaction of the PHOX2B gene product, including mammalian and non-mammalian homologs of PHOX2B, with other intracellular proteins and/or segments of nucleic acid; and (iv) compounds that modulate the activity of PHOX2B gene (i.e., modulate the level of PHOX2B gene expression and/or modulate the level of PHOX2B gene product activity).

[0225] Assays may additionally be utilized which identify compounds which bind to PHOX2B gene regulatory sequences (e.g., promoter sequences). See e.g., Platt, 1994, J. Biol. Chem. 269:28558-28562, which is incorporated herein by reference in its entirety. Also provided is a method for identifying compounds that modulate PHOX2B gene expression, comprising: (a) contacting a test compound with a cell or cell lysate containing a reporter gene operatively associated with a PHOX2B gene regulatory element; and (b) detecting expression of the reporter gene product. Also provided is another method for identifying compounds that modulate PHOX2B gene expression comprising: (a) contacting a test compound with a cell or cell lysate containing PHOX2B transcripts; and (b) detecting the translation of the PHOX2B transcript. Any reporter gene known in the art can

be used, such as but limited to, green fluorescent protein, β -galactosidase, alkaline phosphatase, chloramphenicol acetyltransferase, etc.

5.11.1. In Vitro Screening Assays for Compounds that Bind to the PHOX2B Gene Product

[0226] In vitro systems may be designed to identify compounds capable of interacting with, e.g., binding to, the PHOX2B gene products of the invention and homologs of PHOX2B. Compounds identified may be useful, for example, in modulating the activity of wild type and/or mutant PHOX2B gene products, may be useful in elaborating the biological function of the PHOX2B gene product, may be utilized in screens for identifying compounds that disrupt normal PHOX2B gene product interactions, or may in themselves disrupt such interactions.

[0227] The principle of the assays used to identify compounds that interact with the PHOX2B gene product involves preparing a reaction mixture of the PHOX2B gene product, or fragments thereof and the test compound under conditions and for a time sufficient to allow the two components to interact with, e.g., bind to, thus forming a complex, which can represent a transient complex, which can be removed and/or detected in the reaction mixture. These assays can be conducted in a variety of ways. For example, one method to conduct such an assay would involve anchoring PHOX2B gene product or the test substance onto a solid phase and detecting PHOX2B gene product/test compound complexes anchored on the solid phase at the end of the reaction. In one embodiment of such a method, the PHOX2B gene product or fragment thereof may be anchored onto a solid surface, and the test compound, which is not anchored, may be labeled, either directly or indirectly.

[0228] In practice, microtitre plates may conveniently be utilized as the solid phase. The anchored component may be immobilized by non-covalent or covalent attachments. Non-covalent attachment may be accomplished by simply coating the solid surface with a solution of the protein and drying. Alternatively, an immobilized antibody, preferably a monoclonal antibody, specific for the protein to be immobilized may be used to anchor the protein to the solid surface. The surfaces may be prepared in advance and stored.

[0229] In order to conduct the assay, the nonimmobilized component is added to the coated surface containing the anchored component. After the reaction is complete, unreacted components are removed (e.g., by washing) under conditions such that any complexes formed will remain immobilized on the solid surface. The detection of complexes anchored on the solid surface can be accomplished in a number of ways. Where the previously nonimmobilized component is pre-labeled, the detection of label immobilized on the surface indicates that complexes were formed. Where the previously nonimmobilized component is not pre-labeled, an indirect label can be used to detect complexes anchored on the surface; e.g., using a labeled antibody specific for the previously nonimmobilized component (the antibody, in turn, may be directly labeled or indirectly labeled with a labeled anti-Ig antibody).

[0230] Alternatively, a reaction can be conducted in a liquid phase, the reaction products separated from unreacted components, and complexes detected; e.g., using an immo-

bilized antibody specific for PHOX2B gene product or the test compound to anchor any complexes formed in solution, and a labeled antibody specific for the other component of the possible complex to detect anchored complexes.

5.11.2. Assays for Intracellular Proteins that Interact with the PHOX2B Gene Product

[0231] Any method suitable for detecting protein-protein interactions or protein-nucleic acid interactions may be employed for identifying PHOX2B protein-intracellular protein interactions, especially interactions mediated by the PHOX2B domain.

[0232] Among the traditional methods which may be employed are co-immunoprecipitation, crosslinking and copurification through gradients or chromatographic columns. Surface display library and yeast-based two-hybrid system can also be utilized to isolate the gene encoding such PHOX2B-binding proteins.

[0233] These methods allow the identification of molecules, including intracellular proteins, that interact with PHOX2B gene products. Once isolated, such a protein can be sequenced using techniques well-known to those of skill in the art, such as by Edman degradation (see, e.g., Creighton, 1983, "Proteins: Structures and Molecular Principles," W. H. Freeman & Co., N.Y., pp.34-49). The amino acid sequence obtained may be used as a guide for the generation of oligonucleotide mixtures that can be used to screen for gene sequences encoding such proteins. Screening may be accomplished, for example, by standard hybridization or PCR techniques. Techniques for the generation of oligonucleotide mixtures and the screening are well-known. (See, e.g., Ausubel, supra, and 1990, "PCR Protocols: A Guide to Methods and Applications," Innis, et al., eds. Academic Press, Inc., New York).

[0234] Another method that detects protein interactions *in vivo* is the two-hybrid system, which is described here for illustration only and not by way of limitation. One example of this approach has been described (Chien, et al., 1991, Proc. Natl. Acad. Sci. USA, 88:9578-9582) and a kit is commercially available from Clontech (Palo Alto, Calif.).

[0235] The two-hybrid system or related methodologies may be used to screen activation domain libraries for proteins that interact with a "bait" gene product. By way of example, and not by way of limitation, PHOX2B gene products may be used as the bait gene product. Total genomic or cDNA sequences are fused to the DNA encoding an activation domain. This library and a plasmid encoding a hybrid of a bait PHOX2B gene product fused to the DNA-binding domain are co-transformed into a yeast reporter strain, and the resulting transformants are screened for those that express the reporter gene. For example, a bait PHOX2B gene sequence, such as the open reading frame of the PHOX2B gene, can be cloned into a vector such that it is translationally fused to the DNA encoding the DNA-binding domain of the GAL4 protein. These colonies are purified and the library plasmids responsible for reporter gene expression are isolated. DNA sequencing is then used to identify the proteins encoded by the library plasmids.

[0236] A cDNA library of the cell line from which proteins that interact with bait PHOX2B gene product are to be detected can be made using methods routinely practiced in

the art. According to the particular system described herein, for example, the cDNA fragments can be inserted into a vector such that they are translationally fused to the transcriptional activation domain of GAL4. Such a library can be co-transformed along with the bait PHOX2B gene-GAL4 fusion plasmid into a yeast strain that contains a lacZ gene driven by a promoter that contains GAL4 activation sequence. A cDNA encoded protein, fused to a GAL4 transcriptional activation domain that interacts with bait PHOX2B gene product will reconstitute an active GAL4 protein and thereby drive expression of the HIS3 gene. Colonies that express HIS3 can be detected by their growth on petri dishes containing semi-solid agar based media lacking histidine. The cDNA can then be purified from these strains, and used to produce and isolate the bait PHOX2B gene product-interacting protein using techniques routinely practiced in the art.

[0237] Additionally, methods may be employed which result in the identification of nucleic acids which interact with the PHOX2B protein, i.e., PHOX2B target sites. These methods include, for example, probing gene libraries with labeled PHOX2B protein or fragments thereof (e.g., PHOX2B domain), using PHOX2B protein or fragments thereof in a manner similar to the well-known technique of antibody probing of λ gt11 libraries. Such methods can also be adapted to monitor the interactions of dominant negative mutants of PHOX2B and PHOX2B target sites. Furthermore, novel PHOX2B target sequences can be isolated by the RNA differential display technique and whole genome PCR. See Robinson et al., 1997, Proc. Natl. Acad. Sci. USA, 94:7170-7175.

5.11.3. Assays for Compounds that Interfere with PHOX2B Gene Product/Intracellular Macromolecular Interaction

[0238] The PHOX2B gene products of the invention, fragments thereof, and homologs of PHOX2B may, *in vivo*, interact with one or more intracellular macromolecules, such as proteins and nucleic acid molecules. Such macromolecules may include, but are not limited to DNA, RNA (including polyadenylated (poly(A)) RNA and RNA with the 5' cap structure) and those proteins identified via methods such as those described, above, in Section 5.8.2. For purposes of this discussion, such intracellular macromolecules are referred to herein as "interacting partners". Compounds that disrupt PHOX2B interactions in this way may be useful in regulating the activity of the PHOX2B gene product, including mutant PHOX2B gene products. Such compounds may include, but are not limited to molecules such as peptides, and the like, which would be capable of gaining access to the intracellular PHOX2B gene product.

[0239] The basic principle of the assay systems used to identify compounds that interfere with the interaction between the PHOX2B gene product and its intracellular interacting partner or partners involves preparing a reaction mixture containing the PHOX2B gene product, or fragments thereof, and the interacting partner under conditions and for a time sufficient to allow the two to interact and bind, thus forming a complex. In order to test a compound for inhibitory activity, the reaction mixture is prepared in the presence and absence of the test compound. The test compound may be initially included in the reaction mixture, or may be added at a time subsequent to the addition of PHOX2B gene

product and its intracellular interacting partner. Control reaction mixtures are incubated without the test compound or with a vehicle or carrier. The formation of any complexes between the PHOX2B gene product or fragments thereof and the intracellular interacting partner is then detected. The formation of a complex in the control reaction, but not in the reaction mixture containing the test compound, indicates that the compound interferes with the interaction of the PHOX2B gene protein and the interacting partner. Additionally, complex formation within reaction mixtures containing the test compound and normal PHOX2B gene protein may also be compared to complex formation within reaction mixtures containing the test compound and a mutant PHOX2B gene protein. This comparison may be important in those cases wherein it is desirable to identify compounds that disrupt interactions of mutant but not normal PHOX2B gene proteins.

5.11.4. Cell-Based Assays for Identification of Compounds that Modulate PHOX2B

[0240] Cell-based methods are presented herein which identify compounds capable of treating HSCR by modulating PHOX2B activity. Specifically, such assays identify compounds which affect PHOX2B-dependent processes, such as but not limited to promoting neuron growth and development, development of autonomic crest derivatives, formation, migration and differentiation of neural crest cells into autonomic neurons, coordination of quantitative and qualitative aspects of neurogenesis, generating neurons of the correct phenotype and in proper numbers at the appropriate times and locations, promoting cell cycle exit, cell migration, and initiates cell differentiation.

[0241] In another embodiment, the cell-based assays are based on expression of the PHOX2B gene product in a mammalian cell and measuring the PHOX2B-dependent process. Any mammalian cells that can express the PHOX2B gene and allow the functioning of the PHOX2B gene product can be used. Recombinant expression of the PHOX2B gene in these cells can be achieved by methods described above. In these assays, cells producing functional PHOX2B gene products are exposed to a test compound for an interval sufficient for the compound to modulate the activity of the PHOX2B gene product. The activity of PHOX2B gene product can be measured directly or indirectly through the detection or measurement of PHOX2B-dependent cellular processes such as, for example, the manifestation of a transformed phenotype. As a control, a cell not producing the PHOX2B gene product may be used for comparisons. Depending on the cellular process, any techniques known in the art may be applied to detect or measure it.

5.12. Gene Therapy

[0242] Gene therapy refers to treatment or prevention of a disease performed by the administration of a nucleic acid to a subject. In this embodiment of the invention, the therapeutic nucleic acid produces intracellularly a nucleic acid molecule of the invention that mediates a therapeutic effect by modulating PHOX2B expression. In an embodiment, nucleic acids comprising a sequence encoding a wild-type PHOX2B gene are administered to treat symptoms related to HSCR.

[0243] For general reviews of the methods of gene therapy, see Goldspiel et al., 1993, *Clinical Pharmacy*

12:488-505; Wu and Wu, 1991, *Biotherapy* 3:87-95; Tolsoshev, 1993, *Ann. Rev. Pharmacol. Toxicol.* 32:573-596; Mulligan, 1993, *Science* 260:926-932; and Morgan and Anderson, 1993, *Ann. Rev. Biochem.* 62:191-217; May, 1993, *TIBTECH* 11(5):155-215). Methods commonly known in the art of recombinant DNA technology which can be used are described in Ausubel et al. (eds.), 1993, *Current Protocols in Molecular Biology*, John Wiley & Sons, NY; Kriegler, 1990, *Gene Transfer and Expression, A Laboratory Manual*, Stockton Press, NY; and in Chapters 12 and 13, Dracopoli et al. (eds.), 1994, *Current Protocols in Human Genetics*, John Wiley & Sons, NY.

[0244] Delivery of the nucleic acid into a patient may be either direct, in which case the patient is directly exposed to the nucleic acid or nucleic acid-carrying vector or a delivery complex, or indirect, in which case, cells are first transfected with the nucleic acid in vitro, then transplanted into the patient. These two approaches are known, respectively, as in vivo or ex vivo gene therapy.

[0245] In a specific embodiment, the nucleic acid is directly administered in vivo, where it is expressed to produce the nucleic acid molecule of the invention or encoded functional PHOX2B gene product. This can be accomplished by any of numerous methods known in the art, e.g., by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, e.g., by infection using a defective or attenuated retroviral or other viral vector (see U.S. Pat. No. 4,980,286), or by direct injection of naked DNA, or by use of microparticle bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in biopolymers (e.g., poly- β -1- \rightarrow 4-N-acetylglucosamine polysaccharide; see U.S. Pat. No. 5,635,493), encapsulation in liposomes, microparticles, or microcapsules, or by administering it in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (see e.g., Wu and Wu, 1987, *J. Biol. Chem.* 262:4429-4432), etc. In another embodiment, a nucleic acid-ligand complex can be formed in which the ligand comprises a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted in vivo for cell specific uptake and expression, by targeting a specific receptor (see, e.g., PCT Publications WO 92/06180 dated Apr. 16, 1992 (Wu et al.); WO 92/22635 dated Dec. 23, 1992 (Wilson et al.); WO92/20316 dated Nov. 26, 1992 (Findeis et al.); WO93/14188 dated Jul. 22, 1993 (Clarke et al.); WO 93/20221 dated Oct. 14, 1993 (Young)). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination (Koller and Smithies, 1989, *Proc. Natl. Acad. Sci. USA* 86:8932-8935; Zijlstra et al., 1989, *Nature* 342:435-438).

[0246] In a specific embodiment, a viral vector that contains the PHOX2B nucleic acid molecule of the invention is used. For example, a retroviral vector can be used (see Miller et al., 1993, *Meth. Enzymol.* 217:581-599). These retroviral vectors have been modified to delete retroviral sequences that are not necessary for packaging of the viral genome and integration into host cell DNA. The nucleic acid

molecule of the invention to be used in gene therapy is cloned into the vector, which facilitates delivery of the gene into a patient.

[0247] The form and amount of therapeutic nucleic acid envisioned for use depends on the symptoms presented by the HSCR patient, desired effect, patient state, etc., and can be determined by one skilled in the art.

10 5.13. Pharmaceutical Preparations and Methods of Administration

[0248] The compounds and nucleic acid sequences described herein can be administered to a patient at therapeutically effective doses to treat or prevent cancer. A therapeutically effective dose refers to that amount of a compound sufficient to result in a healthful benefit in the treated subject. Formulations and methods of administration that can be employed when the therapeutic composition comprises a nucleic acid are described below.

[0249] Pharmaceutical compositions for use in accordance with the present invention can be formulated in conventional manner using one or more physiologically acceptable carriers or excipients.

[0250] Thus, the compounds and their physiologically acceptable salts and solvents can be formulated for administration by inhalation or insufflation (either through the mouth or the nose) or oral, buccal, parenteral or rectal administration.

[0251] For oral administration, the pharmaceutical compositions can take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g., pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g., lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g., magnesium stearate, talc or silica); disintegrants (e.g., potato starch or sodium starch glycolate); or wetting agents (e.g., sodium lauryl sulphate). The tablets can be coated by methods well known in the art. Liquid preparations for oral administration can take the form of, for example, solutions, syrups or suspensions, or they can be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations can be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (e.g., lecithin or acacia); non-aqueous vehicles (e.g., almond oil, oily esters, ethyl alcohol or fractionated vegetable oils); and preservatives (e.g., methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations can also contain buffer salts, flavoring, coloring and sweetening agents as appropriate.

[0252] Preparations for oral administration can be suitably formulated to give controlled release of the active compound.

[0253] For buccal administration the compositions can take the form of tablets or lozenges formulated in conventional manner.

[0254] For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from

pressurized packs or a nebulizer, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g. gelatin for use in an inhaler or insufflator can be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

[0255] The compounds can be formulated for parenteral administration (i.e., intravenous or intramuscular) by injection, via, for example, bolus injection or continuous infusion. Formulations for injection can be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions can take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and can contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient can be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

[0256] The compounds can also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

[0257] In addition to the formulations described previously, the compounds can also be formulated as a depot preparation. Such long acting formulations can be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds can be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

5.14. Characterization and Demonstration of Therapeutic or Prophylactic Utility

[0258] Toxicity and efficacy of the prophylactic and/or therapeutic protocols of the instant invention can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD₅₀/ED₅₀. Prophylactic and/or therapeutic agents that exhibit large therapeutic indices are preferred. While prophylactic and/or therapeutic agents that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such agents to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

[0259] The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage of the prophylactic and/or therapeutic agents for use in humans. The dosage of such agents lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any agent used in the method of the invention, the therapeutically effective dose

can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC_{50} (i.e., the concentration of the test compound that achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

[0260] The therapeutics used in accordance with the present invention also can be determined by using various experimental animal models.

[0261] The protocols and compositions of the invention are preferably tested *in vitro*, and then *in vivo*, for the desired therapeutic or prophylactic activity, prior to use in humans. For example, *in vitro* assays which can be used to determine whether administration of a specific therapeutic protocol is indicated, include *in vitro* cell culture assays in which a patient tissue sample is grown in culture, and exposed to or otherwise administered a protocol, and the effect of such protocol upon the tissue sample is observed. A lower level of proliferation or survival of the contacted cells indicates that the Therapeutic is effective to treat the condition in the patient. Alternatively, instead of culturing cells from a patient, Protocols may be screened using cells of a tumor or malignant cell line. Many assays standard in the art can be used to assess such survival and/or growth; for example, cell proliferation can be assayed by measuring 3H -thymidine incorporation, by direct cell count, by detecting changes in transcriptional activity of known genes such as proto-oncogenes (e.g., *fos*, *myc*) or cell cycle markers; cell viability can be assessed by trypan blue staining, differentiation can be assessed visually based on changes in morphology, etc.

[0262] Compounds for use in therapy can be tested in suitable animal model systems prior to testing in humans, including but not limited to in rats, mice, chicken, cows, monkeys, rabbits, etc.

[0263] Further, any assays known to those skilled in the art can be used to evaluate the prophylactic and/or therapeutic utility of the therapies disclosed herein for treatment or prevention of HSCR.

[0264] Efficacy in treating HSCR may be demonstrated by detecting the ability of the Therapeutics of the present invention, or a composition of the invention to reduce or inhibit HSCR in an animal or to ameliorate or alleviate one or more symptoms associated with HSCR. The treatment is considered therapeutic if there is, for example, a reduction in or amelioration of one or more symptoms following administration of the therapeutics, or a composition of the invention.

5.15. Animal Models

[0265] The generation of a mouse or other animal model of HSCR is important for both an understanding the biology of the disease but also for testing of potential therapies. Currently, there is no animal model available for the common nonsyndromic HSCR.

[0266] The present invention provides an animal model of HSCR by introduction of a mutant PHOX2B gene such as, but not limited to the mutant PHOX2B gene comprising the

nucleic acid sequence of SEQ ID NO: 3, in a number of species including mice, rats, pigs, and primates. Methods for specifically mutating genes by homologous recombination in embryonic stem cells (ES cells) have been described (Capecchi 1989, *Science* 244:1288). A wild-type PHOX2B gene may be replaced by a mutated PHOX2B gene. Methods for transgenic procedures are well known in the art. Murphy and Carter, 1992, *Curr. Opin. Cell. Biol.* 4:273-279.

6. EXAMPLES

[0267] The following examples are provided to illustrate certain aspects of the present invention and not intended as limiting the subject matter thereof:

6.1. Collection of Blood Samples from HSCR Patients

[0268] A total of 91 ethnic Chinese patients, diagnosed with HSCR between 1984 and 2001, were included in this study. Diagnosis was made at Queen Mary Hospital, Hong Kong SAR, and was based on histological examination of either biopsy or surgical resection material for absence of enteric plexuses. Eighty-six patients were affected with sporadic HSCR (4 individuals with total aganglionosis, 4 individuals with long-segment aganglionosis and 78 individuals with short-segment aganglionosis) and five patients (with short-segment aganglionosis) also had affected relatives (2 unrelated families in total). Within the sporadic group, 8 patients (1 with total aganglionosis and 7 with short-segment aganglionosis) presented with associated anomalies (5 with Down's syndrome; 1 with Waardenburg syndrome; 1 with renal agenesis and 1 with parathyroid adenoma). Normal controls (39 individuals) were unselected, unrelated, ethnic Chinese subjects from Hong Kong without a diagnosis of HSCR. DNA was extracted from peripheral blood as previously described (Sancandi et al. 2000). All patients and controls assented to molecular analysis. The study was approved by the local ethical committee.

6.2. Amplification and Sequencing the PHOX2B Gene

[0269] Polymerase chain reaction (PCR) was used to amplify the three coding regions (including 5' and 3' flanking regions) of PHOX2B. For exons 1 and 2, PCR was performed in a 25 μ l reaction volume containing 100 ng genomic DNA; a mixture of all four nucleotides (final concentration 0.2 mM for each nucleotide); 2.5 μ l of 10 \times reaction buffer; 0.5 μ M of each appropriate primer; 1 mM of $MgCl_2$, and 1.25 U of AmpliTaq Gold polymerase (Applied Biosystems, Foster City, Calif.). Due to its high GC content, amplification of exon 3 was performed using the GC-RICH PCR System (Roche Molecular Biochemicals, USA). Exon 3 PCR was performed in 25 μ l reaction volume containing 100 ng genomic DNA; a mixture of all four nucleotides (final concentration 0.2 mM for each nucleotide); 5 μ l of 5 \times GC-RICH PCR reaction buffer; 2.5 μ l of 5 M GC-RICH resolution solution; 0.5 μ M of each appropriate primer, and 1.25 U of a blend of Taq DNA Polymerase and Tgo DNA Polymerase.

[0270] Primers used to amplify the three coding regions, including intron-exon boundaries, were derived from the PHOX2B gene sequence available in GenBank (accession number AB015671). The primers were as follow (size of

amplified DNA is in parenthesis; forward primer is first shown in each set): exon 1: GACCTCAGACAAG-GCATCTCA (SEQ ID No. 7) and AATTACCCCTCCCTG-CAATC (SEQ ID No. 8) (586 bp); exon 2: CTGCCGTAT-GACCTGACCTT (SEQ ID No. 9) and ACAGCCACACCAAATCCAGT (SEQ ID No. 10) (442 bp); exon 3: ACCCTAACCGGTGCTTTTCT (SEQ ID No. 11) and ACAATAGCCTTGGGCCTACC (SEQ ID No. 12) (687 bp). The cycling conditions for the three exons were 95° C. for 8 min followed by 35 cycles of 95° C. for 1 min, 62° C. for 1 min and 72° C. for 45 s. A final 10 min extension was included at the end of the 35 cycles. Prior to sequencing, PCR products were column purified (Life Technologies, UK) to remove reaction buffer and unincorporated primers and visualized by running 5 μ l of each sample on 2% agarose gels. PCR products were screened for mutations by direct sequencing using a dye terminator cycle sequencing kit (ABI PRISM® Big Dye™ Terminator v 2.0 Cycle sequencing kit, Applied Biosystems, Foster City, Calif.) and an ABI 3100 automated sequencer (Applied Biosystems, Foster City, Calif.). For those samples in which DNA sequence variation had been observed, PCR amplification from genomic DNA and sequencing using both forward and reverse primers, were repeated up to five times.

[0271] The first polymorphism identified was an A to G transition at position 2250 of the SEQ ID No 1 in intron 2, 100 bp away from the intron/exon boundary. The second polymorphism identified was an A to C change at position 3493 of the SEQ ID No 1, in codon 253 of the PHOX2B exon 3. A to C2607 is a silent transition that does not alter the amino acid sequence. The third sequence variant was a 15 bp deletion at position 3495 of the SEQ ID No 1, in exon 3, starting from codon 254 and just 2 bp downstream of the A to C change at position 3493. This 15 bp deletion occurs in one of the two poly-alanine stretches in the C-terminus, resulting in the loss of five alanine residues (codons 254-258).

6.3. Association of the Newly-Discovered Alleles with HSCR

[0272] Allele and genotype frequencies for each polymorphism in the control group and the unrelated patients were calculated. Allele and genotype frequency comparisons between the control group and the patient group were performed using X^2 (chi-square) tests. Chi-square tests were also performed to determine whether each polymorphism was in Hardy-Weinberg equilibrium within each group.

[0273] The level of association between genotypes involving two or three polymorphic sites and HSCR was assessed using both chi-square contingency tables and the EH program (Terwilliger and Ott, 1994). Haplotype frequencies for each polymorphism (with and without allelic association) were estimated by the EH program. To test whether haplotype frequencies were significantly different between cases and controls, we performed three separate analysis using the EH program: on the case subjects alone, on the control subjects alone, and on the combination of case subjects and controls. The three log-likelihoods obtained (lnLcase, lnLcontrol, and lnLcombined) were used to calculate the relevant statistic test $2[\ln(L_{case}) + \ln(L_{control}) - \ln(L_{combined})]$ which gives an approximate X^2 distribution under the hypothesis that allelic association is allowed. The EH

program, which assumes Hardy-Weinberg equilibrium, was also used to estimate LD between polymorphisms.

TABLE 1

Genotype frequency distribution of the PHOX2B gene polymorphisms		
Polymorphisms	Total patients (91)	Total controls (39)
<u>A/G₂₂₅₀</u>		
AA	0.81 (73)	0.62 (24)
AG	0.16 (15)	0.38 (15)
GG	0.03 (3)	0.00 (0)
	$\chi^2 = 8.28; p = 0.015$	
<u>A/C₃₄₉₃</u>		
AA	0.90 (82)	0.85 (33)
AC	0.10 (9)	0.15 (6)
CC	0.00 (0)	0.00 (0)
	$\chi^2 = 0.72; p = 0.39$	
<u>DEL₃₄₉₅</u>		
NN ^a	0.92 (84)	0.82 (32)
ND ^b	0.08 (7)	0.18 (7)
DD	0.00 (0)	0.00 (0)
	$\chi^2 = 2.99; p = 0.08$	

Number of individuals in brackets. ^aN no deletion; ^bD deletion;

[0274] Comparison of the genotype frequencies of the three new polymorphisms between patient and control groups (Table 1) revealed statistically significant differences for A to G1364 ($X^2=8.28; p=0.015$). For this particular polymorphism, genotypes comprising G were under-represented in the patient group (19% in patients versus 38% in controls). Within each group, no statistically significant differences in genotype frequencies for any of the identified polymorphisms were found between genders. Within each group, each of the three polymorphic sites was in Hardy-Weinberg equilibrium.

6.4. Identification of Additional Polymorphisms in PHOX2B Associated with HSCR

[0275] The procedures for evaluating the candidate gene PHOX2B, as well as for finding additional polymorphisms, will be the same as for those already identified. Mutation detection in the coding sequence of these genes will consist of PCR amplification and sequencing. Mutation detection for the entire genes will include large deletion/insertion analysis of 200-400 bp fragments by SSCP or heteroduplex analysis, and of course sequencing when heterozygosity becomes apparent for any region of the genes. Current Protocols in Human Genetics (John Wiley & Sons, Inc.), Chapter 7, "Searching Candidate Genes for Mutations." Biological samples already isolated from patients with HSCR which did not show any abnormalities in PHOX2B will be screened for polymorphisms mother HSCR genes. Although preferred embodiments have been depicted and described in detail herein, it will be apparent to those skilled in the relevant art that various modifications, additions, substitutions, and the like can be made without departing from the spirit of the invention and these therefore are considered within the scope of the invention.

7. RESULTS

7.1. Polymorphic Sites

[0276] By direct sequencing the entire coding region and portions of 5' and 3' flanking regions, two new SNPs and one

deletion in the PHOX2B gene (FIG. 8) were identified. Nucleotide positions are defined in relation to the first nucleotide of the start codon, which is designated position+1.

[0277] The first polymorphism identified was an A→G transition (referred as AG 2250) in intron 2, 100 bp away from the intron/exon boundary. The second polymorphism identified was an A→C change (referred as AC 3493) of the PHOX2B exon 3. AC 3493 is a silent transition that does not alter the amino acid sequence (GCA GCC). The third sequence variant was a 15 bp deletion (referred as DEL 3495) in exon 3, just two base pair downstream of the AC 3493 polymorphism (FIG. 8). This 15 bp deletion occurs in one of the two poly-alanine stretches in the C-terminus, resulting in the loss of five alanine residues. The poly-alanine region in the normal PHOX2B protein may form α-helical structures and serve as a flexible linker (Shoemaker et al. 1987). Among the 39 normal individuals studied, 7 individuals had this deletion in one of their PHOX2B alleles. Specific PCR amplification enzymes were used to resolve the high GC sequence content of exon 3. In addition, those samples with the deletion were re-amplified and re-sequenced 5 to 10 times.

[0278] Finally, in every sample analyzed there was an A at both positions 1129 and 2304 of the PHOX2B gene (intron 2 and 3 respectively; FIG. 8). The Genbank accession numbers for PHOX2B (AB015671 and AFI 17979) record T and G respectively. These may represent sequencing errors in the GenBank entries. Alternatively, the Chinese population analyzed in this study has these SNPs.

[0279] In addition to the three polymorphisms described in this study, there are three other PHOX2B SNPs recorded in GenBank. Polymorphism ss3136952 (also rs2196822) is a A→C 3004 transition at position 3004 of the PHOX2B gene (intron 2); polymorphism ss1551093 (also rs1063611) is a T→A 5343 transversion at position 5343 (end of 3' untranslated region) and polymorphisms ss1551094 (also rs1063612) is a C→A5403 transversion at position 5403 (end of 3' untranslated region).

7.2. Allelic Variation Between Patients and Controls

[0280] To determine the significance of these three novel polymorphisms in HSCR, a case-control association study was conducted to investigate whether they were directly related to the HSCR phenotype. Differences in allele frequencies between the patient group and the control group were not statistically significant for any of the polymorphisms (AC 2250 X²=0.76; p=0.38; DEL3495 X²=2.82; p=0.09). Also within each group, allele frequencies of each gender were compared. No statistically significant differences for any of the identified polymorphisms were found.

7.3. Haplotype Frequencies and Linkage Disequilibrium Among the Three Polymorphic Sites

[0281] The level of association between HSCR and the genotypes involving two or three polymorphic sites was also assessed. When comparing frequencies of haplotypes generated by the combination AG2250+AC3493 and by the combination AG2250+DEL3495 statistically significant differences between patients and controls were found for both haplotype combinations (X²=8.45; p=0.03 and X²=10.13;

p=0.03 respectively). Contributions to X² were dissected and demonstrated that the major contribution was that of genotypes of the polymorphic site AG2250. In contrast, no statistically significant difference in frequencies was found for the combination AC 3493+DEL3495. When haplotypes comprising the three polymorphisms were analyzed for differences in frequencies, a statistical trend toward association (X²=9.47; p 0.05, 4 df) was found mainly due to the major contribution of the AG 2250 polymorphism. Using the EH program, the haplotype frequencies of the three polymorphisms in 91 patients and 39 controls were estimated (Table 2).

TABLE 2

Estimated haplotype frequencies in the 91 patients and 39 control determined by EH					
Haplotypes					
Intro 2	Exon 3		Frequencies		
	AG 2250	AC 3493	DEL 3495	Patients	Controls
A	A	ND	0.855	0.795	0.828
A	A	D	0.029	0.028	0.029
A	C	ND	0.000	0.014	0.004
A	C	D	0.000	0.000	0.000
G	A	ND	0.056	0.067	0.060
G	A	D	0.009	0.061	0.024
G	C	ND	0.049	0.062	0.053
G	C	D	0.000	0.000	0.000

[0282] The two most common haplotypes for both cases and controls were A-A-ND and G-A-ND (allele designations as shown in Table 1). When haplotype frequencies were compared, none of the haplotype combinations showed statistically significant differences in frequencies between patients and controls (Table 3). Hence, the associations found were driven by the presence of AG 2250.

TABLE 3

Association testing based on the haplotype frequencies of AG 2259, AC 3493 and DEL 3495				
Group	N ^a	In(L)	X ²	p ^b
Patients	91	-90.84	36.54	0.3
Controls	39	-55.89	15.19	
All subjects	130	-150.92	49.24	

^aNumber of individuals; ^bCalculated from 2[ln(Lcase) + ln(Lcontrol) - ln(Lcombined)]; df = 7

[0283]

TABLE 4

Pairwise comparison	Pairwise linkage disequilibrium					
	Patients		Controls		All subjects	
	X ²	P	X ²	p	X ²	p
AG 2250 vs. AC 3493	34.67	0.00	6.36	0.50	38.68	0.00

TABLE 4-continued

Pairwise comparison	Pairwise linkage disequilibrium					
	Patients		Controls		All subjects	
	X ²	P	X ²	p	X ²	p
AG 2250 vs. DEL 3495	0.12	0.90	4.08	0.25	2.13	0.50
AC 3493 vs. DEL 3495	0.72	0.86	1.18	0.75	1.29	0.70

[0284] The AG 2250 polymorphism was in LD with AC 3493 when cases and controls were analyzed combined and in the patient population. AG 2250 was not in LD with AC 3493 in the control group. No LD was found between AG 2250+DEL 3495, nor between AC 3493+DEL 3495, despite being two base pairs apart only. The AG 2250 polymorphism is directly associated with HSCR and that the discrepancies in the analysis of haplotype frequencies are due to incomplete LD.

8. DISCUSSION

[0285] SNPs are used as markers in human genetics studies ranging from comparative population variation (Chakravarti 1999) to disease linkage studies (Kruglyak 1997). SNPs also have potential as direct functional polymorphic variants involved in common and genetically complex human diseases (Collins et al. 1997) and pharmacogenetic traits (Housman and Ledley 1998). In addition, SNPs can also serve as surrogates for unrecognized functional SNPs that may be identified by disease-marker LD studies (Chakravarti 1999; Collins et al. 1997).

[0286] The entire coding region of the PHOX2B gene was sequenced as well as intron/exon boundaries of ethnically matched patients and controls in the search for mutations or polymorphisms that could contribute to the clinical manifestation of HSCR. Two new SNPs (AG 2250, AC 3493) and one deletion (DEL 3495) have been identified. The DEL 3495 showed no association with HSCR despite causing the loss of 5 alanine-residues in the PHOX2B protein. In both patients and controls, the DEL 3495 was always heterozygous. In mice, heterozygous disruption of PHOX2B homeodomain shows no obvious phenotype, but homozygous mutants are embryonic lethal. In those individuals carrying DEL 3495, any functional consequence caused by the DEL 3495 polymorphism may be compensated by the normal allele.

[0287] For AG 2250 statistically significant difference in genotype distribution was found when patients were compared to controls indicating an association of AG 2250 polymorphism with HSCR. This under-representation may indicate that the transition AG 2250 is relatively recent in evolutionary terms, as shown by the relatively low frequency of the allele G in the population. If the minor allele G conferred a protective effect, its frequency in the population would be rising to become the major allele (Cargill 1999).

[0288] The association found may be due to a direct contribution of AG 2250 to the HSCR phenotype or being in LD with another susceptibility locus. Direct involvement of AG 2250 in HSCR may be through the alteration of intronic

sequences crucial for splicing and/or regulation of the expression of PHOX2B gene. Genotypes comprising allele G were under represented in the patient group (19% vs 38%; $X^2=8.28$; $p=0.015$) and in males (19% vs. 44%; $X^2=8.71$; $p=0.012$) compared with male controls. The association of PHOX2B AG 2250 polymorphism with HSCR may represent a protective effect against HSCR phenotype in male or provide a linkage marker for other HSCR susceptibility gene loci.

[0289] There were LD among the three polymorphisms that spread over 1.2 Kb, which was expected given their physical proximity. Pairwise disequilibrium test showed lack of LD between AC 3493 and DEL 3495 polymorphisms which are only two base pairs apart. This indicates that these two polymorphisms are probably situated in a recombination hot spot. Detailed sequence examination of exon 3 (from nucleotide 3176 to 3691) in which these polymorphisms are located showed that there is a high GC content and CGG repeats which may serve as foci for recombination. Recombination across the hot spot would explain the results obtained in the pairwise disequilibrium test and haplotype analysis. This suggests that if there are other polymorphisms or mutations in and around the PHOX2B gene which are also contributing to HSCR, they should be located upstream of the AG 2250 polymorphism. These other possible susceptibility loci may not be in LD with the polymorphisms (AC 3493 and DEL 3495) found in the recombination hot spot that defines the breakpoint of a LD block (Goldstein 2001).

[0290] In humans both the RET and EDNRB signaling pathways are crucial in the development of enteric neurons and have been associated to HSCR. Nevertheless, mutations in RET account for less than 50% of the HSCR patients challenging the understanding of the disease etiology. It is likely that a combination of specific variations in multiple genes of both the RET and EDNRB signaling pathways confer either protection or susceptibility to the disease. Co-existence of RET mutations with mutations or polymorphisms in GDNF and EDNRB genes have been described in patients with HSCR (Hofstra et al. 1997; Salomon et al. 1996). Therefore, there may be a role for PHOX2B in HSCR, acting in combination with mutations in RET or other genes such as EDNRB, EDN3, and GDNF genes.

[0291] All references (including books, articles, papers, patents, and patent applications) cited herein are hereby expressly incorporated by reference in their entirety for all purposes. While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modification, and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice in the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth, and as fall within the scope of the invention and the limits of the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 6

<210> SEQ ID NO 1

<211> LENGTH: 5508

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```
cctgagtctc ttgaaagcac atttagctgc aagaagaaat gcaatttata agatgctttc    60
tctctgtctc tctttgtttt taacccccctc taaccagggtg tattttttaa gaaatccgct   120
tatcgttcac ataaaccccc ttggcccact tactctatgt tacagggcgc ctgagtcttg   180
ccaatgtccc agtcctttat aacatttcat gcacttcggg gggtaggctt gttgttaaat   240
tgagcgtgta acactcttac aaaacagggtt ttctatgaca tcaaggtttc ttctccctaa   300
ccgaggggga aaaaaagaag aggacgaaga agggggaaaa cacacacact atctcaattt   360
atgcctaagg tatatgatca gttaaaaagg cttaaaagct cggggaaatt ggatcaggga   420
gaatcgtcac ccaactttca ttatttccaa gtagtgtgat tgaattaaag ggcagggagc   480
tggttagaag ggaggatcag gggctcgggtg cgtaatgggtg tggattaaa ttctaattag   540
agatgcagga atcaatgata gggaggttgg acagctcagt tcccagtgcc cagcccaata   600
gacggatgag ttattgtcat gtaaaaagcg ccagcaataa gaccaaccgc ttgtctattg   660
tccaagtgga aagagccaag tttattatga ggactatatg ctctagagac ctccagacaag   720
gcatctcata ggagggtttt tcataaaact aggctctgct ggtagtaagg aggccagttt   780
ggaggcaggc gttgagctgt gcacatctcc ccactccagc cacctctctc atatccatct   840
tttattttat ttttccactt ggctgagcca tccagaaacct tttcaatgta taaaatggaa   900
tattcttacc tcaattctct tgcctacgag tcctgtatgg ctgggatgga cacctcagac   960
ctggcttcag cctatgctga cttcagttcc tgcagccagc ccagtggtt ccagtataac  1020
ccgataagga ccaacttttg gccacagtcg ggctgcccct ccctcacgcc gggatcctgc  1080
agcctgggca ccctcaggga ccaccagagc agtccgtacg ccgaggtta ggaccttcag  1140
ctttctcagc ggaggaagcc gcctttccgc ccgtatatag gaagccttga ttgcatttga  1200
aaatggaaa gtgttttagta tttacaaaac gaaatttgct tacacaaatg aaagaattta  1260
tcacgtttga agcgattgca gggaggggta attcaattac agggttacac tatcctagtc  1320
acacccgaac cgcccacaaa attatcttaa gctgccaaaa tgataggcat aatttattta  1380
ctttgcgatg agacgtatag cttagaaaat aattgaatta caaagagtaa agctcattac  1440
tggcagtgtc tcttttttta agaaccgaga gcggctcaca cttcttgggc tggtcatttt  1500
tatgattatt tctttaattt attattattt ttttgcagct ctttcccca acttttgagc  1560
cgggtcaact ttctgagaat tgaaaagtcc ccaaagtggg actgtttggg aacttctttc  1620
cctggtctcc ctgatattcc gactgatggt ttgggatttt tttctctctt ggttttttcc  1680
tgctgaaagc actatctcaa gtccgtcaca tcgctgtgtt tcaatccacc caaaggogct  1740
tgtgccagaa aggactccgc caagcccga gtttgagccc aggtttccgc agataacaaa  1800
tttctcgggt ttcttccgcg agcttctctc ggcaactctc tcgctcgggt gtaggtagcg  1860
gctgccgtat gacctgacct tggagctctc acattctagc tccacggccg gcgagctgcc  1920
```

-continued

ggctgatttg	ctcactttct	gtctcctctg	tcatactcta	gttccttaca	aactcttcac	1980
ggaccacggc	ggcctcaacg	agaagcgcaa	gcagcggcgc	atccgcacca	ctttcaccag	2040
tgccagctc	aaagagctgg	aaagggcttt	cgcgagact	cactaccccg	acatctacac	2100
tcggaggag	ctggccctga	agatcgacct	cacagaggcg	cgagtccag	tacgcgcgcc	2160
tggaaaccga	ccccgctccg	ccgcactgg	ccggggaggt	gtggggtag	ggcgggctgg	2220
tgaaattcga	agtcctggag	cctcgagtga	gaaggaccta	gggcccctg	gccgatcaga	2280
aatactggat	ttgggtggc	tgtgcgttcg	agagaggctt	agagcgcacg	ctcttggcat	2340
tttatttaca	gttgcaagt	gtttcccacc	cgagcagaga	catggggggc	cttgggacgt	2400
ggatgagcga	tgcaatttcg	gggacaggaa	gtgcctgtgg	tggaaagtgt	gcagactttg	2460
ctcccgatt	ataagttttt	ccttctccc	tcccgcccc	caaaaaatg	cctcctaact	2520
caagtgcctt	taacctggcc	ccatggcata	taggttcatt	ttccggaaa	ctgtgacttg	2580
cattagattt	gcaaagggtc	tgtgacttca	tgaaggtcaa	gaacctgac	ttactccaac	2640
ctgttaaca	caggtgcgct	cacgagttgg	ccacagcgc	tttttgggtg	agccccgac	2700
cgagaagcgg	tgcgcacct	tgcacgcttt	tccaggtca	aaggccgggg	atgggcagcg	2760
gagcaaacc	agagaggatc	cctttctctt	ttaccaatta	gagtttaact	ttagaactta	2820
ggcttagggg	tgaatggcga	gctcggggct	tgtcaagaa	gccgactgaa	cagaggccca	2880
ccaaaaaag	gcctccctt	tccgggtctt	tctgggacct	cgggcttttt	aaactctgcc	2940
gcaagcctc	atgtccctgg	cggtctcact	ccccctaaga	aagtttctcc	gaaaatgac	3000
agcaataaga	agcggtagac	ttgggtgatg	tgcgcgggg	ggtgatcaca	gcgcatgggg	3060
aggaggggtg	taaaacaagc	cgaagtagaa	cttgggccac	cctaaccogt	gcttttcttt	3120
cccattttct	tctttctccc	cctgcttcc	cgctctctct	tccgtcttg	gccaggtgtg	3180
gttccagaac	gcgcggcca	agtctcgcaa	gcaggagcgc	gcagcggcag	ccgcagcggc	3240
cgcgccaag	aacggctcct	cgggcaaaaa	gtctgactct	tccagggacg	acgagagcaa	3300
agagccaag	agcactgacc	cggacagcac	tgggggccca	ggtccaatc	ccaacccac	3360
ccccagctg	ggggcgaatg	gagggcggcg	cggcgggcc	agcccggctg	gagctccggg	3420
ggcggcgggg	cccggggggc	cgggagggca	acccggcaag	ggcggcgcag	cagcagcggc	3480
ggcgcccgcg	gcagcggcgg	cggcggcagc	ggcagcggcg	gcagctggag	gcttggtctg	3540
ggctgggggc	cctggacaag	gctgggctcc	cggccccggc	cccacacct	ccatcccggg	3600
ttcgcttggg	ggtcccttcg	gcagcgtcct	atcttcgctc	caaagacca	acggtgcaa	3660
agccgcctta	gtgaagagca	gtatgttctg	atctggaatc	ctgcggcggc	ggcggcggcg	3720
gcgacagcgg	gcgagccagg	gcccggcggg	gcgagtgggc	gagcgggtag	gcccaggct	3780
attgtcgtcg	ctgctgccat	ggctttttca	ttgaggcct	aaagtaatcg	cgctaagaat	3840
aaagggaaaa	cggcgtcgcc	ctcatttcaa	ccccactcct	acccccttc	tcaacccca	3900
aacaaaacaa	acaaaacttc	ctggcttcgc	acctgcctgg	ggcctcgag	cggggccagg	3960
gctccgctg	ctgatogggg	gttgtgagca	gcgcggcctg	gaocggggc	actctcaggg	4020
ggctgtgtct	cgctgtcagt	ttgtgtctgt	ctcggggaat	gtgtgtctgt	ggccaagca	4080
ggtgacagga	agagatgggg	ggcctcaacc	aacttagtga	cttgtttaga	aaaaaaagac	4140
aaaaaagtaa	aaataaaaac	aaaaaagttg	gaaggcagaa	accattaaaa	aacaaaaagc	4200

-continued

```

caacaaccca gaaaggttta aaaaacataa ggaaaaaaaa gacaaattaa aggaggggct 4260
aggggagaag ctgcagctgg agctgaaggc tcgatcttgt gaaccocctaa atccgctccc 4320
toctaacagc acggattctc ttggggctct tcttcaggga agagtaggga cgccgttcca 4380
gccccocctc ctatcgtgto cttgggttcg ggtcactgog gcgacgactt gtcagactg 4440
tcccgccatt agcgcacatc aacccacac agttgcaact cccaaccccg aatccttgca 4560
gccgttcggc cctgaaagat gccctatcca tgagatgcct tttcatctgc aaactctgca 4620
aaatgtgtct catgtttcgc aactcttttt tccccctcg ctcccgccta ccccgctggc 4680
atcttctctc tccaccagct tttactgaac tttttggcac tgctttggat tggggcaat 4740
tgcagtcacc gtaactggct gcagagaaat ctaccgagca aggaaaaggc acacacacac 4800
gtttgcaggg gtgtctcggg ttgcatttct gttggaatga tccgaactgg actcacatcc 4860
tgataggtgg atggactgta tattgagggt tccattcttc gcgcagttta gacatctctg 4920
ttttgattct ttgttgttgt ttttatttta aaaggcacia actctagata ttagttgaat 4980
gttgaggctt taactttttc ggtgtctttc tacaactgtg ttctgtgact caattgtatc 5040
gtgttaatat cagtgcagac tgtctcctct acgtgaccgt ataatgtttt tctcgtottg 5100
tagtctctat ggcgtgtctt tatggtgtaa taaggttctc acggggtaa tcttttgtgt 5160
ttagagaggc cacggttcag acaatgggat atatttttgt tatcagggtc atgtctgtct 5220
gatttctttt ttttctcgtg tggactatgt ttgtgaacat aattgtcata agttatgttt 5280
cagatttttg aatttattta tatgtgttat aatgaatgct tctatttaaa agggaaatat 5340
ttctacatgt gcttatagtt ttccaagat gtaccattaa cttgattggt gataataaaa 5400
accaaaagca agtctagcaa ttgaaactct ctttttcttg attccttttt tttttttttt 5460
tttgggttgg tcattgtttt ttttttttaa gttttttttt aaaaggat 5508

```

<210> SEQ ID NO 2

<211> LENGTH: 314

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

```

Met Tyr Lys Met Glu Tyr Ser Tyr Leu Asn Ser Ser Ala Tyr Glu Ser
 1           5           10           15

```

```

Cys Met Ala Gly Met Asp Thr Ser Ser Leu Ala Ser Ala Tyr Ala Asp
          20           25           30

```

```

Phe Ser Ser Cys Ser Gln Ala Ser Gly Phe Gln Tyr Asn Pro Ile Arg
          35           40           45

```

```

Thr Thr Phe Gly Ala Thr Ser Gly Cys Pro Ser Leu Thr Pro Gly Ser
          50           55           60

```

```

Cys Ser Leu Gly Thr Leu Arg Asp His Gln Ser Ser Pro Tyr Ala Ala
 65           70           75           80

```

```

Val Pro Tyr Lys Leu Phe Thr Asp His Gly Gly Leu Asn Glu Lys Arg
          85           90           95

```

```

Lys Gln Arg Arg Ile Arg Thr Thr Phe Thr Ser Ala Gln Leu Lys Glu
          100          105          110

```

```

Leu Glu Arg Val Phe Ala Glu Thr His Tyr Pro Asp Ile Tyr Thr Arg
          115          120          125

```

```

Glu Glu Leu Ala Leu Lys Ile Asp Leu Thr Glu Ala Arg Val Gln Val

```

-continued

130	135	140
Trp Phe Gln Asn Arg Arg Ala Lys Phe Arg Lys Gln Glu Arg Ala Ala 145 150 155 160		
Ala Ala Ala Ala Ala Ala Ala Lys Asn Gly Ser Ser Gly Lys Lys Ser 165 170 175		
Asp Ser Ser Arg Asp Asp Glu Ser Lys Glu Ala Lys Ser Thr Asp Pro 180 185 190		
Asp Ser Thr Gly Gly Pro Gly Pro Asn Pro Asn Pro Thr Pro Ser Cys 195 200 205		
Gly Ala Asn Gly Gly Gly Gly Gly Gly Pro Ser Pro Ala Gly Ala Pro 210 215 220		
Gly Ala Ala Gly Pro Gly Gly Pro Gly Gly Glu Pro Gly Lys Gly Gly 225 230 235 240		
Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala 245 250 255		
Ala Ala Ala Ala Gly Gly Leu Ala Ala Ala Gly Gly Pro Gly Gln Gly 260 265 270		
Trp Ala Pro Gly Pro Gly Pro Ile Thr Ser Ile Pro Asp Ser Leu Gly 275 280 285		
Gly Pro Phe Gly Ser Val Leu Ser Ser Leu Gln Arg Pro Asn Gly Ala 290 295 300		
Lys Ala Ala Leu Val Lys Ser Ser Met Phe 305 310		

<210> SEQ ID NO 3
 <211> LENGTH: 5508
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

```

cctgagtctc ttgaaagcac atttagctgc aagaagaaat gcaatttata agatgctttc 60
tctctgtctc tctttgtttt taaccccctc taaccagggtg tattttttaa gaaatcogct 120
tatcgttcac ataaaccccc ttggcccact tactctatgt tacagggcgc ctgagtcttg 180
ccaatgtccc agtcctttat aacatttcat gcacttcggg gggtaggctt gttgttaa 240
tgagcgtgta acactcttac aaaacagggt ttctatgaca tcaaggtttc ttctccctaa 300
ccgaggggga aaaaaagaag aggacgaaga agggggaaaa cacacacact atctcaattt 360
atgcctaagg tatatgatca gttaaaaagg cttaaaagct cggggaaatt ggatcaggg 420
gaatcgtcac ccaactttca ttatttccaa gtagtgtgat tgaattaaag ggcagggagc 480
tggttagaag ggaggatcag gggctcgtg cgtaatggtg tggattaaa ttctaattag 540
agatgcagga atcaatgata gggaggttgg acagctcagt tccccagtgc cagccaata 600
gacggatgag ttattgtcat gtaaaaagcg ccagcaataa gaccaaccgc tttgctattg 660
tccaagtgga aagagccaag tttattatga ggactatatg ctctagagac ctgagacaag 720
gcatctcata ggaggctttt tcataaaact aggctctgct gtagtaagg aggccagttt 780
ggagggcagg gttgagctgt gcacatctcc ccaactccgc cacctctccc atatccatct 840
tttatttcat ttttccactt ggctgagcca tccagaacct tttcaatgta taaaatggaa 900
tattcttacc tcaattccto tgcctacgag tcctgtatgg ctgggatgga cacctcagac 960
ctggettcag cctatgctga cttcagttcc tgcagccagg ccagtggctt ccagtataac 1020
    
```

-continued

cogataagga	ccacttttgg	ggccacgtcc	ggetgccctt	ccctcacgcc	gggatcctgc	1080
agcctggggc	ccctcagggc	ccaccagagc	agtccgtacg	ccgcagggtta	ggaccttcag	1140
ctttctcagc	ggaggaagcc	gcctttccgc	ccgtatatag	gaagccttga	ttgcatttga	1200
aatggaaat	gtgttttagta	tttaccaaaac	gaaatttgct	tacacaaatg	aaagaattta	1260
tcacgtttga	agcgattgca	gggaggggta	attcacttac	agggttacac	tatcctagtc	1320
acccccgaac	cgccccaaaa	attatcttaa	gctgcacaaa	tgataggcat	aattttattta	1380
ctttgcatg	agacgtatag	cttagaaaat	aattgaatta	caaagagtaa	agctcattac	1440
tggcagtgtc	tcttttttta	agaaccgaga	gcggctcaca	cttcttgggc	tggtcatttt	1500
tatgattatt	tctttaattt	attattattt	ttttgcagct	ctttccccc	acttttgagc	1560
cggtcaact	ttctgagaat	tgaaaagttc	ccaaagtggg	actgtttgg	aacttctttc	1620
cctggctccc	ctgatattcc	gactgatgtt	ttgggatttt	tttctctct	ggttttttcc	1680
tgctgaaagc	actatctcaa	gtccgtcaca	tcgctgtgtt	tcaatccacc	caaaggcgtc	1740
tggtccagaa	aggactccgc	caagcccga	gtttgagccc	aggtttccgc	agataacaaa	1800
tttctcggg	ttcttccgc	agctctctc	ggcaactctc	tcgctgggt	gtaggtagcg	1860
gctgccgat	gacctgacct	tggagctctc	acattctagc	tccacggccg	gogagctgcc	1920
ggctgatttg	ctcactttct	gtctctctg	tcatactcta	gttccttaca	aactcttcac	1980
ggaccacggc	ggcctcaacg	agaagcgca	gcagcggcgc	atccgcacca	ctttcacccag	2040
tgcccagctc	aaagagctgg	aaaggtctt	cgcgagact	cactaccccg	acatctacac	2100
tcgggaggag	ctggccctga	agatcgacct	cacagaggcg	cgagtccagg	tacgcgcgcc	2160
tggaaaccga	ccccgctccg	ccgcactgg	ccggggaggt	gtgggtgag	ggcggtctgg	2220
tgaaattcga	agtcctggag	cctcgagtgg	gaagcaccta	gggccccatg	gccgatcaga	2280
aatactggat	ttggtgtggc	tgtgcttctg	agagaggctt	agagcgcacg	ctcttggcat	2340
tttatttaca	gttgcaagt	gtttcccacc	cgagcagaga	catggggggc	cttgggacgt	2400
ggatgagcga	tgcaatttcg	gggacaggaa	gtgcctgtgg	tggaaagggt	gcagactttg	2460
ctcccgtatt	ataagttttt	ccttctcccc	tcccggcccc	caaaaaatg	cctcctaact	2520
caagtgtttt	taacctggcc	ccatggcata	taggttcatt	ttcccggaaa	ctgtgacttg	2580
cattagattt	gcaaagggtc	tgtgacttca	tgaaggtaaa	gaacctgac	ttactccaac	2640
ctgttaaa	caggtgcgct	cacgagttgg	ccacagcgcc	tttttgggtg	agccccgac	2700
cgagaagcgg	tgcgaccat	tgcaacgctt	tccaggctca	aaggccgggg	atgggcagcg	2760
gagcaaaccc	agagaggatc	ccttttctct	ttaccaatta	gagtttaact	ttagaactta	2820
ggcttagggg	tgaattggca	gctcggggct	tgctcaagaa	gccgactgaa	cagaggccca	2880
ccaaaaataag	gccttccctt	ttcgggtctt	tctgggacct	gcggcttttt	aaactctgcc	2940
gcaagccttc	atgtccctgg	cggtctcact	ccccctaaga	aagtttctcc	gaaaatgcac	3000
agcaataaga	agcggtagac	ttgggtgatg	tgccgcggg	ggtgatcaca	gcgcatgggg	3060
aggaggggtg	taaaacaagc	ggaagtagaa	cttgggccac	cctaaccggt	gcttttcttt	3120
ccattttct	tctttctccc	cctgcttca	cgctctctct	tcogtcttgg	gccaggtgtg	3180
gttccagaac	cgcccgccca	agtttcgcaa	gcagagcgc	gcagcggcag	ccgcagcggc	3240
cgcgccaag	aacggctcct	cgggcaaaaa	gtctgactct	tccagggacg	acgagagcaa	3300

-continued

agaggccaag	agcactgacc	cggacagcac	tgggggcca	ggtccaatc	ccaaccocac	3360
ccccagctgc	ggggcgaatg	gaggcggcgg	cggcgggccc	agcccggtg	gagctccggg	3420
ggcggcgggg	cccgggggccc	cgggagggca	acccggaag	ggcggcgag	cagcagcggc	3480
ggcggcccg	gcagcggcgg	cggcggcagc	ggcagcggc	gcagctggag	gcctggctgc	3540
ggctggggg	cctggacaag	gctgggctcc	cggccccggc	cccatcacct	ccatcccgga	3600
ttcgcttgg	ggtcccttcg	gcagcgtcct	atcttcgctc	caaagacca	acggtgcca	3660
agccgcctta	gtgaagagca	gtatgttctg	atctggaatc	ctgcggcggc	ggcggcggcg	3720
gcgacagcgg	gcgagccagg	gcccggcgg	gcgagtggc	gagcgggtag	gccaaggct	3780
attgtcgtcg	ctgctgccat	ggctttttca	ttgagggcct	aaagtaatcg	cgctaagaat	3840
aaaggaaaa	cggcgtcgcc	ctcatttcaa	ccccactcct	acccccctcc	tcaaccccca	3900
aacaaaacaa	acaaacttcc	ctggcttcgc	acctgcctgg	ggcctcgag	cggggccagg	3960
gctccgcctg	ctgatcgggg	ggtgtgagca	gcgcgccctg	gacgcggggc	actctcaggg	4020
ggctgtgtct	gcgtgtcagt	ttgtgtctgt	ctcggggaat	gtgtgtctgt	ggcccaagca	4080
ggtgacagga	agagatgggg	ggcctcaacc	aacttagtga	ctgttttaga	aaaaaaagac	4140
aaaaaagtaa	aaataaaaa	aaaaaagttg	gaaggcagaa	accattaaaa	aacaaaaagc	4200
caacaaccca	gaaaggttta	aaaaacataa	ggaaaaaaa	gacaaattaa	aggaggggct	4260
aggggagaag	ctgcagctgg	agctgaaggc	tcgatcttgt	gaaccoccaa	atccgctccc	4320
tcctaacagc	acggattctc	ttggggctct	tcttcaggga	agagttagga	cgccgttcca	4380
gcccccttc	ctatcgtgtc	cttgggttcg	ggtcactgcg	gcgacgactt	gctcagactg	4440
tcccgccggc	cggagtgaact	ttctcgcacc	cccttgctg	tccacctcg	ctgaacacca	4500
tcccgccatt	agcgcctcgg	aaccacacac	agttgcaact	cccaaccccg	aatccttgca	4560
gocgttcggc	cctgaaagat	gccctatcca	tgagatgcct	tttcatctgc	aaactctgca	4620
aatgtgtct	catgtttcgc	aactcttttt	ttccccctcg	ctcccccta	ccccgtggc	4680
atctctctct	tccaccagct	tttactgaac	tttttggcac	tgctttggat	tggggtcaat	4740
tgacgtccac	gtaactggct	gcagagaaat	ctaccgagca	aggaaaaggc	acacacacac	4800
gtttgcaggg	gtgtctcggg	ttgcatttct	gttggaaatga	tccgaactgg	actcacatcc	4860
tgataggtgg	atggactgta	tattgagggt	tccattcttc	gcgcagttaa	gacatctctg	4920
ttttgattct	ttgttgttgt	ttttatttta	aaaggcacia	actctagata	ttagttgaat	4980
gttgaggctt	taactttttc	ggtgtcttcc	tacaactgtg	ttctgtgact	caattgtatc	5040
gtgttaatat	cagtgcagac	tgtctcctct	acgtgaccgt	ataatgtttt	tctcgtcttg	5100
tagtctctat	ggcgtgtctt	tatggtgtaa	taaggttctc	acggggtcaa	tcttttgtgt	5160
ttagagaggc	cacggttcag	acaatggtat	atatttttgt	tatcagggtgc	atgtctgtct	5220
gatttctttt	tttttctgt	tggactatgt	ttgtgaacat	aattgtcata	agttatgttt	5280
cagatttttg	aattttat	tatgtgttat	aatgaatgct	tctattttaa	aggaaatat	5340
ttctacatgt	gcttatagtt	ttccaagagt	gtaccattaa	cttgattggt	gataataaaa	5400
acaaaagca	agtctagcaa	ttgaactctt	ctttttcttg	attctttttt	tttttttttt	5460
tttgggttgg	tcattgtttt	ttttttttta	gttttttttt	aaaaggat		5508

-continued

```

<210> SEQ ID NO 4
<211> LENGTH: 5508
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

cctgagtctc ttgaaagcac atttagctgc aagaagaaat gcaatttata agatgctttc    60
tctctgtctc tctttgtttt taaccccctc taaccagggtg tattttttaa gaaatccgct    120
tatcgttcac ataaaccccc ttggcccact tactctatgt tacagggcgc ctgagtcttg    180
ccaatgtccc agtcctttat aacatttcat gcacttcggg gggtaggctt gttgttaa at    240
tgagcgtgta acactcttac aaaacagggt ttctatgaca tcaaggtttc ttctccctaa    300
ccgaggggga aaaaaagaag aggacgaaga agggggaaaa cacacacact atctcaattt    360
atgcctaagg tatatgatca gttaaaaagg cttaaaagct cggggaaatt ggatcagggg    420
gaatcgtcac ccaactttca ttatttccaa gtagtgtgat tgaattaaag ggcagggagc    480
tggttagaag ggaggatcag gggctcgtg cgtaatggtg tggattataa ttctaattag    540
agatgcagga atcaatgata gggaggttgg acagctcagt tcccagtgcc cagccaata    600
gacggatgag ttattgtcat gtaaaaagcg ccagcaataa gaccaaccgc tttgctattg    660
tccaagtgga aagagccaag tttattatga ggactatatg ctctagagac ctgagacaag    720
gcatctcata ggaggtcttt tcataaaact aggctctgct ggtagtaagg aggccagttt    780
ggaggcaggc gttgagctgt gcacatctcc ccactccagc caocttctcc atatccatct    840
tttatttcat ttttccactt ggctgagcca tccagaacct tttcaatgta taaatggaa    900
tattcttacc tcaattcctc tgcctacgag tcctgtatgg ctgggatgga cacctcgagc    960
ctggcttcag cctatgctga ctccagttcc tgcagccagg ccagtggctt ccagtataac   1020
ccgataagga ccacttttgg gccacagtcc ggctgccctt ccctcacgcc gggatcctgc   1080
agcctggggc ccctcaggga ccaccagagc agtccgtacg ccgcagggtt ggaccttcag   1140
ctttctcagc ggaggaagcc gcctttccgc ccgtatatag gaagccttga ttgcatttga   1200
aaatggaaat gtgttttagta tttaccaaac gaaatttgct tacacaaatg aaagaattta   1260
tcacgtttga agcgattgca gggaggggta attcacttac agggttacac tatcctagtc   1320
acacccgaac cgcccacaaa attatcttaa gctgccaaaa tgataggcat aatttattta   1380
ctttgcatg agacgtatag cttagaaaat aattgaatta caaagagtaa agctcattac   1440
tggcagtgtc tcttttttta agaaccgaga gcggctcaca ctctctgggc tggtcatttt   1500
tatgattatt tctttaattt attattattt ttttgcagct ctttccccca acttttgagc   1560
cgggtcaact ttctgagaat tgaaaagttc ccaaagtggt actgtttggg aacttctttc   1620
cctggctccc ctgatattcc gactgatgtt ttgggatttt tttcctctct gtttttttcc   1680
tgctgaaagc actatctcaa gtccgtcaca tcgcgctgtt tcaatccacc caaaggcgtc   1740
tgtgccagaa aggactccgc caagcccga gtttgagccc aggtttccgc agataacaaa   1800
tttctcgggt ttcttccgc agcttctctc ggcaactctc tcgcgcgggt gtaggtagcg   1860
gctgccgtat gacctgacct tggagtctc acattctagc tccacggccg gcgagctgcc   1920
ggctgatttg ctactttct gtctcctctg tcatactcta gttccttaca aactcttcac   1980
ggaccacggc ggcctcaacg agaagcgcaa gcagcggcgc atccgcacca ctttaccag   2040
tgcccagctc aaagagctgg aaagggctt cgcgagagact cactaccccg acatctacac   2100

```

-continued

togggaggag	ctggcctga	agatcgacct	cacagaggcg	cgagtccag	tacgcgcgcc	2160
tggaaaaccg	ccccctccg	ccgactggg	ccggggagg	gtgggtgag	ggcggtctg	2220
tgaaattcga	agtccctgg	cctcagtg	gaaggacct	gggccccat	gccgatcaga	2280
aatactggat	ttggtgtgg	tgtgcgttc	agagaggct	agagcgcac	ctcttgcat	2340
tttatttaca	gttgcaagt	gtttcccac	cgagcagaga	catgggggg	cttgggacgt	2400
ggatgagcga	tgcaatttc	gggacagg	gtgcctgtg	tggaaagtgt	gcagactttg	2460
ctcccgtatt	ataagtttt	ccttctccc	tcccgcgcc	caaaaaatg	cctcctaact	2520
caagtgcctt	taacctggc	ccatggcata	taggttcatt	ttcccggaa	ctgtgacttg	2580
cattagattt	gcaaagggt	tgtgacttc	tgaagtgca	gaacctgac	ttactccaac	2640
ctgttaaaaa	cagggtcgc	cacgagttg	ccacagcgc	tttttgggt	agccccgac	2700
cgagaagcgc	tgcgcacct	tgcacgctt	tccaggctc	aaggccggg	atgggcagcg	2760
gagcaaaacc	agagaggat	cctttctct	ttaccaatta	gagtttaact	ttagaactta	2820
ggcttagggg	tgaatggcg	gctcggggt	tgtcaagaa	gccgactgaa	cagaggccca	2880
ccaaaaaag	gccttccct	ttcgggtct	tctgggacct	gcggctttt	aaactctgcc	2940
gcaagccttc	atgtccctg	cgtgctcact	ccccctaaga	aagtttctc	gaaaatgcac	3000
agcaataaga	agcggtagc	ttggtggat	tgcgcgcgg	ggtgatcaca	gcgcatggg	3060
aggagggtgt	taaaacaag	cgaagtaga	cttgggccac	cctaaccggt	gcttttctt	3120
cccattttct	tctttctcc	cctgcttcac	cgtctctct	tcogtcttg	gccaggtgtg	3180
gttccagaac	cgcccgcca	agtttcgca	gcaggagcg	gcagcggcag	ccgcagcgg	3240
cgcgccaag	aacggtcct	cgggcaaaa	gtctgactt	tccagggacg	acgagagcaa	3300
agaggccaag	agcactgac	cggacagc	tgggggcca	ggtccaatc	ccaacccac	3360
ccccagctgc	ggggcgaat	gaggcggcg	cggcggccc	agcccggctg	gagctccgg	3420
ggcgcgggg	cccgggggc	cgggagcga	accggcga	ggcgcgag	cagcagcgg	3480
ggcgccgcg	gcccgggcg	cggcggcag	ggcagcggc	gcagctggg	gcttggtgc	3540
ggctggggg	cctggacaag	gctgggctc	cggccccgc	ccatcacct	ccatccgga	3600
ttcgtctggg	ggtcccctc	gcagcgtct	atcttcgct	caaagacca	acggtgcaa	3660
agccgcctta	gtgaagagc	gtatgttct	atctggaat	ctgcggcgc	ggcgcgcg	3720
gcgacagcgc	gcgagccag	gcccggcgg	gcgagtggt	gagcgggtg	gcccaaggct	3780
attgctgctg	ctgctgccat	ggctttttc	ttgaggcct	aaagtaatc	cgctaagaat	3840
aaagggaaaa	cggcgtcgc	ctcatttca	ccccactct	acccccttc	tcaacccca	3900
aacaaaacaa	acaaacttc	ctggcttcg	acctgcctg	ggcctcgag	cggggccagg	3960
gctccgcctg	ctgatgggg	gttgtgagc	gcgggcctg	gacgcgggg	actctcagg	4020
ggctgtgtct	gcgtgtcag	ttgtgtctg	ctcggggaat	gtgtgtctg	ggccaagca	4080
ggtgacagga	agagatggg	ggcctcaacc	aacttagtga	ctgttttaga	aaaaaaagc	4140
aaaaaagtaa	aaataaaaa	aaaaaagtt	gaaggcagaa	accattaaaa	aacaaaaagc	4200
caacaaccca	gaaagttta	aaaaacata	ggaaaaaaa	gacaaattaa	aggaggggct	4260
aggggagaag	ctgcagctg	agctgaagg	tcgatcttg	gaaccctaa	atccgctccc	4320
toctaacagc	acggattctc	ttggggctct	tcttcaggga	agagtaggga	cgccgttcca	4380

-continued

```

gcccccttc ctatcgtgtc cttgggttcg ggtcactgcg gcgacgactt gctcagactg 4440
tccccggcggc cggagtgact ttctcgcacc cccttgccctg tcccacctcg ctgaacacca 4500
tccccccatt agcgcacgag aacccccacac agttgcaact cccaaccccg aatccttgca 4560
gocgttcggc cctgaaagat gccctatcca tgagatgcct tttcatctgc aaactctgca 4620
aaatgtgtct catgtttcgc aactcttttt ttccccctcg ctcccgccta ccccgctggc 4680
atcttcttct tccaccagct tttactgaac tttttggcac tgctttggat tgggttcaat 4740
tgcagtcac gtaactggct gcagagaaat ctaccgagca aggaaaaggc acacacacac 4800
gtttgcaggg gtgtctcgtt ttgcatttct gttggaatga tcogaactgg actcacatcc 4860
tgtatgggtg atggactgta tattgagggt tccattcttc gcgcagtta gacatctctg 4920
ttttgattct ttgttgttgt ttttatttta aaaggcacia actctagata ttagttgaat 4980
gttagaggct taactttttt ggtgtcttct tacaactgtg ttctgtgact caattgtatc 5040
gtgttaatat cagtgcagac tgtctcctct acgtgaccgt ataatgtttt tctcgtcttg 5100
tagtctctat ggcgtgtctt tatgggtgaa taaggttctc acggggtcaa tcttttgtgt 5160
ttagagaggc cacggttcag acaatggtat atatttttgt tatcagggtc atgtctgtct 5220
gatttctttt ttttctctgt tggactatgt ttgtgaacat aattgtcata agttatgttt 5280
cagatttttg aatttattta tatgtgttat aatgaatgct tctattttaa agggaaatat 5340
ttctacatgt gcttatagtt ttccaagat gtaccattaa ctgattgtt gataataaaa 5400
accaaaagca agtctagcaa ttgaactctt ctttttcttg attctttttt tttttttttt 5460
tttgggttgg tcattgtttt ttttttttaa gttttttttt aaaaggat 5508

```

<210> SEQ ID NO 5

<211> LENGTH: 5493

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

```

cctgagtctc ttgaaagcac atttagctgc aagaagaaat gcaatttata agatgctttc 60
tctctgtctc tctttgtttt taacccccctc taaccagggtg tattttttaa gaaatccgct 120
tatcgttcac ataaaccccc ttggcccact tactctatgt tacagggcgc ctgagtcttg 180
ccaatgtccc agtcctttat aacatttcat gcacttcggg gggtaggctt gttgttaaat 240
tgagcgtgta acactcttac aaaacagggtt ttctatgaca tcaaggtttc ttctccctaa 300
ccgaggggga aaaaaagaag aggcagaaga agggggaaaa cacacacact atctcaattt 360
atgcctaagg tatatgatca gttaaaaagg cttaaagct cggggaaatt ggatcagggg 420
gaatcgtcac ccaactttca ttatttccaa gtagtgtgat tgaattaaag ggcagggagc 480
tggttagaag ggaggatcag gggctcgttg cgtaatggtg tggattataa ttctaattag 540
agatgcagga atcaatgata gggagggttg acagctcagt tcccagtgcc cagcccaata 600
gacggatgag ttattgtcat gtaaaaagcg ccagcaataa gaccaaccgc tttgctattg 660
tccaagtgga aagagccaag tttattatga ggactatag ctctagagac ctacagacaag 720
goatctcata ggaggotttt tcataaaact aggctctgct gtagtaagg aggccagttt 780
ggaggcaggc gttgagctgt gcacatctcc ccactccagc cacctctctc atatccatct 840
tttatttcat ttttccactt ggctgagcca tccagaacct tttcaatgta taaaatggaa 900

```

-continued

tattcttacc	tcaattcctc	tgccctacgag	tcctgtatgg	ctgggatgga	cacctcagac	960
ctggcttcag	cctatgctga	cttcagttcc	tcagccagc	ccagtggtt	ccagtataac	1020
ccgataagga	ccacttttgg	ggccacgtcc	ggctgccctt	ccctcacgcc	gggatcctgc	1080
agcctgggca	ccctcagga	ccaccagagc	agtccgtacg	ccgcaggtta	ggaccctcag	1140
ctttctcagc	ggaggaagcc	gcctttccgc	ccgtatatag	gaagccttga	ttgcatttga	1200
aaatggaaat	gtgttttagta	tttaccaaaac	gaaatttgct	tacacaaatg	aaagaattta	1260
tcacgtttga	agcgattgca	gggaggggta	attcaacttac	agggttacac	tatcctagtc	1320
acacccgaac	cgccccaaaa	attatcttaa	gctgccaaaa	tgataggcat	aatttattta	1380
ctttcgcgatg	agacgtatag	cttagaaaat	aattgaatta	caaagagtaa	agctcattac	1440
tggcagtgtc	tcttttttta	agaaccgaga	gcggctcaca	cttcttgggc	tggtcatttt	1500
tatgattatt	tctttaattt	attatatttt	ttttgcagct	ctttccccc	acttttgagc	1560
cggtcaact	ttctgagaat	tgaaaagtto	ccaaagtggg	actgtttgg	aacttctttc	1620
cctggctccc	ctgatattcc	gactgatgtt	ttgggatttt	tttcctctct	ggttttttcc	1680
tgctgaaagc	actatctcaa	gtccgtcaca	tcgcctgtt	tcaatccacc	caaaggcgt	1740
tgtgccagaa	aggactccgc	caagcccga	gtttgagccc	aggtttccgc	agataacaaa	1800
tttcctcgtt	ttcttcccgc	agcttctctc	ggcaactctc	tcgcgcgggt	gtaggtagcg	1860
gctgccgat	gacctgacct	tggagtctc	acattctagc	tcacagcccg	gcgagctgcc	1920
ggctgatttg	ctcactttct	gtctcctctg	tcatactcta	gttccttaca	aactcttcac	1980
ggaccacggc	ggcctcaacg	agaagcgcaa	gcagcggcgc	atccgcacca	ctttcaccag	2040
tgcccagctc	aaagagctgg	aaaggtctt	cgcgagact	cactacccc	acatctacac	2100
tcgggaggag	ctggccctga	agatcgacct	cacagaggcg	cgagtccag	tacgcgcgcc	2160
tggaaaccga	ccccgctccg	ccgcactggt	ccggggaggt	gtgggtgag	ggcgctgg	2220
tgaaattcga	agtcctggag	cctcgagtga	gaaggacct	gggcccctatg	gccgatcaga	2280
aatactgat	ttggtgtggc	tgtgcgttcg	agagaggctt	agagcgcacg	ctcttgcat	2340
tttatttaca	gttgcaagt	gtttcccacc	cgagcagaga	catggggggc	cttgggacgt	2400
ggatgagcga	tgcaatttcg	gggacaggaa	gtgcctgtgg	tggaaagtgt	gcagactttg	2460
ctcccgtatt	ataagttttt	ccttctcccc	tcccgcctcc	caaaaaaatg	cctcctaact	2520
caagtgcctt	taacctggcc	ccatggcata	taggttcatt	ttcccggaaa	ctgtgacttg	2580
cattagattt	gcaaaggttc	tgtgacttca	tgaaggtcaa	gaacctgac	ttactccaac	2640
ctgttaaaca	caggtgcgct	cacgagttgg	ccacagcgcc	tttttgggtg	agccccgac	2700
cgagaagcgg	tgcgcacct	tgacagcttt	tccaggtcca	aaggccgggg	atgggcagcg	2760
gagcaaacc	agagaggatc	ccttttctct	ttaccaatta	gagtttaact	ttagaaactta	2820
ggcttagggg	tgaatggcga	gctcggggct	tgctcaagaa	gcgcactgaa	cagaggocca	2880
caaaaaaag	gccttccctt	ttcgggtctt	tctgggacct	gcggcttttt	aaactctgcc	2940
gcaagccttc	atgtccctgg	cggtctcact	ccccctaaga	aagtttctcc	gaaaatgcac	3000
agcaataaga	agcggtagac	ttggtggatg	tgccgcggg	ggtgatcaca	gcgcatgggg	3060
aggaggtgt	taaaacaagc	cgaagtagaa	cttgggccc	cctaaccggt	gcttttcttt	3120
cccattttct	tctttctccc	cctgcttca	cgtctctct	tcogtcttg	gccaggtgtg	3180

-continued

gttcagaac	cgccgogcca	agtttcgcaa	gcaggagcgc	gcagcggcag	ccgcagcggc	3240
cgcgcccaag	aacggctcct	cgggcaaaaa	gtctgactct	tccagggacg	acgagagcaa	3300
agaggccaag	agcactgacc	cggacagcac	tgggggccca	ggtccaatc	ccaacccac	3360
ccccagctgc	ggggcgaatg	gaggcggcgg	cgggggccc	agcccggctg	gagctccggg	3420
ggcgcggggg	cccgggggccc	cgggagggca	accocggcaag	ggcgcgcgag	cagcagcggc	3480
ggcgcccgcg	gcagcggcag	cggcgcgagc	tggagccctg	gctgcggctg	ggggccctgg	3540
acaaggctgg	gctcccggcc	ccggcccat	cacctccatc	ccggattcgc	ttgggggtcc	3600
cttcggcagc	gtcctatctt	cgctccaaag	acccaacggt	gcaaaagccg	ccttagtgaa	3660
gagcagtatg	ttctgatctg	gaatctcgcg	gcggcggcgg	cggcgcgac	agcggcgag	3720
ccagggcccg	ggcgggcgag	tggcgagcgc	ggtaggccca	aggctattgt	cgctcgtgct	3780
gccatggctt	tttcattgag	ggcctaaagt	aatcgcgcta	agaataaagg	gaaaacggcg	3840
tgcacctcat	ttcaacccca	ctcctacccc	cttcctcaac	ccccaaaca	aacaaacaaa	3900
cttcocctggc	ttcgcacctg	cctggggcct	cgcagcgggg	ccagggctcc	gcctgctgat	3960
cgggggttgt	gagcagcgcg	gcctggacgc	ggggcactct	cagggggctg	tgtctcgtg	4020
tcagttttgt	tctgtctcgg	ggaatgtgtg	tctgtggccc	aagcaggtga	caggaagaga	4080
tggggggcct	caaccaactt	agtgaactgt	ttagaaaaaa	aagacaaaa	agtaaaaaata	4140
aaaacaaaa	agttggaagg	cagaaacctt	taaaaaaca	aaagccaaca	accagaaaag	4200
gtttaaaaaa	cataaggaaa	aaaaagacaa	attaaaggag	gggctagggg	agaagctgca	4260
gctggagctg	aaggctcgat	cttgtgaacc	cctaaatccg	ctccctccta	acagcacgga	4320
ttctcttggg	gctcttcttc	agggaaagag	agggacgccc	ttocagcccc	ccttcctatc	4380
gtgtccttgg	gttcgggtca	ctgcgcgac	gacttgctca	gactgtcccg	gcggccggag	4440
tgacttttct	gcaccccctt	gcctgtccca	cctcgtgaa	caccatcccg	ccattagcgc	4500
atcggaaacc	cacacagttg	caactcccaa	cccgaatct	ttgcagccgt	tcggccctga	4560
aagatgccct	atccatgaga	tgccctttca	tctgcaaaact	ctgcaaaatg	tgtctcatgt	4620
ttcgcacact	tttttttccc	cctcgtccc	gcctaccccg	tcggcatttt	cttcttcac	4680
cagcttttac	tgaacttttt	ggcactgctt	tggattgggg	tcaattgcag	tccacgtaac	4740
tggctgcaga	gaaatctacc	gagcaaggaa	aaggcacaca	cacacgtttg	caggggtgtc	4800
tcggtttgca	tttctgttgg	aatgatccga	actggactca	catcctgtat	ggtggatgga	4860
ctgtatattg	agggttccat	tcttcgcgca	gttttagacat	ctctgttttg	attctttgtt	4920
gttgttttta	ttttaaaagg	cacaaactct	agatattagt	tgaatgttga	ggctttaact	4980
ttttcgggtg	ctttctacaa	ctgtgttctg	tgactcaatt	gtatcgtgtt	aatatcagtg	5040
cagactgtct	cctctacgtg	accgtataat	gtttttctcg	tctttagtgc	tctatggcgt	5100
gtctttatgg	tgtaataagg	ttctcacggg	gtcaatcttt	tgtgtttaga	gaggccacgg	5160
ttcagacaa	ggtatataat	ttgttatca	ggtgcatgtc	tgtctgattt	cttttttttt	5220
cctgttggac	tatgtttgtg	aacataattg	tcataagtta	tgtttcagat	ttttgaattt	5280
atttatatgt	gttataatga	atgcttctat	ttaaaaggga	aatatttcta	catgtgotta	5340
tagttttcca	agagtgtacc	attaacttga	ttgttgataa	taaaaaccaa	aagcaagtct	5400
agcaattgaa	ctcttctttt	tcttgattct	tttttttttt	tttttttttg	gttggtcatt	5460

-continued

gttttttttt tttaagtttt tttttaaaag gat

5493

<210> SEQ ID NO 6

<211> LENGTH: 309

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

Met Tyr Lys Met Glu Tyr Ser Tyr Leu Asn Ser Ser Ala Tyr Glu Ser
 1 5 10 15

Cys Met Ala Gly Met Asp Thr Ser Ser Leu Ala Ser Ala Tyr Ala Asp
 20 25 30

Phe Ser Ser Cys Ser Gln Ala Ser Gly Phe Gln Tyr Asn Pro Ile Arg
 35 40 45

Thr Thr Phe Gly Ala Thr Ser Gly Cys Pro Ser Leu Thr Pro Gly Ser
 50 55 60

Cys Ser Leu Gly Thr Leu Arg Asp His Gln Ser Ser Pro Tyr Ala Ala
 65 70 75 80

Val Pro Tyr Lys Leu Phe Thr Asp His Gly Gly Leu Asn Glu Lys Arg
 85 90 95

Lys Gln Arg Arg Ile Arg Thr Thr Phe Thr Ser Ala Gln Leu Lys Glu
 100 105 110

Leu Glu Arg Val Phe Ala Glu Thr His Tyr Pro Asp Ile Tyr Thr Arg
 115 120 125

Glu Glu Leu Ala Leu Lys Ile Asp Leu Thr Glu Ala Arg Val Gln Val
 130 135 140

Trp Phe Gln Asn Arg Arg Ala Lys Phe Arg Lys Gln Glu Arg Ala Ala
 145 150 155 160

Ala Ala Ala Ala Ala Ala Ala Lys Asn Gly Ser Ser Gly Lys Lys Ser
 165 170 175

Asp Ser Ser Arg Asp Asp Glu Ser Lys Glu Ala Lys Ser Thr Asp Pro
 180 185 190

Asp Ser Thr Gly Gly Pro Gly Pro Asn Pro Asn Pro Thr Pro Ser Cys
 195 200 205

Gly Ala Asn Gly Gly Gly Gly Gly Gly Pro Ser Pro Ala Gly Ala Pro
 210 215 220

Gly Ala Ala Gly Pro Gly Gly Pro Gly Gly Glu Pro Gly Lys Gly Gly
 225 230 235 240

Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Ala Gly
 245 250 255

Gly Leu Ala Ala Ala Gly Gly Pro Gly Gln Gly Trp Ala Pro Gly Pro
 260 265 270

Gly Pro Ile Thr Ser Ile Pro Asp Ser Leu Gly Gly Pro Phe Gly Ser
 275 280 285

Val Leu Ser Ser Leu Gln Arg Pro Asn Gly Ala Lys Ala Ala Leu Val
 290 295 300

Lys Ser Ser Met Phe
 305

What is claimed is:

1. An isolated nucleic acid molecule comprising a nucleic acid sequence of SEQ ID NOS: 3, 4, or 5.

2. An isolated nucleic acid molecule comprising the nucleic acid sequence of SEQ ID NO: 1, wherein a nucleotide at position 2250 is substituted by a nucleotide base that is not adenine.

3. An isolated nucleic acid molecule comprising the nucleic acid sequence of SEQ ID NO: 1, wherein the nucleic acid molecule comprises one or more nucleotide substitutions from nucleotide positions 1-2250.

4. An isolated nucleic acid molecule comprising the nucleic acid sequence of SEQ ID NO: 1, wherein the nucleic acid molecule comprises one or more nucleotide deletions from nucleotide positions 1-2250.

5. An isolated nucleic acid molecule comprising the nucleic acid sequence of SEQ ID NO: 1, wherein the nucleic acid molecule comprises one or more nucleotide insertions from nucleotide positions 1-2250.

6. An isolated nucleic acid molecule comprising the nucleic acid sequence of SEQ ID NO: 1, wherein the nucleic acid molecule comprises a nucleotide substitution at nucleotide position 1129.

7. An isolated nucleic acid molecule comprising the nucleic acid sequence of SEQ ID NO: 1, wherein the nucleic acid molecule comprises a nucleotide substitution at nucleotide position 2304.

8. An isolated nucleic acid molecule comprising the nucleic acid sequence of SEQ ID NO: 1, wherein a nucleotide at position 3494 is substituted by a nucleotide base that is not adenine.

9. An isolated nucleic acid molecule comprising the nucleic acid sequence of SEQ ID NO: 1, wherein nucleotides from nucleotide positions 3495 to 3499 are deleted.

10. An isolated nucleic acid molecule comprising at least 6 contiguous nucleotides of SEQ ID NO: 1 including a nucleotide substitution at position 2250 by a nucleotide base that is not adenine.

11. An isolated nucleic acid molecule comprising at least 6 contiguous nucleotides of SEQ ID NO: 1 including a nucleotide substitution at position 3494 by a nucleotide base that is not adenine.

12. An isolated nucleic acid molecule comprising at least 6 contiguous nucleotides of SEQ ID NO: 1 including one or more nucleotides at positions 3495 to 3499.

13. An isolated nucleic acid molecule comprising at least 6 contiguous nucleotides of SEQ ID NO: 1 including a nucleotide substitution at position 2250 by a nucleotide base that is not adenine and that hybridizes to a nucleic acid molecule consisting of: (a) the nucleic acid sequence of SEQ ID NO: 1; or (b) a nucleotide sequence that encodes the amino acid sequence of SEQ ID NO: 2, under moderately stringent conditions, wherein the moderately stringent conditions comprise hybridization in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate, 1 mM EDTA at 65° C., and washing in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate, 1 mM EDTA at 65° C., and washing in 0.2×SSC/0.1% sodium dodecyl sulfate at 42° C., or the complement thereof.

14. An isolated nucleic acid molecule comprising at least 6 contiguous nucleotides of SEQ ID NO: 1 including a nucleotide substitution at position 3493 by a nucleotide base that is not adenine and that hybridizes to a nucleic acid molecule consisting of: (a) the nucleic acid sequence of SEQ ID NO: 1; or (b) a nucleotide sequence that encodes the

amino acid sequence of SEQ ID NO: 2, under moderately stringent conditions, wherein the moderately stringent conditions comprise hybridization in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate, 1 mM EDTA at 65° C., and washing in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate, 1 mM EDTA at 65° C., and washing in 0.2×SSC/0.1% sodium dodecyl sulfate at 42° C., or the complement thereof.

15. An isolated nucleic acid molecule comprising at least 6 contiguous nucleotides of SEQ ID NO: 1 including one or more nucleotides at positions 3495 to 3499 and that hybridizes to a nucleic acid molecule consisting of: (a) the nucleic acid sequence of SEQ ID NO: 1; or (b) a nucleotide sequence that encodes the amino acid sequence of SEQ ID NO: 2, under moderately stringent conditions, wherein the moderately stringent conditions comprise hybridization in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate, 1 mM EDTA at 65° C., and washing in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate, 1 mM EDTA at 65° C., and washing in 0.2×SSC/0.1% sodium dodecyl sulfate at 42° C., or the complement thereof.

16. A vector comprising the nucleic acid molecule of claim 1, 2, 13, 14, or 15.

17. An expression vector comprising the nucleic acid molecule of claim 1, 2, 13, 14, or 15 operatively associated with a regulatory nucleic acid controlling the expression of the nucleic acid molecule in a host cell.

18. A host cell genetically engineered to express the nucleic acid molecule of claim 1, 2, 13, 14, or 15.

19. A host cell genetically engineered to express the nucleic acid molecule of claim 1, 2, 13, 14, or 15, operatively associated with a regulatory nucleic acid controlling the expression of the nucleic acid molecule in the host cell.

20. A method for producing a polypeptide comprising introducing into a cell an expression vector comprising the nucleic acid molecule of claim 19 operatively associated with a regulatory nucleic acid controlling the expression of the nucleic acid molecule in the cell; and culturing the cell such that the polypeptide encoded by the nucleic acid molecule is produced.

21. A method for producing a polypeptide comprising culturing a cell genetically engineered to express the nucleic acid molecule of claim 19 such that the polypeptide encoded by the molecule is produced.

22. A peptide comprising at least 6 consecutive amino acid residues of SEQ ID NO: 6 including one or more amino acids at positions 245 to 249.

23. A polypeptide comprising the amino acid sequence of SEQ ID NO: 6.

24. A method for screening a subject predisposed or susceptible to HSCR associated with a genetic polymorphism in the PHOX2B gene, said method comprising:

- (a) providing a biological sample from the subject; and
- (b) testing the sample for the presence of a nucleotide mutation at nucleotide position 2250 of SEQ ID NO: 1, wherein the presence of the mutation indicates that the subject is genetically predisposed to HSCR.

25. A method for screening a subject predisposed or susceptible to HSCR associated with a genetic polymorphism in the PHOX2B gene, said method comprising:

- (a) providing a biological sample from the subject; and
- (b) testing the sample for the presence of one or more nucleotide mutations from nucleotide positions 1-2250

of SEQ ID NO: 1, wherein, the presence of the mutation indicates that the subject is genetically predisposed to HSCR.

26. A method for diagnosing a subject with increased risk of HSCR associated with a genetic polymorphism in the PHOX2B gene, said method comprising:

- (a) providing a biological sample from the subject; and
- (b) testing the sample for the presence of a nucleotide mutation at nucleotide position 2250 of SEQ ID NO: 1, wherein, the presence of the mutation indicates that the subject is genetically predisposed to HSCR.

27. A method for diagnosing a subject with increased risk of HSCR associated with a genetic polymorphism in the PHOX2B gene, said method comprising:

- (a) providing a biological sample from the subject; and
- (b) testing the sample for the presence of one or more nucleotide mutations from nucleotide positions 1-2250 of SEQ ID NO: 1, wherein, the presence of the mutation indicates that the subject is genetically predisposed to HSCR.

28. The method of claim 24 or 26 wherein the subject is heterozygous for the PHOX2B mutation.

29. The method of claim 24 or 26 wherein the subject is homozygous for the PHOX2B mutation.

30. The method of claim 24 or 26 comprising amplifying a fragment of the PHOX2B gene comprising two or more nucleotides including a nucleotide at position 2250.

31. The method of claim 30 wherein the fragment is amplified using PCR primers comprising nucleotide sequence of SEQ ID NOS: 9 and 10.

32. The method of claim 24 or 26, wherein the biological sample is selected from the group consisting of blood, saliva, amniotic fluid, and urine.

33. The method according to claim 24 or 26, wherein the subject is a human.

34. The method according to claim 24 or 26, further comprising testing the sample for the presence of mutation in one or more genes selected from the group consisting of RET, EDNRB, EDN3 and GDNF.

35. A method for treating a subject diagnosed as having HSCR comprising modulating the level of PHOX2B polypeptide having amino acid sequence of SEQ ID NO: 2 in the subject.

36. An animal model for HSCR comprising a mammal comprising the nucleic acid sequence of SEQ ID NO: 1, wherein a nucleotide at position 2250 is substituted by a nucleotide base that is not adenine.

37. A method for screening potential therapeutic agents for modulating the activity of PHOX2B gene, comprising:

(a) contacting a test compound with a cell or cell lysate containing PHOX2B gene product; and

(b) detecting the activity of the PHOX2B gene.

38. An oligonucleotide of at least 8 nucleotides in length selected from nucleotides 1-46, 48-123; 120-369; 365-394; 390-540; 538-646; 643-1004; 1001-1080; 1083-1109; 1106-1304; 1301-1366; 1363-1386; 1389-1514; 1516-1778; 1773-1917; 1921-2010; 2051-2146; 2154-2209; 2234-2368; 2367-2422; 2420-2464; 2465-2491; 2488-2568; 2872-2901; 2902-2934; 2936-2954; 2449-3001; 3000-3042; 3420-3435; 3451-3708; 3703-3754; 3750-3770; 3774-3840; 3840-3962; 3964-3978; 3974-3992; 3990-4157; 4153-4251; 4257-4282; 4284-4321; 4316-4333; 4337-4391; 4386-4400; 4398-4436; 4444-4547; 4572-4714; 4709-4777; 5165-5397; 5394-5508 of SEQ ID NO: 1, 3 or 4.

39. An oligonucleotide pair comprising an oligonucleotide of claim 38 and an oligonucleotide of at least 8 nucleotides in length selected from SEQ ID NO: 1, 3, or 4.

40. A kit for detection of a polymorphism in the PHOX2B gene in a subject sample, the kit comprising at least one oligonucleotide of at least 8 nucleotides in length selected from the group consisting of SEQ ID NOS: 1, 3, or 4, wherein the oligonucleotide is used to amplify a fragment of the PHOX2B DNA or RNA in a subject sample.

41. A method of determining the source of a first biological sample from a subject compared to a second biological sample, said method comprising:

- (a) providing a first biological sample from the subject;
- (b) testing the first biological sample for the presence of a mutation of the PHOX2B gene at nucleotide positions 2250, 3493, or 3495-3499 of SEQ ID NO: 1;
- (c) testing the second biological sample for the presence of a mutation of the PHOX2B gene at nucleotide positions 2250, 3493, or 3495-3499 of SEQ ID NO: 1; and
- (d) comparing the mutations present in step (b) and (c); wherein the presence of identical mutation in the first biological sample compared to the second biological sample indicates that the first and second biological samples are obtained from an identical source.

42. A method for producing a polypeptide comprising culturing a host cell genetically engineered to express the nucleic acid molecule of claim 1, 2, 13, 14, or 15 operatively associated with a regulatory nucleic acid and controlling the expression of the nucleic acid molecule in the host cell such that the polypeptide encoded by the molecule is produced.

* * * * *