

Title: Re-examining the factor structure and psychometric properties of the Mini-Mental Adjustment to Cancer Scale in a sample of 364 Chinese cancer patients

Authors: Ted C. T. Fong*, Rainbow T. H. Ho

Affiliation: Centre on Behavioral Health and Dept of Social Work and Social Administration,
The University of Hong Kong, Hong Kong

Corresponding author: *Ted Chun Tat Fong, Centre on Behavioral Health, 2/F, Hong Kong
Jockey Club Building for Interdisciplinary Research, 5 Sassoon Road, Pokfulam, HKSAR
Phone no: 852-2831 5174; E-mail: ttaatt@hku.hk; Fax no: 852-2816 6710

Keywords: mental adjustment; cancer; Mini-MAC; factor structure; exploratory factor analysis;
Chinese

Word count (exc. figures/tables): 2,441; number of tables: 5; number of figures: 0

Conflict of Interest: There are no potential conflicts of interest.

Abstract

Purpose: The Mini-Mental Adjustment to Cancer Scale (Mini-MAC) is widely used to evaluate cancer patients' psychological responses. Validation studies of the scale have shown methodological shortcomings and inconsistency in the factor solutions. The aim of this study was to examine the factor structure and psychometric properties of the Mini-MAC.

Methods: A large sample of 364 Chinese patients with breast or colorectal cancer completed the Mini-MAC and psychosocial measures (general health, perceived stress, anxiety, and depression). Exploratory factor analyses examined the relative fit of two- to six-factor models using robust weighted least square estimation and oblique target rotation. Convergent validity was evaluated via correlations between the Mini-MAC factor scores and the psychosocial outcomes.

Results: The five-factor model showed the best model fit and largely replicated the original Mini-MAC's helpless/hopeless (HH), anxious preoccupation (AP), fighting spirit (FS), fatalism (FA), and cognitive avoidance (CA) subscales. The five factors had acceptable reliability (Cronbach's $\alpha = .67 - .88$) and 4-month test-retest reliability ($r = .45 - .64$). HH, AP, and CA were positively associated with the psychosocial outcomes ($r = .19 - .60$). Modest and negative correlations were found between the psychosocial outcomes and FS and FA.

Conclusions: The results support the Mini-MAC's original five-factor structure with satisfactory reliability and convergent validity. The results demonstrate that the Mini-MAC is a valid measure for assessing psychological responses in cancer patients.

Keywords: mental adjustment; cancer; Mini-MAC; factor structure; exploratory factor analysis; Chinese

Introduction

Mental adjustment to cancer is defined as patients' cognitive appraisals of the life-threatening condition and their behavioral coping responses [1]. Coping responses have been associated with various psychosocial outcomes in cancer patients [2,3].

Valid and reliable assessment of the adjustment response to cancer is vital for assessing the risk of later psychological adjustment difficulties and for evaluating how the illness affects clinical practice. The Mental Adjustment to Cancer Scale (MAC) is a 40-item self-rating measure of adjustment styles in cancer patients [4]. It assesses adjustment along five dimensions: helpless/hopeless, anxious preoccupation, fighting spirit, fatalism, and cognitive avoidance. Later, Watson *et al.* [5] included additional items to evaluate cognitive avoidance and performed exploratory factor analysis (EFA) in a larger sample of 573 cancer patients, resulting in the Mini-Mental Adjustment to Cancer Scale (Mini-MAC) as a 29-item shortened version.

The Mini-MAC, often adopted in preference to the MAC, is widely used to measure psychological responses to cancer in research and clinical settings. Researchers have translated the Mini-MAC into Chinese [6], Italian [7], Greek [8], Korean [9], and Norwegian [10] and examined its psychometric properties in different contexts. The results of these validation studies indicate that the scale has adequate test-retest reliability and convergent validity with psychosocial outcomes. Despite the proposed five-factor structure for the Mini-MAC [5], various factor structures have emerged in the validation studies (Table 1), including three- [6], four- [9,10], and five-factor solutions [7,8]. Four of the validation studies [6,7,9,10] obtained the results via principal component analysis and varimax rotation. Principal component analysis fails to differentiate between shared and unique variance [11], and thus is a biased estimator for factor analysis [12]. The assumption of uncorrelated factors in varimax rotation is often unrealistic and can result in misleading factor structures [12].

1 The inappropriate use of the extraction and rotation methods casts considerable
2 doubts over the credibility of the results.
3

4
5 [Insert Table 1 about here]
6

7 The most recent validation study [13] adopted more psychometrically sound
8 criteria (principal axis factoring and oblimin rotation) for the Mini-MAC and found
9 support for a four-factor structure. In particular, oblique rotation provides a precise
10 and realistic representation of the factor structure. However, similar to previous
11 studies [6,8,9], this study had a small sample size ($n \leq 200$), which was sub-optimal. A
12 larger sample size is preferable to produce a more accurate and robust factor solution
13 [11]. Overall, the methodological limitations of existing validation studies point to the
14 need for further psychometric analysis of the scale. The present study aimed to
15 systematically examine the factor structure of the Mini-MAC among a large sample of
16 364 Chinese cancer patients. The psychometric properties in terms of internal
17 consistency, test-retest reliability, and convergent validity were scrutinized for the best
18 factor solution.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

39 **Methods**

40 **Sample**

41 The data used in this study were taken from a larger data set collected for a clinical
42 psychotherapy trial among cancer patients over a 24-month period. Eight hundred and
43 seventy-two cancer patients who had their treatment completed were invited to join
44 the study via mail and phone calls. In the invitation mail, the study's purpose,
45 procedures, and potential risks were explained. The exclusion criteria included a
46 recurrent cancer diagnosis, inability to understand Chinese, or life expectancy of less
47 than 6 months. A total of 364 cancer patients were recruited post treatment from four
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 community cancer support centers in Hong Kong, indicating a response rate of 41.7%.
2
3 The participants completed a paper-and-pencil questionnaire at baseline and a
4
5 subsample of 291 patients provided follow-up data 4 months later. Ethical approval
6
7 was obtained from the local institutional review board and the participants provided
8
9 written informed consent.
10

11 Measures

12
13 The Chinese Mini-MAC [6] is a 29-item, 4-point self-report instrument for assessing
14
15 the cognitive and behavioral responses of cancer patients with five proposed factors:
16
17 helpless/hopeless (HH, 8 items), anxious preoccupation (AP, 8 items), fighting spirit
18
19 (FS, 4 items), fatalism (FA, 5 items), and cognitive avoidance (CA, 4 items).
20
21

22 Satisfactory reliability was found for HH ($\alpha = .83 - .87$), AP ($\alpha = .81 - .88$), and CA (α
23
24 = .65 - .82) while marginal reliability was found for FS ($\alpha = .21 - .76$) and FA ($\alpha = .52$
25
26 - .71) in previous studies [6-10]. A series of psychosocial outcomes, namely general
27
28 health, perceived stress, anxiety, and depression, was evaluated in this study. General
29
30 health was assessed by the 12-item Chinese General Health Questionnaire [14]. Item
31
32 responses were rated using a 2-point format, with the total scale score ranging from 0
33
34 to 12. Perceived stress was measured by the 10-item Chinese Perceived Stress Scale
35
36 [15]. Item responses were rated using a 5-point format, with the total scale score
37
38 ranging from 0 to 40. Anxiety and depression were assessed using the 14-item
39
40 Chinese Hospital Anxiety and Depression Scale [16]. Item responses were rated using
41
42 a 4-point format, with the total scores for anxiety (7 items) and depression (7 items)
43
44 ranging from 0 to 21. Higher scores denoted higher levels of distress for the
45
46 psychosocial outcomes. Cronbach's alpha (α) coefficient values were satisfactory for
47
48 the measurement scales of general health ($\alpha = .875$), perceived stress ($\alpha = .788$),
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 anxiety ($\alpha = .848$), and depression ($\alpha = .742$).
2
3
4

5 Data analysis 6

7 EFA was performed using Mplus version 7.11 [17] to investigate the factor structure
8 of the Mini-MAC. In preliminary analyses, 13 of the 29 Mini-MAC items exhibited
9 floor and ceiling effects, with the response proportion at both tails exceeding 30%.

10 Given the non-normality, factor extraction was conducted on the categorical
11 indicators using robust weighted least square estimators [18,19] and oblique target
12 rotation [20]. Popular criteria such as Kaiser's eigenvalue greater than 1 and scree
13 tests are prone to factor overextraction. To decide the optimal dimensionality, we
14 systematically specified five EFA models (two- to six-factor models) and compared
15 their model fit.
16
17
18
19
20
21
22
23
24
25
26
27

28 Model fit was assessed based on the following criteria on the goodness-of-fit
29 indices [21]: comparative fit index (CFI) $\geq .95$; Tucker-Lewis index (TLI) $\geq .95$; root
30 mean square error of approximation (RMSEA) $\leq .06$; and standardized root mean
31 square residual (SRMR) $\leq .08$. Chi-square (χ^2) difference tests were conducted to
32 compare the fit of adjacent EFA models (e.g. two- versus three-factor models). Model
33 selection was based on theoretical considerations with reference to the original
34 five-factor structure. Factor loadings greater than 0.40 were considered major and
35 practically significant, and items without any major factor loadings were removed
36 from the model.
37
38
39
40
41
42
43
44
45
46
47
48
49

50 The best-fitting factor model was then evaluated using the 4-month follow-up
51 data for model validation. The internal consistency for each factor was assessed by
52 Cronbach's α . Test-retest reliability was evaluated by correlating the derived factor
53 scores at baseline with those at the 4-month follow-up. The convergent validity of the
54 Mini-MAC was examined through correlation analyses of the Mini-MAC factor
55
56
57
58
59
60
61
62
63
64
65

1 scores with general health, perceived stress, anxiety, and depression. The rate of
2 missing data was minimal (no more than 2% of any item response). The statistical
3 significance level was set at .01.
4
5
6
7

8 9 10 **Results**

11 Sample characteristics

12
13 The study sample comprised patients with breast ($n = 196$) or colorectal ($n = 168$)
14 cancer. Of the 364 participants, 69.5% were female and the sample mean age was 53.5
15 years ($SD = 11.0$). The majority of the participants was married (79.7%), had
16 completed secondary education (56.3%), and had received chemotherapy (69.0%) and
17 surgery (65.7%). Compared with the participants who provided follow-up data, the
18 participants who did not respond at follow-up were significantly more likely to be
19 male, have colorectal cancer, and have a lower level of anxiety at baseline.
20
21
22
23
24
25
26
27
28
29
30
31

32 Factor structure

33
34 As shown in the goodness-of-fit indices in Table 2, mediocre fit was found for the
35 two- and three-factor models, and the four-factor model displayed a marginally
36 acceptable fit to the data. In contrast, the five-factor model showed an adequate model
37 fit and fitted the data significantly better than the four-factor model via the chi-square
38 difference test ($\Delta\chi^2 = 166.2$, $\Delta df = 25$, $p < .01$). The six-factor model showed a
39 significantly better fit than the five-factor model ($\Delta\chi^2 = 81.5$, $\Delta df = 24$, $p < .01$).
40
41 However, the sixth factor consisted of only one major factor loading from item 21 and
42 was not correlated with any of the other five factors, thus providing little incremental
43 value over the five-factor model.
44
45
46
47
48
49
50
51
52
53
54
55
56

57 [Insert Table 2 about here]
58

59 The eigenvalues of the five factors were 9.760, 4.386, 1.585, 1.469, and 1.342.
60
61
62
63
64
65

1 All 29 items showed major factor loadings, ranging from .401 to .820 (Table 3). The
2 factor loading pattern was congruent with the original five-factor structure [5] with
3 two exceptions. Item 7 had a major factor loading on HH rather than AP and Item 26
4 loaded significantly onto both FS and CA. The five-factor model provided an
5 acceptable fit to the data at follow-up ($\chi^2 = 519.0$, $df = 271$, $p < .01$, CFI = .975, TLI
6 = .962, RMSEA = .056, and SRMR = .038). In view of the findings, the five-factor
7 model was chosen to represent the factor structure and further tests of reliability and
8 convergent validity were conducted.
9

10 [Insert Table 3 about here]
11
12
13
14
15
16
17
18
19
20
21
22
23

24 Reliability and convergent validity

25 Cronbach's α coefficient, interfactor correlations, and test-retest reliability were
26 assessed for the five Mini-MAC factors (Table 4). HH and AP were found to exhibit
27 high reliability ($\alpha > .85$) while acceptable reliability was found for FS, FA, and CA (α
28 ~ .70). Good test-retest reliability was found for the factors across the 4-month
29 interval ($r = .449 - .640$). CA was positively correlated with the other factors except
30 FS ($r = .188 - .449$). FA were positively associated with FS ($r = .548$) and negatively
31 correlated with HH ($r = -.221$).
32
33
34
35
36
37
38
39
40
41
42

43 [Insert Table 4 about here]
44
45

46 Regarding the convergent validity of the Mini-MAC (Table 5), positive and
47 moderately strong correlations were found for HH and AP with the psychosocial
48 outcomes ($r = .431 - .601$). Higher levels of HH and AP were linked to worse general
49 health and higher levels of perceives stress, anxiety, and depression. Both FS and FA
50 were negatively and modestly correlated with the psychosocial outcomes ($r = -.175 -$
51 $-.328$). Higher levels of FS and FA were associated with better health and lower levels
52 of psychological distress. CA had positive and weak correlations with the outcomes (r
53
54
55
56
57
58
59
60
61
62
63
64
65

1 = .194 - .277), suggesting higher level of CA to be linked to worse general and mental
2
3 health.

4
5 [Insert Table 5 about here]
6
7

8 9 10 **Discussion**

11
12 Given the inconsistency in the factor solutions and the methodological shortcomings
13 of previous validation studies of the Mini-MAC, we performed a systematic
14 investigation of its factor structure using a robust weighted least square estimator with
15 an oblique target rotation. Comprehensive analyses and comparisons of various factor
16 structures lend support to the five-factor solution as the underlying factor structure.
17
18 The five-factor model, which provides a significantly better fit than the three- and
19 four-factor models, largely replicates the original five-factor structure proposed by
20 Watson *et al.* [5]. The modest to moderate inter-factor correlations support adequate
21 discriminant validity for the structure.
22
23

24
25 In our five-factor model, all 29 items contributed significantly to a clear factor
26 structure, with only Item 26 (“I make a positive effort not to think about my illness”)
27 loading significantly on both FS and CA. Item 7 (“It is a devastating feeling”), which
28 was originally hypothesized as an AP indicator, may better measure the dimension of
29 HH. Despite these minor disparities, HH and AP showed good reliability and
30 convergent validity with general health, perceived stress, anxiety, and depression.
31
32 These results are in line with previous findings and support the two factors as good
33 indicators of poor adjustment. Despite the recent concerns [10,13] over the utility of
34 FS due to its low reliability and lack of convergent validity, this subscale showed
35 adequate reliability and convergent validity as measured against the psychosocial
36 outcomes.
37
38

39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60 Cronbach’s α for the FA subscale (.674) was slightly better than those in the
61
62
63
64
65

1 original study [5] and other validation studies [7,9,10,13]. Researchers [6,9,10] have
2
3 posited that fatalism may reflect not only the acceptance of an unavoidable fate, but
4
5 also an optimistic and active form of coping. Similarly, Fitzpatrick *et al.* [22] found
6
7 FA to be positively correlated with spirituality in a sample of breast cancer patients. In
8
9 the present study, FA was negatively associated with general health, perceived stress,
10
11 anxiety, and depression. This finding suggests that fatalism does not necessarily mean
12
13 resignation coping, but could enable patients to endure the situation better, thereby
14
15 reducing their level of distress. Given the distinct feature of this subscale in assessing
16
17 potentially positive coping styles, future studies could adopt this subscale and
18
19 examine its predictive validity on psychosocial outcomes.
20
21
22
23

24 The CA subscale showed satisfactory reliability and was positively associated
25
26 with HH, AP, FA, and the psychosocial outcomes. These results, which are generally
27
28 in line with the findings of the original study [5], appear to suggest that CA is an
29
30 indicator of poor adjustment in Chinese cancer patients. As Bredal [10] points out, CA
31
32 is not necessarily detrimental to patients. The CA items have been described as
33
34 distractions and mental disengagements of active thinking about the illness and can be
35
36 viewed as emotion-focused coping strategies [6]. The effective use of such strategies
37
38 can assist in affective regulation and problem-focused coping in certain situations.
39
40
41
42

43 From an analytical perspective, it should be noted that eight of the ten
44
45 inter-factor correlations were significant ($r = .188 - .677$) in our factor solution,
46
47 confirming the use of an oblique rotation. We agree with the view of
48
49 Hulbert-Williams *et al.* [13] that orthogonal rotations (e.g. varimax rotation) should
50
51 no longer be used with these data to avoid distorting the underlying factor structure.
52
53 Despite its common use in previous studies, principal component analysis has been
54
55 shown to be a biased and outdated extraction method in the psychometric literature
56
57 [11,12,19], and its future use is highly discouraged. It is recommended that
58
59
60
61
62
63
64
65

1 researchers adopt more up-to-date estimators such as principal axis factoring or
2 maximum likelihood for continuous indicators, or robust weighted least square
3 estimation for categorical indicators.
4
5

6
7 A limitation of the present study is that it was based on Chinese patients with
8 breast or colorectal cancer. The findings may not be generalizable to patients with
9 other types of cancer or in other cultural contexts. The present study was based on
10 self-report data and may be subject to common method bias. Future longitudinal
11 studies could elucidate the predictive validity of the scale with psychosocial
12 outcomes. Despite these limitations, the present study is the first of its kind to apply
13 systematic factor analysis in evaluating the scale's factor structure. The results
14 demonstrate the original five-factor structure with satisfactory reliability and
15 convergent validity. Confirmatory factor analysis in further research is recommended
16 to verify the psychometric properties of the Mini-MAC and to assess the measurement
17 invariance of the scale across gender, cancer type, and cultural context.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34
35
36 **Acknowledgements:** We would like to thank Hong Kong Cancer Fund, Queen Mary
37 Hospital, Pamela Youde Nethersole Eastern Hospital, Dr. M.Y. Luk, Ms. Amy Choi,
38 Ms. Venus Tang, Mr. Ryder Chan, and all the study participants for their help in data
39 collection.
40
41
42
43
44

45
46
47 **Conflict of Interest:** There are no potential conflicts of interest.
48
49
50

51 52 53 54 55 56 57 58 59 60 61 62 63 64 65

1. Greer S, Moorey S, Watson M (1989) Patients' adjustment to cancer: the Mental Adjustment to Cancer (MAC) Scale vs clinical ratings. *J Psychosom Res* 33

1 (3):373-377. doi:10.1016/0022-3999(89)90027-5

2
3 2. Seok JH, Choi WJ, Lee YS, Park CS, Oh YJ, Kim JS, Chang HS (2013)

4 Relationship between Negative Mental Adjustment to Cancer and Distress in Thyroid

5 Cancer Patients. *Yonsei Med J* 54 (3):658-664. doi:10.3349/ymj.2013.54.3.658

6
7 3. Johansson M, Ryden A, Finizia C (2011) Mental adjustment to cancer and its

8 relation to anxiety, depression, HRQL and survival in patients with laryngeal cancer -

9 A longitudinal study. *Bmc Cancer* 11. doi:10.1186/1471-2407-11-283

10
11 4. Watson M, Greer S, Young J, Inayat Q, Burgess C, Robertson B (1988)

12 Development of a questionnaire measure of adjustment to cancer: the MAC scale.

13
14 *Psychol Med* 18 (1):203-209

15
16 5. Watson M, Law M, Dossantos M, Greer S, Baruch J, Bliss J (1994) The Mini-MAC

17 - Further development of the Mental Adjustment to Cancer Scale. *Journal of*

18
19 *Psychosocial Oncology* 12 (3):33-46. doi:10.1300/j077v12n03_03

20
21 6. Ho SMY, Fung WK, Chan CLW, Watson M, Tsui YKY (2003) Psychometric

22 properties of the Chinese version of the Mini-Mental Adjustment to Cancer

23
24 (Mini-MAC) scale. *Psychooncology* 12 (6):547-556. doi:10.1002/pon.672

25
26 7. Grassi L, Buda P, Cavana L, Annunziata MA, Torta R, Varetto A (2005) Styles of

27 coping with cancer: The Italian version of the mini-mental adjustment to cancer

28
29 (Mini-MAC) scale. *Psychooncology* 14 (2):115-124. doi:10.1002/pon.826

30
31 8. Anagnostopoulos F, Kolokotroni P, Spanea E, Chrysoschoou M (2006) The

32 mini-mental adjustment to cancer (Mini-MAC) scale: Construct validation with a

33
34 Greek sample of breast cancer patients. *Psychooncology* 15 (1):79-89.

35
36 doi:10.1002/pon.924

37
38 9. Kang JI, Chung HC, Kim SJ, Choi HJ, Ahn JB, Jeung HC, Namkoong K (2008)

39 Standardization of the Korean version of Mini-Mental Adjustment to Cancer

40
41 (K-Mini-MAC) scale: Factor structure, reliability and validity. *Psychooncology* 17

1 (6):592-597. doi:10.1002/pon.1277

2 10. Bredal IS (2010) The Norwegian version of the Mini-Mental Adjustment to

3 Cancer Scale: Factor structure and psychometric properties. *Psychooncology* 19

4 (2):216-221. doi:10.1002/pon.1564

5 11. Costello AB, Osborne JW (2005) Best practices in exploratory factor analysis:

6 four recommendations for getting the most from your analysis. *Practical Assessment,*

7 *Research and Evaluation* 10 (7):1-9

8 12. Fabrigar LR, Wegener DT, MacCallum RC, Strahan EJ (1999) Evaluating the use

9 of exploratory factor analysis in psychological research. *Psychol Meth*, 4 (3):272-299.

10 doi:10.1037/1082-989x.4.3.272

11 13. Hulbert-Williams NJ, Hulbert-Williams L, Morrison V, Neal RD, Wilkinson C

12 (2012) The Mini-Mental Adjustment to Cancer Scale: Re-analysis of its psychometric

13 properties in a sample of 160 mixed cancer patients. *Psychooncology* 21 (7):792-797.

14 doi:10.1002/pon.1994

15 14. Pan PC, Goldberg D (1990) A comparison of the validity of GHQ-12 and CHQ-12

16 in Chinese primary care patients in Manchester. *Psychol Med* 20 (4):931-940

17 15. Leung DYP, Lam TH, Chan SSC (2010) Three versions of Perceived Stress Scale:

18 Validation in a sample of Chinese cardiac patients who smoke. *Bmc Public Health*

19 10:513. doi:51310.1186/1471-2458-10-513

20 16. Fong TCT, Ho RTH (2013) Factor analyses of the Hospital Anxiety and

21 Depression Scale: a Bayesian structural equation modeling approach. *Qual Life*

22 *Res*:1-7. doi:10.1007/s11136-013-0429-2

23 17. Muthén LK, Muthén B (1998-2013) *Mplus user's guide*. 7th edn. Muthen &

24 Muthen, Los Angeles, CA

25 18. Beauducel A, Herzberg PY (2006) On the performance of maximum likelihood

26 versus means and variance adjusted weighted least squares estimation in CFA.

1 Structural Equation Modeling 13 (2):186-203. doi:10.1207/s15328007sem1302_2

2
3 19. Schmitt TA (2011) Current Methodological Considerations in Exploratory and
4
5 Confirmatory Factor Analysis. Journal of Psychoeducational Assessment 29
6
7 (4):304-321. doi:10.1177/0734282911406653
8
9

10 20. Browne MW (2001) An overview of analytic rotation in exploratory factor
11
12 analysis. Multivariate Behav Res 36 (1):111-150. doi:10.1207/s15327906mbr3601_05
13

14 21. Hu LT, Bentler PM (1999) Cutoff criteria for fit indexes in covariance structure
15
16 analysis: conventional criteria versus new alternatives. Structural Equation Modeling
17
18 6 (1):1-55. doi:10.1080/10705519909540118
19
20

21 22. Fitzpatrick C, Levine EG, Heide F, Zelman D, Targ E (2000) Re-examining the
22
23 construct of fatalism in women with breast cancer: Stoic resignation versus
24
25 spiritually-focused acceptance. Psychooncology 9 (5):S90-S90
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1 Summary of psychometric results on the dimensionality of the Mini-MAC

Study author	Year	<i>N</i>	Analysis type	Extraction	Rotation	No. of factors
Watson (UK) [1]	1994	573	Exploratory	PCA+ML	Oblique	5
Ho (Hong Kong) [2]	2003	115	Exploratory	PCA	Varimax	3
Grassi (Italy) [3]	2005	430	Exploratory	PCA	Varimax	5
Anagnostopoulos (Greece) [4]	2006	198	Confirmatory	ML	/	5
Kang (Korea) [5]	2008	201	Exploratory	PCA	Varimax	4
Bredal (Norway) [6]	2010	400	Exploratory	PCA	Varimax	4
Hulbert-Williams (UK) [7]	2012	153	Exploratory	PAF	Oblique	4

PCA, principal component analysis; ML, maximum likelihood; PAF, principal axis factoring. In the confirmatory factor analysis study by Anagnostopoulos [4], six cross-loadings and three residual covariances were additionally estimated.

Table 2 Goodness-of-fit indices for various EFA models of Mini-MAC ($n = 364$)

Model	χ^2	df	CFI	TLI	RMSEA	SRMR	$\Delta\chi^2(\Delta df)$
2-factor	1304.9*	349	.899	.883	.087	.068	
3-factor	983.3*	322	.930	.912	.075	.057	281.7(27)*
4-factor	741.1*	296	.953	.936	.064	.045	218.1(26)*
5-factor	556.3*	271	.970	.955	.054	.035	166.2(25)*
6-factor	490.0*	247	.974	.958	.052	.032	81.5(24)*

df , degree of freedom; CFI, comparative fit index; TLI, Tucker-Lewis index; RMSEA, root mean square error of approximation; SRMR, standardized root mean square residual; $\Delta\chi^2$, change in chi-square compared to previous model. * $p < .01$.

Table 3 Factor loading matrix of the five-factor solution using robust weighted least square estimator with target rotation ($n = 364$)

Item	Factor				
	HH	AP	FS	FA	CA
4	.820*				
6	.676*				
12	.766*				
14	.660*				
15	.783*				
16	.658*				
20	.680*				
21	.720*				
7	.541*				
5		.409*			
9		.688*			
13		.543*			
22		.743*			
25		.423*			
28		.692*			
29		.792*			
2			.474*		
10			.504*		
18			.600*		
23			.781*		
1				.780*	
3				.401*	
8				.512*	
19				.792*	
24				.458*	
11					.417*
17					.745*
26			.541*		.484*
27					.757*

Item-factor loadings with magnitude less than .40 are not shown.

HH, Helplessness/Hopeless; AP, Anxious Preoccupation; FS, Fighting Spirit; FA, Fatalism; CA, Cognitive Avoidance. * $p < .01$.

Table 4 Cronbach's alpha, correlations, and test-retest reliability among the Mini-MAC factors ($n = 364$)

Factor	α	HH	AP	FS	FA	CA
HH (9 items)	.876	<i>.533*</i>				
AP (7 items)	.871	<i>.677*</i>	<i>.640*</i>			
FS (5 items)	.692	<i>-.410*</i>	<i>-.231*</i>	<i>.454*</i>		
FA (5 items)	.674	<i>-.221*</i>	<i>.010</i>	<i>.548*</i>	<i>.449*</i>	
CA (4 items)	.705	<i>.437*</i>	<i>.449*</i>	<i>.024</i>	<i>.188*</i>	<i>.523*</i>

Test-retest reliability coefficients are shown in italics on the diagonal ($n = 291$).

α , Cronbach's alpha; HH, Helplessness/Hopeless; AP, Anxious Preoccupation; FS, Fighting Spirit; FA, Fatalism; CA, Cognitive Avoidance. * $p < .01$.

Table 5 Correlations between the Mini-MAC factors and psychosocial outcomes ($n = 359-364$)

Outcome	Factor score				
	HH	AP	FS	FA	CA
General health	<i>.441*</i>	<i>.443*</i>	<i>-.240*</i>	<i>-.199*</i>	<i>.194*</i>
Perceived stress	<i>.488*</i>	<i>.512*</i>	<i>-.298*</i>	<i>-.175*</i>	<i>.218*</i>
Anxiety	<i>.563*</i>	<i>.601*</i>	<i>-.328*</i>	<i>-.201*</i>	<i>.277*</i>
Depression	<i>.558*</i>	<i>.431*</i>	<i>-.308*</i>	<i>-.242*</i>	<i>.265*</i>

HH, Helplessness/Hopeless; AP, Anxious Preoccupation; FS, Fighting Spirit; FA, Fatalism; CA, Cognitive Avoidance. * $p < .01$.