

## O-5

**Heterozygous Overexpression of Preproendothelin-1 in Endothelial Cells Enhances Thromboxane-Prostanoid Receptor-Induced Contractions in the Renal Artery of Obese Mice**Oliver Baretella<sup>1</sup>, Sookja K. Chung<sup>2,4</sup>, Aimin Xu<sup>1,3,4</sup>, Paul M. Vanhoutte<sup>1,4</sup><sup>1</sup>Department of Pharmacology & Pharmacy, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong, China, <sup>2</sup>Department of Anatomy, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong, China, <sup>3</sup>Department of Medicine, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong, China, <sup>4</sup>Research Centre of Heart, Brain, Hormone & Healthy Aging, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong, China

Circulating levels of the endothelium-derived peptide endothelin-1 (ET-1) are elevated in human obesity, and ET-1 mediated vascular tone is increased. The renal artery is important in controlling intrarenal blood flow and is highly sensitive to ET-1. Whether or not ET-1 affects renal artery tone in obesity is unknown. To investigate the role of endogenous ET-1, a mouse model with *tie-1* promoter-driven endothelium-restricted heterozygous overexpression of preproendothelin-1 was used (TET+/-). Obesity was induced in TET+/- and WT littermates by feeding a high fat diet for seven months; lean controls were kept on standard chow. The main renal arteries were studied in wire myographs testing contractions (in the presence of L-NAME) to ET-1, serotonin (5-HT), and U46619, targeting ET<sub>A</sub>, 5-HT<sub>2</sub>, and TP receptors, respectively. Contractions to ET-1 were comparable between groups ( $PD_2$  8.29±0.05,  $n=6-8$ ); 5-HT-induced responses were facilitated at lower concentrations in obese mice leading to a shift in  $PD_2$  (lean 7.08±0.02 vs. obese 7.23±0.07,  $n=5-8$ ,  $P<0.01$ ). Responses to U46619 were significantly shifted to the left in renal arteries of obese animals ( $PD_2$  8.57±0.06 vs. lean 8.21±0.05,  $n=5-8$ ,  $P<0.001$ ), and the area under the curve was significantly different between lean and obese TET+/- mice (AUC 418±23 vs. lean 319±25,  $n=5$ ,  $P<0.05$ ). Thus, TET+/- had no effect on responses in lean animals. By contrast, in obesity heterozygous overexpression of ppET-1 enhanced TXA<sub>2</sub>-mediated, but not 5-HT or ET-1 induced contractions of the renal artery.

## O-6

**The Effect of Proteinuria-Mediated Endothelin-1 Downregulation of PKC $\alpha$  Signalling in Proximal Tubular Cells and Its Successful Treatment is Measurable Using microRNA15a as Biomarker in Vitro and in Vivo**Heike Loeser<sup>1</sup>, Melanie von Brandenstein<sup>1</sup>, Maike Wittersheim<sup>1</sup>, Volker Burst<sup>2</sup>, Claudia Richter<sup>1</sup>, Bernd Hoppe<sup>3</sup>, Jochen W.U. Fries<sup>1</sup><sup>1</sup>Institute of Pathology, University Hospital Cologne, Cologne, Germany, <sup>2</sup>Department of Internal Medicine II, Division of Nephrology, University Hospital Cologne, Cologne, Germany, <sup>3</sup>Institute of Pediatrics, Division of Nephrology, University Hospital Cologne, Cologne, Germany

In proteinuric diseases, stimulation of proximal tubule cells (RPTECs) by protein and endothelin-1 result in the activation of different signal pathways, ultimately causing renal insufficiency. Therapeutic interventions are hampered by the lack of specific and easily detectable markers. We described a regulatory pathway in which nuclear migration of protein kinase C  $\alpha$  controls the release of pri-miRNA15a. After endothelin-1 stimulation the migration of PKC $\alpha$  is inhibited, and mature miRNA15a is made. Using qRT-PCR we detect miRNA15a in the urine of adult and pediatric patients with membranous or minimal change nephropathy. By laser-microdissection this miRNA is predominantly located in the proximal tubules. In cell culture, human RPTECs produce the highest miRNA15a levels after ET-1 stimulation. In rats after 5/6 nephrectomy, miRNA15a is increased in the urine. By graded sieving and qRT-PCR, the highest amount of miRNA15a is found in the tubular fraction. Selegiline treatment upregulates PKC $\alpha$  in vitro and in the murine adriamycin model, significantly downregulating ET-1 induced miRNA15a production. Thus measuring urinary miRNA15a levels: i) indicates the regulation of a signal pathway in RPTECs in vivo in proteinuric conditions; ii) allows for the first time to control the effectiveness of a therapy aiming to protect proximal tubules.