

RESEARCH ARTICLE

Open Access



Towards microbiome transplant as a therapy for periodontitis: an exploratory study of periodontitis microbial signature contrasted by oral health, caries and edentulism

Alex E. Pozhitkov^{1*}, Brian G. Leroux¹, Timothy W. Randolph³, Thomas Beikler⁴, Thomas F. Flemmig⁵ and Peter A. Noble^{1,2}

Abstract

Background: Conventional periodontal therapy aims at controlling supra- and subgingival biofilms. Although periodontal therapy was shown to improve periodontal health, it does not completely arrest the disease. Almost all subjects compliant with periodontal maintenance continue to experience progressive clinical attachment loss and a fraction of them loses teeth. An oral *microbial transplant* may be a new alternative for treating periodontitis (inspired by fecal transplant). First, it must be established that microbiomes of oral health and periodontitis are distinct. In that case, the health-associated microbiome could be introduced into the oral cavity of periodontitis patients. This relates to the goals of our study: (i) to assess if microbial communities of the entire oral cavity of subjects with periodontitis were different from or oral health contrasted by microbiotas of caries and edentulism patients; (ii) to test *in vitro* if safe concentration of sodium hypochlorite could be used for initial eradication of the original oral microbiota followed by a safe neutralization of the hypochlorite prior transplantation.

Methods: Sixteen systemically healthy white adults with clinical signs of one of the following oral conditions were enrolled: periodontitis, established caries, edentulism, and oral health. Oral biofilm samples were collected from sub- and supra-gingival sites, and oral mucosae. DNA was extracted and 16S rRNA genes were amplified. Amplicons from the same patient were pooled, sequenced and quantified. Volunteer's oral plaque was treated with saline, 16 mM NaOCl and NaOCl neutralized by ascorbate buffer followed by plating on blood agar.

Results: Ordination plots of rRNA gene abundances revealed distinct groupings for the oral microbiomes of subjects with periodontitis, edentulism, or oral health. The oral microbiome in subjects with periodontitis showed the greatest diversity harboring 29 bacterial species at significantly higher abundance compared to subjects with the other assessed conditions. Healthy subjects had significantly higher abundance in 10 microbial species compared to the other conditions. NaOCl showed strong antimicrobial properties; nontoxic ascorbate was capable of neutralizing the hypochlorite.

Conclusions: Distinct oral microbial signatures were found in subjects with periodontitis, edentulism, or oral health. This finding opens up a potential for a new therapy, whereby a health-related entire oral microbial community would be transplanted to the diseased patient.

Keywords: Bacteriotherapy, Microbial transplant, Caries, Edentulism, Periodontitis, Red complex

* Correspondence: alexander.pozhitkov@evolbio.mpg.de

¹Department of Oral Health Sciences, University of Washington, Box 3574444, Seattle, WA 98195-7444, USA

Full list of author information is available at the end of the article

Background

The human oral microbiome is composed of a wide variety of microorganisms that play important roles in health and disease. Bacteria, for example, maintain oral and systemic health [1], but can also cause disease. Bacteriophages shape microbial diversity [2]. Protozoans and fungi consume food debris and other microbes [3, 4] and archaea utilize fermentation byproducts to produce methane [5]. Yet, in terms of microbial species, our current understanding of the human oral microbiome is limited to a few well-studied microorganisms. Based on what is known about these microbes, researchers have developed conceptual models with the explicit purpose of determining the etiology of oral diseases (e.g., [6, 7]). These models suggest putative interactions among the microorganisms as well as between microorganisms and the host. Although these studies have significantly advanced the field, recent studies show that the oral cavity is a complex and dynamic habitat consisting of hundreds of different interacting species [8, 9].

The conceptual model for periodontitis is that *Porphyromonas gingivalis*, *Tannerella forsythia*, *Treponema denticola* and *Aggregatibacter actinomycetemcomitans* are responsible for the disease [6, 10, 11]. Evidence from animal models indicated that these bacteria may disrupt tissue homeostasis by manipulating signaling pathways of the host and once the innate immunity is compromised, cause a shift in the relative abundance of microbes, resulting in inflammation and bone loss [12, 13]. Yet, it is well established that these bacteria i.e., *P. gingivalis*, *T. forsythia*, *T. denticola*, *A. actinomycetemcomitans*, are also found in healthy oral cavities (e.g., [14–19]). Moreover, *P. gingivalis* is not found in up to half of the patients with chronic or aggressive periodontitis [20]. The paradox of these microbes being present in both health and disease questions the validity of the conceptual underpinnings of the etiology of periodontitis, which invites for alternative ways of thinking about periodontitis as well as other oral diseases.

Conventional periodontal therapies aim at controlling supra- and subgingival biofilms and managing prognostic factors such as poor glycemic control in subjects with diabetes mellitus and active smoking [21, 22]. Although periodontal therapy has been shown to improve periodontal health and reduce the rate of further clinical attachment loss [23, 24] and tooth loss due to periodontitis [25–27], it falls short of completely arresting the disease. Almost all subjects compliant with periodontal maintenance care continue to show signs of progressive clinical attachment loss, and one- to two- thirds of them, lose one or more teeth during an extended period of periodontal maintenance care [28, 29]. We believe new thinking is necessary for finding successful ways of treating periodontitis.

The knowledge from a different field indicates that an appropriate microbial community is a key for maintaining

resistance against infection. A classic example is the gut microbiome, which protects the host from *Clostridium difficile* (CD) infection. Specifically, some patients develop CD colonization of their gut upon antibiotic treatment, which eradicates out their innate microbiome. A successful strategy for combating CD by transplanting gut microbiota from a healthy donor was first medically recorded more than 50 years ago [30]. (The origins of the procedure date back to the 4th century in China [31]). Specifically, a suspension of stool taken from a systemically healthy intimate partner, relative or friend is introduced into the gastrointestinal tract of the recipient via colonoscopy, enema or a nasogastric tube. Recent systematic review reports approx 90 % efficiency of eradicating the CD infection by a fecal transplant [32]. Besides the CD infection, other potentially dysbiotic diseases were found to be responsive to the microbial transplantation, such as inflammatory bowel disease, obesity. For a review on current developments in the gut microbial transplant see Kelly et al. [33].

Inspired by the success of the fecal transplant, here we put forward a concept of a microbial transplant as a potential therapy for periodontitis. We envision the therapy of consisting of three steps: (i) harvesting sub- and supra-gingival microbiota from a healthy donor, e.g., spouse or a partner; (ii), performing deep cleaning, root planning and applying a broad-spectrum antimicrobial agent to the periodontitis patient; and (iii) neutralizing the antimicrobial agent immediately following by a rinsing with a microbial suspension harvested from the healthy donor in the periodontitis patient.

The objectives of the present study were two-fold. The first objective was to assess if the entire oral microbial community of periodontitis was different from established caries, edentulism, and oral health. The second objective was to test an approach for applying a broad-spectrum safe antimicrobial agent (NaOCl) to significantly reduce the load of existing microbiome followed by neutralization of this agent for subsequent microbial transplantation.

Methods

Study subjects

Adult subjects were recruited at the Department of Periodontics, University of Washington, Seattle, WA, U.S.A. and Section of Periodontics, University of Düsseldorf, Germany. A written informed consent for participation in the study was obtained from the subjects. The study was approved by the University of Washington Institutional Review Board, ref. number 3570. Subjects were enrolled if they had one of the following clinical conditions: severe periodontitis, caries, edentulism, or oral health. A periodontitis case was defined as having at least 2 interproximal sites at different teeth with clinical

attachment loss (CAL) of 6 mm or greater and at least 1 interproximal site with probing depth (PD) of 5 mm or greater [34] and a minimum of 20 permanent teeth, not including 3rd molars. Subjects were excluded from the periodontitis group if they had multiple established caries lesion or wore a removable partial denture. A caries case was defined as having the following number of teeth with established caries lesions: 6 or more teeth in subjects 20 to 34 years of age; 4 or more teeth in subjects 35 to 49 years of age; and 3 or more teeth in subjects 50 years of age and older. Established caries was defined as a class 4 lesion according to the International Caries Detection and Assessment System. The number of teeth with caries lesion in caries cases was greater than one standard deviation above the mean of caries extent in respective age group the U.S.A. [35]. Exclusion criteria for a caries case were interproximal sites with CAL of 4 mm or greater or PD of 5 mm or greater [34]. An edentulous case had to be completely edentulous in both jaws and their teeth had to be extracted more than one year before the enrollment in the study. A healthy case was defined as having 28 teeth, not counting 3rd molars, or 24 or more teeth, not counting 3rd molars if premolars had been extracted for orthodontic reasons or were congenitally missing with no signs of oral disease. Exclusion criteria for a healthy case included: smoking, loss of permanent teeth due to caries or periodontitis, any interproximal sites with CAL of 4 or greater or PD of 5 mm or greater, or any established caries lesions. Exclusion criteria for all groups included: oral mucosal lesions, systemic diseases, and use of antibiotics or local antiseptics within 3 months prior to the study.

Sample collection

For all but the edentulous patients, supra- and subgingival plaque was collected from six sites with the deepest probing depth in each sextant. One sterile paper point per site was inserted into the deepest aspect of the periodontal pocket or gingival sulcus. Biofilm from oral mucosae was collected by swiping a sterile cotton swab over the epithelial surfaces of the lip, left and right buccal mucosae, palate, and dorsum of the tongue [36]. Samples were stored at -80°C .

Molecular methods

Microbial DNA was isolated from cells by physical and chemical disruption using zirconia/silica beads and phenol-chloroform extraction in a FastPrep-24 bead beater [37]. Prokaryotic 16S rRNA genes were amplified using universal primers (27 F and 1392R) using the GemTaq kit from MGQuest (Cat# EP012). The PCR program involved a pre-amplification step of 10 cycles with annealing temperature of 56°C followed by 20 amplification cycles with annealing temperature 58°C .

In each cycle, elongation time was 1 min 10s, at 72°C . PCR was finalized by extended elongation for 5 min. PCR products were purified with DNA Clean & Concentrator columns (Zymo Research, USA) and quantified using the NanoDrop (Agilent, USA).

Due to technical reasons, for some subjects, subgingival and mucosal microbiotas were pooled together into one vial, while for other subjects the subgingival and mucosal microbiotas were stored separately. Since our goal was to investigate the entire microbial community, for the separately stored supra and subgingival microbiota samples, equal quantities of PCR product derived from swab and paper point samples were pooled together for each patient. For edentulous patients, there were no paper point samples. Each purified PCR product, 500 ng, was labeled with a Multiplex Identifier (MID) during the Roche Rapid Library preparation step. Four MID-tagged sequences, representing each of the conditions, were combined in equimolar concentrations and subjected to emPCR and DNA sequencing protocols as specified by the manufacturer's recommendations for the Roche 454 Jr. instrument.

Data analysis

The obtained sequences were separated out by their respective Multiplex Identifier (MID) and uploaded to the MG-RAST web server [38]. The MG-RAST pipeline assessed the quality of sequences, removed short sequences (multiplication of standard deviation of length cutoff of 2.0) and removed sequences with ambiguous bp (non-ACGT; maximum allowed number of ambiguous base pair was set to 5). The pipeline annotated the sequences and allowed the integration of the data with previous metagenomic and genomic samples. The RDP database was used as annotation source, with minimum sequence identity of 97 %, maximum e-value cutoff at 10^{-5} , and minimum sequence length of 100 bases. Alpha diversity analysis was conducted in MG-RAST.

Orthogonal transformation of the annotated rRNA genes to their principal components (PC) was conducted using normalized abundances [39, 40]. Normalization of the abundance was performed identically to the procedure used by MG-RAST. Specifically, abundances were increased by one, log₂ transformed, and centered to produce relative values. Relative values were standardized by dividing them by the standard deviation of the log₂ values [38]. The data were graphed on a 2 dimensional ordination plot. To determine the relative contribution of the microbial species to the plot, we let X denote the 16 by 578 (patients by species) matrix of the normalized abundance values. The matrix X was used to produce a 16×16 matrix D of distances between all pairs of subjects. Principal coordinate analysis (PCoA, i.e., multidimensional scaling) was achieved by performing principal

component analysis (PCA) on the matrix of distances, D. The Euclidean distance metric was used to create this matrix, but this metric produced results similar to those when the alternative Bray-Curtis distance was used. To investigate and visualize differences between the four patient groups, the first two principal components, PC1 and PC2, of the distance matrix D were retained. To establish which species were most prominently responsible for the groupings, the projection of each species onto the (PC1, PC2) plane was calculated; those species with the largest projections are displayed in the right panel of Fig. 3. That is, this figure shows a biplot of the species most highly correlated PC1 and PC2.

Mann-Whitney test ($\alpha = 0.05$) was used to investigate significant difference in normalized relative abundances. The abundance data were normalized in two different ways: MG-RAST normalization as described above and raw abundances normalized to the total number of reads in a sample. Differences in alpha diversities by condition were determined using ANOVA. Two-tailed T-tests, assuming unequal variance, were used to investigate if there was a significant difference in the means ($\alpha = 0.05$). These analyses were performed using SAS JMP.

Sodium hypochlorite experiments

According to previously suggested concentration [41], sodium hypochlorite experiments were conducted with a 1:50 dilution of the 6 % household bleach (Clorox, The Clorox Company, USA), the “hypochlorite working solution”. The dilution corresponds to 16 mM NaOCl. The neutralizing agent was a sodium ascorbate buffer produced by adjusting pH of a 23 mM ascorbic acid solution with concentrated NaOH until pH 5.3.

Three samples of dental plaque from a volunteer were subjected to three challenges: (i) resuspension in saline, (ii) resuspension in the hypochlorite working solution and (iii) resuspension in the hypochlorite working solution that was previously neutralized with equal volume of the ascorbate buffer. Resuspended plaque samples were plated on blood agar plates.

Active chlorine concentration was crudely assessed by iodine colorimetry. A colorimetric solution contained 1 volume of 0.1 % stabilized starch solution (Fisher Scientific) mixed with 0.1 volume of 1 M potassium iodide. Two volumes of the colorimetric solution were mixed with 1 volume of the solution containing NaOCl. Intense blue color indicated detectable active chlorine. Iodine colorimetry was calibrated with a dilution series of the hypochlorite working solution with the following dilutions 1:200, 1:400, 1:800, 1:1600, 0. The dilution 1:800 still showed a hint of blue, while 1:1600 was completely colorless.

Results

Demographics

A total of 16 subjects, 4 subjects with each of the assessed oral conditions, were enrolled into the study. Ten subjects were from Germany and 6 from the U.S.A.; 9 were females and 7 males. The age of the enrolled subjects ranged from 28 to 92 years. Subjects with periodontitis had a median of 34 sites (range 15 to 48) with PD of 4 to 6 mm and 5 sites (range 4 to 26) with PD of 7 mm or greater while none of the subjects with caries and none of the healthy subjects had any sites with PD of 4 mm or greater. Subjects with caries had a median of 14 teeth (range 9 to 17) with established caries while periodontitis and healthy subjects were mostly caries free (Table 1).

Microbial signatures

The average (\pm std) number of 16S rRNA amplicon sequences per individual oral microbiota was $13,104 \pm 5,533$ (Additional file 1: Table S1). The length and GC content of the sequences was similar for all individuals: 514 ± 10 bp and 53 ± 1 bp, respectively. Comparison of the number of sequences, sequence length, or GC content revealed no significant differences by oral condition or gender, indicating a balanced data set. The rarefaction curve of most samples approached saturation, indicating sufficient reads for comparisons by condition and gender (Fig. 1).


Alpha diversity of the oral microbiome in periodontitis subjects was significantly ($p < 0.05$) greater compared to that found in subjects with caries and edentulous subjects. Although the oral microbiome was more diverse in periodontitis subjects compared to healthy subjects, the difference missed statistical significance ($p = 0.06$) (Fig. 2). The composition of the oral microbiome in subjects with periodontitis was distinctly different from that of all other oral conditions. Specifically, the periodontitis condition had greater abundances of Bacteroidetes (32 %),

Table 1 Demographics of the patients population

Parameter	Condition			
	Healthy	Periodontitis	Caries	Edentulous
America/Europe	0/4	2/2	2/2	2/2
Male/Female	2/2	2/2	2/2	1/3
Ages	41 (31–52) ^a	47 (28–58)	39 (29–49)	77 (58–92)
Number of teeth	27 (25–28)	28 (22–30)	28 (22–31)	0
ICDAS ≥ 4	0	0 (0–1)	14 (9–17)	–
PD ≤ 3 mm	100	62 (26–79)	100	–
PD 4–6 mm	0	34 (15–48)	0	–
PD ≥ 7 mm	0	5 (4–26)	0	–

ICDAS International Caries Detection and Assessment System, 4 denotes established decay (dentine shadow), PD pocket depth


^aMedian (min-max)



Fusobacteria (7 %), Spirochaetes (5 %) and Synergisetes (1 %) and fewer Actinobacteria (14 %) (Fig. 3). Actinobacteria were most abundant in the oral microbiome of healthy (28 %) and edentulous subjects (34 %), followed by Firmicutes, which occurred in abundances of 22 % and 30 %, respectively (Fig. 3). Subjects with caries showed somewhat lower abundances of Actinobacteria (19 % vs. 29 %) and slightly greater abundances of Fusobacteria (5 % vs. 3 %) in the oral microbial communities compared to healthy subjects (Fig. 3).

“The red complex”

An ordination plot based on the 16S rRNA genes revealed that the oral microbiomes in subjects with periodontitis, edentulism, or oral health formed distinct groups (Fig. 4, left panel). The ordination plot explained 78 % of the observed variability. These results, along with the high diversity values, suggest that oral microbiome in subjects with periodontitis, edentulous subjects,



and healthy subjects were very different from each other. Permutation MANOVA (pseudo F test, refs. [42, 43]) showed significant differences in microbial species abundances between the four oral conditions (Bonferroni-adjusted p-value of 0.0083). The Bonferroni adjustment refers to the 6 comparisons, i.e., all pairs of the 4 groups. In order to determine if the “red complex” bacteria (*P. gingivalis*, *T. denticola* and *T. forsythia*) were absolute determinants for the microbial signature of periodontitis, the abundances of these organisms were removed from the original data set. Non-parametric MANOVA and ordination plot analyses of the data set without “red complex” bacteria did not alter the findings shown in Fig. 4a or the p-value.

Projecting the top 23 contributing bacteria on the PCoA plot (i.e., biplot) shows the relative contribution of bacteria to the ordination (Fig. 4, right). Since the purpose of the biplot was predominantly illustrative, the number of displayed species, 23, was chosen arbitrary based on the upper tail of the histogram of projection sizes (Additional file 1: Figure S1). *P. endodontalis*, *Prevotella intermedia*, *T. vincentii*, and an uncultured *Porphyromonas* sp. were found to contribute to the microbial composition in periodontitis while a complex of *P. melaninogenica*, *Fusobacterium nucleatum*, *P. catoniae*, *Capnocytophaga ochracea* and *C. sputigena*, *C. gingivalis*, *Haemophilus parainfluenzae* and *Neisseria elongata* contributed to the microbial composition of orally healthy subjects. Members of the *Streptococcus* and *Lactobacillus* were associated with edentulism.

The species contributing to the differences in diversity and the groupings in the ordination plot are shown in Table 2. Twenty-nine out of the 587 microbial species had higher abundances in patients with periodontitis than those in the other assessed conditions. In the healthy subjects, 10 out of the 587 microbial species were found in higher abundances compared to the non-healthy subjects (Table 3).

Antimicrobial agent and its neutralization

Antimicrobial activity of sodium hypochlorite was tested with a volunteer’s dental plaque. Resuspension of the plaque in 1:50 dilution of the household bleach, i.e., “the hypochlorite working solution” resulted in a complete inhibition of bacterial growth on a blood agar. In contrast, the viability of whole dental plaque bacteria was maintained following exposure to the hypochlorite working solution inactivated for 7 min with 23 mM sodium ascorbate buffer (Fig. 5). Starch-iodine colorimetry assay revealed that sodium ascorbate - ascorbic acid buffer reduced the concentration of active chlorine by a factor of at least 800–1600, i.e., below 10–20 μM , after 30 s of inactivation.




Fig. 3 Percent abundance of microbes by phylum and condition

Discussion

Potential for the periodontitis bacteriotherapy

The first step towards developing a bacteriotherapy for periodontitis is to establish the existence of a microbial community characteristic of periodontitis as well as the community characteristic of oral health. Two additional communities from edentulous and caries patients, whose microbial communities are expected to be distinct, were added for contrasting purposes.

Several studies have compared microbial communities isolated from different sites in the oral cavity and from patients with different clinical conditions (e.g., [14, 15, 17, 19, 44–47]). One of these studies revealed distinct partitioning of bacterial communities in subgingival biofilms from healthy and diseased periodontal sites [45]. Another study indicated that not only the oral

microbial species were drastically different between periodontal health and disease but also their gene expression [48]. Moreover, in subjects with periodontitis, disease-associated bacteria were found on the buccal mucosae, tongue, and saliva indicating that the entire oral microbiome might be altered [49–53].

Since the bacteriotherapy would likely be performed by attempting to transplant both sub- and supra-gingival microbes, the present study was set to test the hypothesis that overall compositions of the oral microbial communities were distinct between the conditions of periodontitis, edentulism, caries, and oral health. In fact, the entire oral microbiome of each subject was obtained by pooling together several oral cavity sub-microbiomes such as subgingival, supragingival, and mucosal membranes.




Fig. 4 Left panel: PCoA ordination plot of the 16 human oral samples by condition. Condition: red, periodontitis; blue, caries; black, edentulous; green, healthy. Right panel: Projection of major microbial species contributing to the groups

Table 2 Bacterial species that had significantly different abundances (%) between individuals with the non-periodontitis (NP; e.g., healthy, caries, edentulous) and periodontitis (P) abundances based on Mann–Whitney test (alpha = 0.05)

Phylum/class	Genus/species	NP	P
Actinobacteria	<i>Atopobium vaginae</i>	0.01	0.35*
	<i>Actinomyces georgiae</i>	0.05	0.35**
	<i>Actinomyces meyeri</i>	0.10	0.49**
Bacteroidetes	<i>Porphyromonas endodontalis</i>	0.12	2.91*
	<i>uncultured Porphyromonas sp.</i>	0.04	1.26*
	<i>Porphyromonas gingivicanis</i>	0.00	0.04**
	<i>Tannerella forsythia</i>	0.20	0.83*
	<i>Prevotella intermedia</i>	0.12	3.23*
	<i>Prevotella pallens</i>	0.14	0.85*
	<i>Prevotella pleuritidis</i>	0.00	0.01**
	<i>Prevotella salivae</i>	0.20	0.74**
Firmicutes/Clostridia	<i>Parvimonas micra</i>	0.12	1.19*
	<i>Eubacterium saburreum</i>	0.07	0.28*
	<i>Pseudobutyrvibrio xylanivorans</i>	0.00	0.06*
	<i>Peptostreptococcus anaerobius</i>	0.07	0.50*
Firmicutes/Negativicutes	<i>Megasphaera micronuciformis</i>	0.12	0.92*
Fusobacteria	<i>Leptotrichia wadei</i>	0.10	0.32**
Proteobacteria/Deltaproteobacteria	<i>uncultured Desulfobulbus sp.</i>	0.00	0.02*
Proteobacteria/Epsilonproteobacteria	<i>Campylobacter rectus</i>	0.03	0.14*
	<i>Campylobacter concisus</i>	0.03	0.07**
Spirochaetes/Spirochaetales	<i>Treponema denticola</i>	0.07	1.60*
	<i>Treponema maltophilum</i>	0.01	0.25*
	<i>Treponema medium</i>	0.03	0.29*
	<i>Treponema socranskii</i>	0.05	0.21*
	<i>Treponema sp.</i>	0.03	0.10*
	<i>Treponema vincentii</i>	0.07	0.71*
	<i>Treponema pectinovorum</i>	0.00	0.04**
Synergistetes	<i>Synergistetes bacterium SGP1</i>	0.05	0.23*
	<i>Aminobacterium colombiense</i>	0.01	0.05*

*Significant difference using both normalization methods (i.e., raw species abundances were log₂ transformed, normalized to produce relative values ((raw values-average)/std) versus abundances normalized to the total number of reads in a sample)

**Significant difference for abundances using data normalized only to the total number of reads in a sample

The key finding of this study, critically important for the potential bacteriotherapy, is that microbial compositions (i.e., signatures) of the entire (or substantial subset of) oral microbiomes of subjects with periodontitis contrasted by edentulism and oral health are distinct. In addition, the microorganisms which are usually considered as pathogens of periodontitis, *P. gingivalis*, *T. denticola* and *T. forsythia*, are not the sole representatives of the periodontitis microbial signature. In particular, this study provides evidence that a broad microbial-community model of oral health may offer a valuable perspective when considered in conjunction with (or in contrast to) more common single-pathogen models ([7] and refs. therein). With this perspective, it is reasonable to

expect that microbial community composition is a function of interspecies competition and cooperation, which provides optimism for further developing the bacteriotherapy. Below, we expand on the motivation for choosing molecular methods and sampling procedures.

Methodological validity

Historically, subgingival oral microbiome was predominantly assessed by a checkerboard hybridization technique revealing signals from approx. 40 species (for review, see [54]). Pathogenicity of a few species, i.e., “colored complexes”, was presumed to account for oral conditions, such as periodontitis. With the advent of next generation sequencing, extensive cataloging of the subgingival

Table 3 Microbial species that had significantly different abundances (%) between individuals with the non-healthy (NH; periodontitis, caries, edentulous) and healthy (H) abundances based on Mann–Whitney test (alpha = 0.05)

Phylum/class	Genus/species	NH	H
Actinobacteria	<i>Micrococcus lylae</i>	0.00	0.03**
Bacteroidetes	<i>Prevotella marshii</i>	0.01	0.09**
Firmicutes/Bacilli	<i>Abiotrophia para-adiacens</i>	0.22	0.60**
	<i>Granulicatella elegans</i>	0.20	0.66*
	<i>Granulicatella adiacens</i>	0.60	1.27**
	<i>Streptococcus iniae</i>	0.04	0.15*
Firmicutes/Negativicutes	<i>Selenomonas ruminantium</i>	0.00	0.10*
Proteobacteria/Betaproteobacteria	<i>Neisseria polysaccharea</i>	0.00	0.02*
Proteobacteria/Gammaproteobacteria	<i>Aggregatibacter segnis</i>	0.01	0.21**
	<i>Haemophilus parainfluenzae</i>	0.17	0.46*

*Significant difference determined using both normalization methods (i.e., raw species abundances were log2 transformed, normalized to produce relative values ((raw values-average)/std) versus abundances normalized to the total number of reads in a sample)

**Significant difference for abundances using data that was normalized to only the total number of reads in a sample

microbiome has been conducted (e.g., [17, 45, 55]). In our study we also utilized high-throughput sequencing technology, because it provides an in-depth qualitative and quantitative characterization of the microbial communities, which is essential for discovery of microbial signatures.

The data for the analysis of microbial signatures were the relative abundances of microorganisms measured by the number of sequencing reads obtained from high-throughput sequencing of each subject’s oral microbiome. Contrary to what one might expect, it is important to note, that absolute abundances are not

attainable by the high throughput sequencing due to the biases in amplification and other physicochemical reasons [56, 57]. After sequencing reads are processed, each subject is represented by a p-dimensional vector (p = 597) in the multidimensional space, with each coordinate corresponding to the relative abundance of each microbial species. The ordination analysis [58] presented in the Results section is a standard dimension-reducing method, which reveals the structure of the 3 groups (Fig. 4). Interpretation of these groups is based on the subjects’ microbial abundance profiles that are most similar and hence grouped closer to each other. Although the sample size is




Fig. 5 Dental plaque treated with saline, the hypochlorite working solution or inactivated hypochlorite working solution

rather small ($n = 16$), the quality of the clusters and their statistical significance (based on permutation) suggests that the observed grouping may be reproducible in a larger, independent cohort. Other researchers, e.g., Kumar et al. [47], showed with a larger group that there is a clear separation between microbial communities for healthy and periodontitis and peri-implantitis patients. Another potential issue with generalization of our findings could be due to the fact that all health subjects were from Europe. Nevertheless, as was discovered by Nasidze et al. [59], the compositions of the oral microbial communities obtained from 12 worldwide locations of 10 individuals each were “larger among individuals from the same location than among individuals from different locations”. Hence the differences in the composition among individuals were not determined by geography.

Periodontitis pathogenesis: “red complex” or entire community shift?

This study revealed grouping of subjects based on the composition of their entire oral microbiomes, which is consistent with previously reported results based on the microbial composition of their subgingival plaque samples [47]. Specifically, it has been suggested that there exist distinct microbial signatures characteristic of periodontitis, peri-implantitis, and oral health. Our research expands on these findings by investigating the entire oral microbiome including combined together subgingival, supragingival, and mucosal sub-microbiomes. Building upon a recent report by Griffen et al. [45], which showed a high microbial diversity of sub-gingival biofilms in subjects with periodontitis compared to oral health, our study also revealed a similarly increased diversity in the entire oral microbiome of subjects with periodontitis compared to edentulism, oral health and caries. Among the four conditions, our study showed the lowest microbial diversity in oral microbiomes of subjects with edentulism. This is presumably due to the absence of teeth and subgingival pockets in which otherwise mature biofilms develop [45]. It is important to note that there are other studies focused on microbial signatures of periodontitis (e.g., [60]), however these studies did not attempt to contrast several conditions.

Our observations support the polymicrobial nature of periodontitis contrary to the “old-school” long-standing notion of specific pathogens believed to cause periodontitis. Indeed, the oral microbial signature in periodontitis subjects included *P. gingivalis*, *T. denticola* and *T. forsythia*, which are collectively referred to as the “red complex”. As our results showed, red complex bacteria were not the absolute determinants of the periodontitis microbial signature. The presence of the red complex bacteria in oral health and other conditions could be attributed to differences in: (i) the host immune response,

(ii) genes encoding virulence factors among different strains, and (iii) gene regulation [51]. The findings of distinct oral microbial signatures between subjects with periodontitis and oral health supported the notion that periodontitis is associated with a dysbiosis (i.e., microbial imbalance) of the entire oral microbiome. The oral dysbiosis may be due to the assembly of a synergistic microbial community that tilts the balance away from a health-associated microbial homeostasis [51]. It could be initiated by the acquisition of virulence factors by microbes and/or changes in the regulation/expression of genes within the community [17, 61]. Perhaps one or several microorganisms, e.g., *P. gingivalis*, are capable of initiating synergistic interactions that ultimately result in the emergence of disease-provoking microbial community [7], however a mechanistic proof is lacking.

The idea for the potential bacteriotherapy of periodontitis is inspired by the success of the fecal microbial transplant treatment of *Clostridium difficile* (CD) infection. It should be noted that while the diversity of bacteria associated with periodontal disease is very high [55, 62], the diversity of bacteria associated with CD infections is very low [63]. The reason to emphasize these differences is that the goal of bacteriotherapy for CD is to increase bacterial diversity whereas the goal of bacteriotherapy for periodontitis is not the same since the bacterial diversity is already high. Presumably in both diseases, replacement of healthy stable microbial communities is the answer to prevent future dysbiosis. However, without performing actual experiments, one cannot speculate on the success or failure of bacteriotherapy for periodontal disease.

Antimicrobial treatment before the transplant

It is reasonable to assume that for the bacteriotherapy to be successful, the recipient’s oral cavity has to have lowest possible load of microorganisms to allow the new microbiome to flourish without interference from the previous disease-associated microbiome. Administration of adjunctive systemic antibiotics [64, 65] or a full-mouth disinfection approach that reduces the bacterial reservoir in all habitats of the oral cavity will result in more pronounced shifts in the microbial composition and significantly reduces the incidence of clinical attachment loss compared to supra- and subgingival debridement alone [37, 66, 67]. Still, it is desirable to administer such an antimicrobial agent that is effective against a broad spectrum of microbes and is easy to neutralize with another nontoxic agent, because as most antiseptics remain active in the oral cavity for quite some time [68, 69] and therefore, may affect the viability of an oral microbial transplant. We have been able to demonstrate that sodium hypochlorite (NaOCl), which is bactericidal against a panel of oral biofilm microorganisms [68, 70, 71], can be inactivated by a nontoxic sodium

ascorbate – ascorbic acid buffer. Although some critics have said that NaOCl treatment is antiquated and dangerous, a recent study proved the opposite [72]. Moreover, the Galvan et al. study [72] showed safety and a significant reduction of plaque and bleeding on probing in the group that used NaOCl rinses compared to the control group.

Conclusions

The present exploratory study revealed statistically significant microbial signatures of oral health and periodontitis. This finding presents a path forward for a potential new therapy for periodontitis, which could be based on substituting periodontitis-associated microbiome with the health-associated one. At the beginning (in 1950s), the fecal transplant therapy was successfully used without a deep mechanistic understanding. Currently, insights into the gene expression and metabolic effects of the therapy have been made (e.g., [73]). If the proposed therapy is proven to be successful, deep mechanistic studies will be of great interest.

Additional file

Additional file 1: Table S1. Number of sequences obtained from each condition and sample. **Table S2.** Comparison of oral microbial community composition in different studies. **Figure S1.** Determination of an arbitrary cut-off for displaying species on a biplot (Fig. 4, right panel). (PDF 136 kb)

Abbreviations

PCA: Principal component analysis; PCR: Polymerase chain reaction; MID: Multiplex identifier; PCoA: Principal coordinate analysis; bp: Base pairs.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

AEP, TB, TFF, PAN designed the experiments; AP, PAN conducted the experiments; BGL, TWR conducted statistical analysis; AP, PAN wrote the paper. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by National Science Foundation (NSF) grant DBI 10388671 to PAN, intramural funding of the Medical School, Heinrich-Heine-University Dusseldorf to TB and NIH training grant 5 T90 D021984-03 to AEP.

Author details

¹Department of Oral Health Sciences, University of Washington, Box 3574444, Seattle, WA 98195-7444, USA. ²PhD Program in Microbiology, Alabama State University, Montgomery, AL 36101, USA. ³Fred Hutchinson Cancer Research Center, 1100 Fairview Ave. N., PO Box 19024, Seattle, WA 98109, USA. ⁴Section of Periodontics, School of Medicine, Heinrich-Heine-University, Moorenstrasse 5, 40225 Dusseldorf, Germany. ⁵Faculty of Dentistry, The University of Hong Kong, Prince Philip Dental Hospital, 34 Hospital Road, Sai Ying Pun, Hong Kong, SAR, Peoples' Republic of China.

Received: 4 June 2015 Accepted: 6 October 2015

Published online: 14 October 2015

References

1. Wade WG. The oral microbiome in health and disease. *Pharmacol Res.* 2013;69:137–43.

2. Pride DT, Salzman J, Haynes M, Rohwer F, Davis-Long C, White 3rd RA, et al. Evidence of a robust resident bacteriophage population revealed through analysis of the human salivary virome. *ISME J.* 2012;6:915–26.
3. Wantland WW, Wantland EM, Remo JW, Winquist DL. Studies on human mouth protozoa. *J Dent Res.* 1958;37:949–50.
4. Ghannoum MA, Jurevic RJ, Mukherjee PK, Cui F, Sikaroodi M, Naqvi A, et al. Characterization of the oral fungal microbiome (mycobiome) in healthy individuals. *PLoS Pathog.* 2010;6, e1000713.
5. Lepp PW, Brinig MM, Ouverney CC, Palm K, Armitage GC, Relman DA. Methanogenic Archaea and human periodontal disease. *Proc Natl Acad Sci.* 2004;101:6176–81.
6. Holt SC, Ebersole JL. *Porphyromonas gingivalis*, *Treponema denticola*, and *Tannerella forsythia*: the “red complex”, a prototype polybacterial pathogenic consortium in periodontitis. *Periodontol* 2000. 2005;38:72–122.
7. Darveau RP, Hajishengallis G, Curtis MA. *Porphyromonas gingivalis* as a potential community activist for disease. *J Dent Res.* 2012;91:816–20.
8. Faust K, Sathirapongsasuti JF, Izard J, Segata N, Gevers D, Raes J, et al. Microbial co-occurrence relationships in the human microbiome. *PLoS Comput Biol.* 2012;8, e1002606.
9. Diaz PI, Dupuy AK, Abusleme L, Reese B, Obergfell C, Choquette L, et al. Using high throughput sequencing to explore the biodiversity in oral bacterial communities. *Mol Oral Microbiol.* 2012;27:182–201.
10. Dewhirst FE, Chen T, Izard J, Paster BJ, Tanner AC, Yu WH, et al. The human oral microbiome. *J Bacteriol.* 2010;192:5002–17.
11. Kumar PS, Griffen AL, Barton JA, Paster BJ, Moeschberger ML, Leys EJ. New bacterial species associated with chronic periodontitis. *J Dent Res.* 2003;82:338–44.
12. Darveau RP. Periodontitis: a polymicrobial disruption of host homeostasis. *Nat Rev Microbiol.* 2010;8:481–90.
13. Hajishengallis G, Liang S, Payne MA, Hashim A, Jotwani R, Eskan MA, et al. Low-abundance biofilm species orchestrates inflammatory periodontal disease through the commensal microbiota and complement. *Cell Host Microbe.* 2011;10:497–506.
14. Bik EM, Long CD, Armitage GC, Loomer P, Emerson J, Mongodin EF, et al. Bacterial diversity in the oral cavity of 10 healthy individuals. *ISME J.* 2010;4:962–74.
15. Diaz PI, Chalmers NI, Rickard AH, Kong C, Milburn CL, Palmer Jr RJ, et al. Molecular characterization of subject-specific oral microflora during initial colonization of enamel. *Appl Environ Microbiol.* 2006;72:2837–48.
16. Eick S, Pietkiewicz M, Sculean A. Oral microbiota in Swiss adolescents. *Clin Oral Investig.* 2013;17:79–86.
17. Liu B, Faller LL, Klitgord N, Mazumdar V, Ghodsi M, Sommer DD, et al. Deep sequencing of the oral microbiome reveals signatures of periodontal disease. *PLoS One.* 2012;7, e37919.
18. Papaioannou W, Gizani S, Haffajee AD, Quirynen M, Mamai-Homata E, Papagiannoulis L. The microbiota on different oral surfaces in healthy children. *Oral Microbiol Immunol.* 2009;24:183–9.
19. Zaura E, Keijsers BJ, Huse SM, Crielaard W. Defining the healthy “core microbiome” of oral microbial communities. *BMC Microbiol.* 2009;15:259.
20. Beikler T, Prior K, Ehmke B, Flemmig TF. Specific antibiotics in the treatment of periodontitis – a proposed strategy. *J Periodontol.* 2004;75:169–75.
21. Kaldahl WB, Kalkwarf KL, Patil KD, Molvar MP, Dyer JK. Long-term evaluation of periodontal therapy, II Incidence of sites breaking down. *J Periodontol.* 1996;67:103–8.
22. Tervonen T, Karjalainen K. Periodontal disease related to diabetic status. A pilot study of the response to periodontal therapy in type 1 diabetes. *J Clin Periodontol.* 1997;24:505–10.
23. Axelsson P, Lindhe J. The significance of maintenance care in the treatment of periodontal disease. *J Clin Periodontol.* 1981;8:281–94.
24. Haffajee AD, Socransky SS. Frequency distributions of periodontal attachment loss. Clinical and microbiological features. *J Clin Periodontol.* 1986;13:625–37.
25. Axelsson P, Lindhe J, Nystrom B. On the prevention of caries and periodontal disease. Results of a 15-year longitudinal study in adults. *J Clin Periodontol.* 1991;18:182–9.
26. Axelsson P, Nystrom B, Lindhe J. The long-term effect of a plaque control program on tooth mortality, caries and periodontal disease in adults. Results after 30 years of maintenance. *J Clin Periodontol.* 2004;31:749–57.
27. Hujoel PP, Leroux BG, Selipsky H, White BA. Non-surgical periodontal therapy and tooth loss. A cohort study. *J Periodontol.* 2000;71:736–42.
28. Konig J, Plagmann HC, Ruhling A, Kocher T. Tooth loss and pocket probing depths in compliant periodontally treated patients, a retrospective analysis. *J Clin Periodontol.* 2002;29:1092–100.

29. Rosling B, Serino G, Hellstrom MK, Socransky SS, Lindhe J. Longitudinal periodontal tissue alterations during supportive therapy. Findings from subjects with normal and high susceptibility to periodontal disease. *J Clin Periodontol*. 2001;28:241–9.
30. Eiseman B, Silen W, Bascom GS, Kauvar AJ. Fecal enema as an adjunct in the treatment of pseudomembranous enterocolitis. *Surgery*. 1958;44:854–9.
31. Zhang F, Luo W, Shi Y, Fan Z, Ji G. Should we standardize the 1,700-year-old fecal microbiota transplantation? *Am J Gastroenterol*. 2012;107:1755.
32. Kassam Z, Lee CH, Yuan Y, Hunt RH. Fecal microbiota transplantation for *Clostridium difficile* infection: systematic review and meta-analysis. *Am J Gastroenterol*. 2013;108:500–8.
33. Kelly CR, Kahn S, Kashyap P, Laine L, Rubin D, Atreja A, et al. Update on fecal microbiota transplantation 2015: indications, methodologies, mechanisms, and outlook. *Gastroenterology*. 2015;149:223–37.
34. Page RC, Eke PI. Case definitions for use in population-based surveillance of periodontitis. *J Periodontol*. 2007;78:1387–99.
35. Dye BA, Tan S, Smith V, Lewis BG, Barker LK, Thornton-Evans G, et al. Trends in oral health status: United States, 1988–1994 and 1999–2004. National Center for Health Statistics. *Vital Health Stat*. 2007;11:248.
36. Beikler T, Schnitzer S, Abdeen G, Ehmke B, Eisenacher M, Flemmig TF. Sampling strategy for intraoral detection of periodontal pathogens before and following periodontal therapy. *J Periodontol*. 2006;77:1323–3132.
37. Flemmig TF, Arushanov D, Daubert D, Rothen M, Mueller G, Leroux BG. Randomized controlled trial assessing efficacy and safety of glycine powder air polishing in moderate to deep periodontal pockets. *J Periodontol*. 2012;83:444–52.
38. Meyer F, Paarmann D, D'Souza M, Olson R, Glass EM, Kubal M, et al. The metagenomics RAST server - a public resource for the automatic phylogenetic and functional analysis of metagenomes. *BMC Bioinf*. 2008;9:386.
39. Legendre P, Legendre L.F.J. Numerical ecology, Volume 24. 3rd ed. Oxford: Elsevier; 2012.
40. Di Bella JM, Bao Y, Gloor GB, Burton JP, Reid G. High throughput sequencing methods and analysis for microbiome research. *J Microbiol Methods*. 2013;95:401–14.
41. Slots J. Selection of antimicrobial agents in periodontal therapy. *J Periodontol Res*. 2002;37:389–98.
42. Anderson MJ. A new method for non-parametric multivariate analysis of variance. *Austral Ecol*. 2001;26:32–46.
43. McArdle BH, Anderson MJ. Fitting multivariate models to community data: a comment on distance-based redundancy analysis. *Ecology*. 2001;82:290–7.
44. Belda-Ferre P, Alcaraz LD, Cabrera-Rubio R, Romero H, Simón-Soro A, Pignatelli M, et al. The oral metagenome in health and disease. *ISME J*. 2012;6:46–56.
45. Griffen AL, Beall CJ, Campbell JH, Firestone ND, Kumar PS, Yang ZK, et al. Distinct and complex bacterial profiles in human periodontitis and health revealed by 16S pyrosequencing. *ISME J*. 2012;6:1176–85.
46. Keijser BJ, Zaura E, Huse SM, van der Vossen JM, Schuren FH, Montijn RC, et al. Pyrosequencing analysis of the oral microflora of healthy adults. *J Dent Res*. 2008;87:1016–20.
47. Kumar PS, Mason MR, Brooker MR, O'Brien K. Pyrosequencing reveals unique microbial signatures associated with healthy and failing dental implants. *J Clin Periodontol*. 2012;39:425–33.
48. Duran-Pinedo AE, Chen T, Teles R, Starr JR, Wang X, Krishnan K, et al. Community-wide transcriptome of the oral microbiome in subjects with and without periodontitis. *ISME J*. 2014;8:1659–72.
49. Cortelli JR, Aquino DR, Cortelli SC, Fernandes CB, de Carvalho-Filho J, Franco GC, et al. Etiological analysis of initial colonization of periodontal pathogens in oral cavity. *J Clin Microbiol*. 2008;46:1322–9.
50. Davies RM, Ellwood RP, Davies GM. The effectiveness of a toothpaste containing triclosan and polyvinyl-methyl ether maleic acid copolymer in improving plaque control and gingival health: a systematic review. *J Clin Periodontol*. 2004;31:1029–33.
51. Hajishengallis G, Lamont RJ. Beyond the red complex and into more complexity: the polymicrobial synergy and dysbiosis (PSD) model of periodontal disease etiology. *Mol Oral Microbiol*. 2012;27:409–19.
52. Saygun I, Nizam N, Keskiner I, Bal V, Kubar A, Açikol C, et al. Salivary infectious agents and periodontal disease status. *J Periodontol Res*. 2011;46:235–9.
53. Teles R, Sakellari D, Teles F, Konstantinidis A, Kent R, Socransky S, et al. Relationships among gingival crevicular fluid biomarkers, clinical parameters of periodontal disease, and the subgingival microbiota. *J Periodontol*. 2010;81:89–98.
54. Socransky SS, Haffajee AD. Periodontal microbial ecology. *Periodontol* 2000. 2005;38:135–87.
55. Ge X, Rodriguez R, Trinh M, Gunsolley J, Xu P. Oral microbiome of deep and shallow dental pockets in chronic periodontitis. *PLoS One*. 2013;8, e65520.
56. Amend AS, Seifert KA, Bruns TD. Quantifying microbial communities with 454 pyrosequencing: does read abundance count? *Mol Ecol*. 2010;19:5555–65.
57. Harrison A, Binder H, Buhot A, Burden CJ, Carlon E, Gibas C, et al. Physico-chemical foundations underpinning microarray and next-generation sequencing experiments. *Nucleic Acids Res*. 2013;41:2779–96.
58. Jolliffe IT. Principal component analysis. New York: Springer; 2002.
59. Nasidze I, Li J, Quinque D, Tang K, Stoneking M. Global diversity in the human salivary microbiome. *Genome Res*. 2009;19:636–43.
60. Lourenço TG, Heller D, Silva-Boghossian CM, Cotton SL, Paster BJ, Colombo AP. Microbial signature profiles of periodontally healthy and diseased patients. *J Clin Periodontol*. 2014;41:1027–36.
61. Frias-Lopez J, Duran-Pinedo A. Effect of periodontal pathogens on the metatranscriptome of a healthy multispecies biofilm model. *J Bacteriol*. 2012;194:2082–95.
62. Paster BJ, Olsen I, Aas JA, Dewhirst FE. The breadth of bacterial diversity in the human periodontal pocket and other oral sites. *Periodontol* 2000. 2006;42:80–7.
63. Song Y, Garg S, Girotra M, Maddox C, von Roseninge EC, Dutta A, et al. Microbiota dynamics in patients treated with fecal microbiota transplantation for recurrent *Clostridium difficile* infection. *PLoS One*. 2013;8, e81330.
64. Haffajee AD, Teles RP, Socransky SS. The effect of periodontal therapy on the composition of the subgingival microbiota. *Periodontol* 2000. 2006;42:219–58.
65. Zandbergen D, Slot DE, Cobb CM, Van der Weijden FA. The clinical effect of scaling and root planning and the concomitant administration of systemic amoxicillin and metronidazole: a systematic review. *J Periodontol*. 2013;84:332–51.
66. De Soete M, Mongardini C, Peuwels M, Haffajee A, Socransky S, van Steenberghe D, et al. One-stage full-mouth disinfection. Long-term microbiological results analyzed by checkerboard DNA-DNA hybridization. *J Periodontol*. 2001;72:374–82.
67. Ehmke B, Moter A, Beikler T, Milián E, Flemmig TF. Adjunctive antimicrobial therapy of periodontitis, long-term effects on disease progression and oral colonization. *J Periodontol*. 2005;76:749–59.
68. Lobene RR, Soparkar PM, Hein JW, Quigley GA. A study of the effects of antiseptic agents and a pulsating irrigating device on plaque and gingivitis. *J Periodontol*. 1972;43:564–8.
69. Cousido MC, Tomás Carmona I, García-Caballero L, Limeres J, Alvarez M, Diz P. *In vivo* substantivity of 0.12 % and 0.2 % chlorhexidine mouthrinses on salivary bacteria. *Clin Oral Investig*. 2010;4:397–402.
70. Adcock JE, Berry Jr WC, Kalkwarf KL. Effect of sodium hypochlorite solution on the subgingival microflora of juvenile periodontitis lesions. *Pediatr Dent*. 1983;5:190–4.
71. Spratt DA, Pratten J, Wilson M, Gulabivala K. An *in vitro* evaluation of the antimicrobial efficacy of irrigants on biofilms of root canal isolates. *Int Endod J*. 2001;34:300–7.
72. Galván M, Gonzalez S, Cohen CL, Alonzaiz FA, Chen CT, Rich SK, et al. Periodontal effects of 0.25 % sodium hypochlorite twice-weekly oral rinse. A pilot study. *J Periodontol Res*. 2014;49:696–702.
73. Kellermayer R, Nagy-Szakal D, Harris RA, Luna RA, Pitashny M, Schady D, et al. Serial fecal microbiota transplantation alters mucosal gene expression in pediatric ulcerative colitis. *Am J Gastroenterol*. 2015;110:604–6.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

