

1 **A molecular arms race between host innate antiviral response and emerging human coronaviruses**

2

3 Lok-Yin Roy Wong, Pak-Yin Lui and Dong-Yan Jin

4

5 School of Biomedical Sciences, The University of Hong Kong, Pokfulam, Hong Kong

6

7

8

9

10

11

12

13

14

15

16

17

18 **Correspondence:** DY Jin, School of Biomedical Sciences, The University of Hong Kong, 3/F Laboratory

19 Block, Faculty of Medicine Building, 21 Sassoon Road, Pokfulam, Hong Kong. Phone: +852-3917-9491;

20 Fax: +852-2855-1254; E-mail: [dyjin@hku.hk](mailto:dyjin@hku.hk).

21 **Abstract**

22 Coronaviruses have been closely related with mankind for thousands of years. Community-acquired human  
23 coronaviruses have long been recognized to cause common cold. However, zoonotic coronaviruses are now  
24 becoming more a global concern with the discovery of highly pathogenic severe acute respiratory syndrome  
25 (SARS) and Middle East respiratory syndrome (MERS) coronaviruses causing severe respiratory diseases.  
26 Infections by these emerging human coronaviruses are characterized by less robust interferon production.  
27 Treatment of patients with recombinant interferon regimen promises beneficial outcomes, suggesting that  
28 compromised interferon expression might contribute at least partially to the severity of disease. The  
29 mechanisms by which coronaviruses evade host innate antiviral response are under intense investigations.  
30 This review focuses on the fierce arms race between host innate antiviral immunity and emerging human  
31 coronaviruses. Particularly, the host pathogen recognition receptors and the signal transduction pathways  
32 to mount an effective antiviral response against SARS and MERS coronavirus infection are discussed. On  
33 the other hand, the counter-measures evolved by SARS and MERS coronaviruses to circumvent host  
34 defense are also dissected. With a better understanding of the dynamic interaction between host and  
35 coronaviruses, it is hoped that insights on the pathogenesis of newly-identified highly pathogenic human  
36 coronaviruses and new strategies in antiviral development can be derived.

## 37 Introduction

38 Coronaviruses (CoVs) are classified into four genera, namely *alpha-*, *beta-*, *gamma-* and *deltacoronavirus*,  
39 under the family of *Coronaviridae* and the order of *Nidovirales* (Woo et al., 2012). The first three genera  
40 were previously known as groups I, II and III, respectively (Lau et al., 2006; Zhong et al., 2012). CoVs  
41 have been shown to infect many different hosts including bats, birds, dogs, mice and human (Woo et al.,  
42 2009; de Groot et al., 2013). The infections are commonly zoonotic in nature (Chan et al., 2013). In the past  
43 50 years, several human CoVs (HCoVs) were identified. HCoV-229E and HCoV-OC43, belonging to  
44 *alpha-* and *betacoronavirus* respectively, were the first two HCoVs identified in the mid-1960s (Tyrrell and  
45 Bynoe, 1965; Hamre and Procknow, 1966; McIntosh et al., 1967). Healthy individuals infected with either  
46 HCoV-OC43 or HCoV-229E develop illnesses within the range of typical common colds with good  
47 prognosis (Bradburne et al., 1967). Since the identification of these two HCoVs, extensive studies were  
48 conducted to understand their pathogenicity. However, almost all studies showed that HCoV-OC43 and  
49 HCoV-229E caused mild illnesses with high titers of neutralizing antibodies (Bradburne et al., 1967). The  
50 idea of HCoV being a relatively weak respiratory disease-causing agent was therefore presented to the field.

51 This idea was generally accepted until the outbreak of SARS in 2003. SARS-CoV was the first HCoV  
52 identified to cause acute respiratory distress syndrome (ARDS) (Cheng et al., 2007; Graham et al., 2013).  
53 According to World Health Organization (WHO), a total of 8096 cases from 29 countries were reported  
54 with a case mortality rate of 9.6%. The SARS outbreak changed the landscape of CoV studies entirely and  
55 marked the new era of combating infectious diseases. Tremendous efforts have been put into understanding  
56 SARS-CoV pathogenicity, opening a new page of CoV biology. Despite advances in infection control and  
57 quarantine measures in the past decade, another HCoV causing ARDS was identified in Saudi Arabia as a  
58 novel lineage C *betacoronavirus* in September 2012 (Zaki et al., 2012). The newly identified HCoV was  
59 later named MERS-CoV. Up to October 2015, 1611 laboratory-confirmed cases were reported to WHO  
60 with 575 related deaths in 26 countries, including a recent outbreak involving 186 cases and 37 deaths in

61 South Korea. MERS-CoV is closely related phylogenetically to two bat CoVs, HKU4 and HKU5, shedding  
62 light on the possible zoonotic reservoir of MERS-CoV (Zaki et al., 2012; Memish et al., 2013).

63 Together with HCoV-HKU1 identified in 2005 (Woo et al., 2005) and HCoV-NL63 discovered in 2004  
64 (Fouchier et al., 2004; van der Hoek et al., 2004), 6 HCoVs have been documented up to date. These 6  
65 HCoVs present diseases with a range of clinical severity from typical common cold in HCoV-OC43,  
66 HCoV-229E, HCoV-HKU1 and HCoV-NL63 to ARDS in SARS-CoV and MERS-CoV. Why these CoVs  
67 show dramatically different pathogenicity in human is an important but unanswered question in the field.  
68 One model to explain this difference is based on adaptation and host immunity. According to this model,  
69 bats are reservoir of various CoVs. Bat CoVs constantly emerge in human via intermediate hosts such as  
70 civets and dromedaries. Exposure of immunologically naïve human populations to these CoVs commonly  
71 causes severe diseases plausibly due to aberrant activation of innate immunity and lack of immune memory.  
72 When some CoVs become better adapted in human by acquiring the ability to transmit from human to  
73 human readily, pandemics could arise. Meanwhile, as they become fully adapted, the CoVs might only  
74 cause mild diseases in human. Existing evidence supports the origin of HCoV-OC43, HCoV-229E, HCoV-  
75 HKU1 and HCoV-NL63 from bats and other animals (Woo et al., 2009; Huynh et al., 2012; Corman et al.,  
76 2015). Adaptation and virus-host interaction are also known to be major determinants in CoV pathogenesis  
77 (Pepin et al., 2010; Chan et al., 2013). It will therefore be of great interest to see whether emerging human  
78 CoVs might be particularly capable of evading innate antiviral response while activating pathological  
79 inflammation. In other words, we need to determine whether the more severe clinical presentations might  
80 be accounted for by the specific interaction between host and emerging human CoVs, namely SARS-CoV  
81 and MERS-CoV. In this review, the host innate antiviral response to CoV infection is particularly focused.  
82 In addition, the viral strategies adopted by SARS-CoV and MERS-CoV to subvert innate immunity are also  
83 summarized to provide inspiring insights that may explain the discrepancies in virulence (Figure 1).

84

## 85 An overview of CoV biology

86 CoVs are polycistronic positive-sense single-stranded RNA (ssRNA) viruses with genomes of about 30kb  
87 in size. The 5' most two-thirds of CoV genome encodes polyprotein 1a (pp1a) and pp1ab replicase  
88 polyproteins, which are further cleaved by viral proteases to yield non-structural proteins (nsps), while the  
89 3' end of the genome encodes structural and lineage-specific proteins (Durai et al., 2015). The CoV life  
90 cycle begins with the binding to cellular receptor followed by membrane fusion as well as viral RNA and  
91 protein synthesis in the cytoplasm. The pp1a and pp1ab polyproteins are co-translationally processed  
92 resulting in the formation of the replicase complex. A set of nested subgenomic mRNAs and genomic RNA,  
93 which possess both the same 3' end and a common 5' leader sequence derived from the 5' end of the genome,  
94 is then transcribed. Normally, only the 5' end of each mRNA is translated. Virion assembly is achieved by  
95 budding into intracellular membranes and virion release is accomplished through the secretory pathway  
96 (Cheng et al., 2007; Durai et al., 2015).

97 The coronaviral spike (S) protein is responsible for binding to specific host receptor on cell surface and  
98 fusing viral envelope with lipid membrane of host upon infection (Bosch et al., 2003; Rota et al., 2003;  
99 Chen et al., 2013). HCoV-NL63 and SARS-CoV from  $\alpha$ - and  $\beta$ -genera respectively recognize angiotensin-  
100 converting enzyme 2 (ACE2) (Li et al., 2003; Pyrc et al., 2007; Frieman et al., 2008; Chen et al., 2013)  
101 while MERS-CoV infects cells through another cell surface enzyme dipetidyl peptidase 4 (DPP4) (Chen et  
102 al., 2013; Raj et al., 2013). Aminopeptidase N (APN) has also been found to be recognized by some  $\alpha$ -  
103 genus CoVs like HCoV-229E (Yeager et al., 1992). Cell surface receptor binding dictates species-specific  
104 viral entry as well as tropism. This also confines the direction of cellular antiviral response. We and others  
105 have shown the ability of CoV S proteins to activate unfolded protein response and endoplasmic reticulum  
106 stress (Chan et al., 2006; Fung et al., 2014; Siu et al., 2014b). The activity of S might also be functionally  
107 related to coronaviral perturbation of innate antiviral response including IFN and cytokine production.

108

**109 Detection of CoV by host innate immune sensors**

110 Pattern recognition receptors (PRRs) constitute an indispensable part of the host innate immune defense  
111 mechanism by the detection of foreign, non-self patterns from invading microbes distinct from host. These  
112 pathogen-associated molecular patterns (PAMPs) are usually biomolecules derived from the surface or  
113 generated during the life cycle of the microbes. The detection of PAMPs by host PRRs activates innate  
114 immune response including the expression of type I IFNs and cytokines for clearance of invading microbes.  
115 During CoV infection, retinoic acid-inducible gene I (RIG-I)-like receptors (RLRs) and Toll-like receptors  
116 (TLRs) are believed to bear pivotal importance in stimulating host type I IFN induction. It is therefore  
117 essential to review the sensing mechanism of the PRRs to understand viral evasion mechanisms and provide  
118 insights on the development of potential viral antagonists.

**119 *RIG-I-like receptors***

120 After viral entry, CoV genomes are exposed in the cytoplasm for expression of viral proteins, providing an  
121 opportunity for viral RNA sensing by host. RLRs are ubiquitously expressed cytoplasmic RNA helicases  
122 of DExD/H box family responsible for sensing double-stranded RNA (dsRNA) (Yoneyama et al., 2005).  
123 Three types of RLRs have been identified up to now, including RIG-I, melanoma differentiation-associated  
124 gene 5 (MDA5) and laboratory of genetics and physiology 2 (LGP2) (Loo and Gale, 2011). RIG-I and  
125 MDA5 consist of N-terminal caspase activation and recruitment domain (CARD) in two tandem copies, a  
126 central DExD/H box helicase domain and a C-terminal domain (CTD) (Yoneyama et al., 2004, 2005). The  
127 N-terminal CARDS are the effector domain of RLRs to mediate downstream transduction, which is held by  
128 the CTD when unstimulated (Jiang et al., 2011; Kowalinski et al., 2011; Luo et al., 2011). However, in the  
129 presence of residual amount of cytoplasmic dsRNA, RLRs bind to dsRNA through the central DExD/H  
130 box helicase domain and CTD with ATP, causing a conformational change that exposes the N-terminal  
131 CARDS for signal transduction (Yoneyama et al., 2004; Jiang et al., 2011). LGP2 lacking the N-terminal  
132 CARDS is thought to act as co-factor that augments the function of RIG-I and MDA5 (Sato et al., 2010;

133 Bruns et al., 2014). Exposure of CARDs leads to oligomerization of RIG-I or MDA5 to form filamentous  
134 structure (Berke et al., 2012; Peisley et al., 2013; Wu et al., 2013). The CARD filament recruits and further  
135 initiates similar filamentous structure formation of CARD on MAVS, an adaptor protein which further  
136 recruits downstream effectors tumor necrosis factor receptor-associated factor 3 (TRAF3), TANK-binding  
137 kinase 1 (TBK1) and I $\kappa$ B kinase  $\epsilon$  (IKK $\epsilon$ ) (Loo and Gale, 2011; Wu et al., 2014). TBK1 and IKK $\epsilon$  form a  
138 complex of activated protein kinase for phosphorylation and activation of not only MAVS adaptor (Liu et  
139 al., 2015a), but also IRF3 transcription factor (Loo and Gale, 2011). Activated IRF3 are phosphorylated,  
140 dimerized and eventually translocated to the nucleus. On the other hand, TRAF2/6 is also recruited to  
141 MAVS for NF- $\kappa$ B activation. Specifically, canonical NF- $\kappa$ B inhibitor I $\kappa$ B is phosphorylated and then  
142 degraded through proteasomes in a ubiquitination-dependent fashion (Loo and Gale, 2011). I $\kappa$ B  
143 degradation exposes nuclear localization signal on NF- $\kappa$ B dimer for nuclear translocation. Activated IRF3  
144 and NF- $\kappa$ B together with other transcription factors including c-Jun assemble the enhanceosome that binds  
145 to IFN- $\beta$  promoter for IFN- $\beta$  expression (Ford et al., 2010; Loo and Gale, 2011). Infection with mouse  
146 hepatitis virus induces RIG-I expression. In addition, the activation of type I IFN production by this CoV  
147 in oligodendrocytes requires both RIG-I and MDA5 (Li et al., 2010). Thus, RLRs might play an important  
148 role in the sensing of CoV infection.

149 Several critical questions concerning RLR recognition of CoVs merit further investigations. First, the role  
150 of RLRs in CoV sensing should be studied in RLR-null and CoV-susceptible cells and animals. When  
151 necessary CRISPR/Cas9 technology might be used to disrupt RLR genes in target cells (Hsu et al., 2014;  
152 Yuen et al., 2015). Second, the CoV PAMPs recognized by RLRs should be identified and characterized.  
153 Particularly, it will be of interest to see whether and how common and highly structured regions in  
154 coronaviral genome, such as the aforementioned 5' leader sequence, might be recognized by RLRs. For  
155 example, a polyuridine motif in the 3' untranslated region of hepatitis C virus genome and the panhandle  
156 structure in RNA viruses such as influenza A virus have previously been shown to be RIG-I agonists (Saito  
157 et al., 2008; Weber et al., 2013; Kell et al., 2015; Liu et al., 2015b). In addition, possible involvement of

158 viral proteins such as nucleocapsid (N) in this recognition as in the case of other RNA viruses (Saito et al.,  
159 2008; Weber et al., 2013) should also be clarified. Finally, comparative analysis of SARS-CoV, MERS-  
160 CoV and other HCoVs for their ability to activate RLRs will shed light on whether RLR activation would  
161 be a critical determinant in CoV virulence.

### 162 *Toll-like receptors*

163 CoVs have been observed to infect host cells through more than one pathway. While CoV entry by the  
164 fusion of viral envelope and host membrane has been described, the endosomal pathway is still considered  
165 the classical entry pathway for CoVs. In this pathway the activation of S protein cleavage by cathepsin L  
166 and transmembrane serine protease TMPRSS2 occurs in the absence of cell surface proteases in certain cell  
167 types (Shirato et al., 2013; Burkard et al., 2014). In this regard, TLR family may play an essential role in  
168 sensing CoV infection through the endosomal pathway. TLR family was identified as another PRR  
169 homologous to *Drosophila* Toll receptor (Boehme and Compton, 2004), sensing various PAMPs within the  
170 endosome which leads to induction of cytokines and IFNs. In human, each of the 11 TLRs is known to  
171 specifically recognize a particular PAMP and preferentially resides in either plasma or endosomal  
172 membrane. The cellular localization of TLRs defines their functions in detecting different PAMPs. For  
173 example, TLRs critically involved in viral nucleic acid sensing, including TLR3 for dsRNA, TLR7 and  
174 TLR8 for ssRNA, and TLR9 for unmethylated CpG island of dsDNA viruses, are mainly localized in  
175 endosomal membrane while other members having a role in sensing other biomolecules derived from  
176 microbial surface components localized to plasma membrane of infected cells (Xagorari and Chlichlia,  
177 2008; Kawai and Akira, 2010). TLR family members being type 1 transmembrane proteins share a similar  
178 structure with a single transmembrane domain. TLR specificity is determined by the ectodomain made up  
179 of various number of leucine-rich repeats (LRRs) that bind the corresponding PAMP directly (Boehme and  
180 Compton, 2004). Signal transduction begins with ligand binding to LRRs in the ectodomain, thus recruiting  
181 cytosolic adaptor protein MyD88 with cytoplasmic Toll/IL-1 receptor (TIR) domain by homotypic TIR-  
182 TIR domain interaction (Xagorari and Chlichlia, 2008). The TLR-MyD88 complex then recruits and



183 activates interleukin 1R-associated kinase (IRAK) by phosphorylation. The activated IRAK then in turn  
184 associates with TRAF6 and activates a series of downstream effectors leading to the activation of a range  
185 of cytokines and IFN-stimulated genes (ISGs), while activation of type I IFN expression by TLR3 is  
186 independent of MyD88 but dependent on TRIF (Boehme and Compton, 2004; Xagorari and Chlichlia,  
187 2008). TLR pathway is significantly involved in the suppression of CoV replication and induction of type  
188 I IFN expression. Mice deficient of either TLR3 or TLR4 were more prone to SARS-CoV pathogenesis  
189 (Mazaleuskaya et al., 2012; Totura et al., 2015). Notably, disruption of either MyD88 or TRIF arm of the  
190 TLR signaling pathway causes lethal SARS-CoV disease, indicating the importance of both arms in host  
191 innate immunity against SARS-CoV (Totura et al., 2015). Full characterization of the role of TLRs in host  
192 innate antiviral response against SARS-CoV and MERS-CoV versus other HCoVs will not only provide  
193 new knowledge about how TLR activation might impact CoV pathogenesis, but might also identify new  
194 strategies for antiviral and vaccine development. For example, synthetic TLR agonists could potentially  
195 serve as antivirals and vaccine adjuvants in the prevention and control of CoVs.

196

### 197 **Host innate immune response against CoV infection**

198 Innate antiviral response is the first line of defense against CoV infection. Type I IFNs are important  
199 antiviral and immunomodulatory agents. Type I IFNs function by binding to IFN- $\alpha$  receptor-1 (IFNAR-1)  
200 and IFNAR-2 receptor complex, thus activating *Janus* family tyrosine kinase (JAK), leading to the  
201 phosphorylation of signal transducer and activator of transcription (STAT), a family of transcription factors  
202 regulating the expression of ISGs. Activated STAT and IRF9 form IFN-stimulated gene factor 3 (ISGF3),  
203 stimulating expression of ISGs by binding to IFN-stimulated response element (ISRE) in promoters of ISGs  
204 (Levy et al., 2001; Samuel, 2001). Viral induction of ISGs was abrogated in STAT1<sup>-/-</sup> mice infected with  
205 SARS-CoV. The viral infection could not be cleared resulting in severe disease, extensive lung injury and

206 100% mortality (Frieman et al., 2010; Zornetzer et al., 2010). This indicates the importance of STAT1 in  
207 SARS-CoV pathogenesis.

208 ISGs are the workhorses of the innate antiviral response with diverse functions including direct antiviral  
209 activities and regulation of adaptive immune system (Schneider et al., 2014). For example, IFN-inducible  
210 gene *p53* evokes apoptosis in virus-infected cells (Takaoka et al., 2003). IFN-inducible protein kinase PKR,  
211 2', 5'-oligoadenylate synthetase (OAS) and RNase L are important modulators involved in dsRNA sensing,  
212 viral gene expression and replication. They act sequentially to trigger viral RNA degradation and  
213 suppression of viral activities (Samuel, 2001). Other ISGs encoding antiviral effectors such as Mx proteins,  
214 cholesterol-25-hydroxylase, IFITM proteins, TRIM proteins, viperin, tetherin, cGAMP synthase and STING  
215 could also be highly relevant to CoV infection (Schneider et al., 2014; Schoggins et al., 2014; Ma et al.,  
216 2015a; Ma et al., 2015b). Inflammatory responses triggered by inflammatory cytokines like tumor necrosis  
217 factor  $\alpha$  (TNF- $\alpha$ ) and IFN- $\gamma$  are also found to be IFN-dependent (Samuel, 2001). IFNs do not only exert  
218 antiviral effects through activation of innate immunity but also act as modulators of adaptive immunity.  
219 Adaptive immune response is activated by increased level of IFNs. The levels of major histocompatibility  
220 complex (MHC) proteins class I and II are found up-regulated by IFNs. This facilitates efficient antigen  
221 presentation and hence cellular immune response to CoV infection (Samuel, 1991, 2001; Ivashkiv and  
222 Donlin, 2014). In addition, the roles of non-conventional ISGs including microRNAs, long non-coding  
223 RNAs and alternatively spliced isoforms have been increasingly recognized in recent years (Schneider et  
224 al., 2014). It will be of importance to determine whether SARS-CoV and MERS-CoV might be unique in  
225 ISG activation as suggested in a recent study, which demonstrated that MERS-CoV induces repressive  
226 histone modifications to down-regulate specific subsets of ISGs (Menachery et al., 2014b). In relation to  
227 this, two areas concerning ISG activation by CoVs might require more attention and research efforts. First,  
228 unbiased and large-scale screening of antiviral ISGs using RNA interference or CRISPR/Cas9 technology  
229 might be carried out to identify key cellular factors that restrict SARS-CoV and MERS-CoV replication  
230 and infection. Second, small-molecule compounds that activate antiviral ISGs could be identified and tested

231 for inhibition of SARS-CoV and MERS-CoV replication and infection. For example, establishing the  
232 significance of cGAS and STING in CoV infection might lead to the development of cyclic dinucleotides  
233 such as c-di-GMP and cGAMP as novel anti-CoV agents.

234

### 235 **Evasion of innate immune response by CoV**

236 CoVs have been reported to directly or indirectly suppress IFN production and signaling pathways by a  
237 subset of viral proteins via various mechanisms. In many cases, infected patients have shown diminished  
238 levels of type I IFNs. This is especially true for SARS and MERS patients with severe diseases (Faure et  
239 al., 2014). It was also shown that SARS-CoV and MERS-CoV were capable of evading type I IFN  
240 production and signaling to different extents in cultured cells (Kindler et al., 2013). When the deficiency in  
241 type I IFN production in CoV-infected cells was remedied by IFN- $\alpha$  treatment, CoV replication was  
242 inhibited (Falzarano et al., 2013). Combination of IFN- $\alpha$  with other antiviral drugs further improves the  
243 survival of infected patients (Omrani et al., 2014). This evidence suggests an essential role of type I IFNs  
244 in the antiviral effect against CoV infection. CoVs have evolved strategies to counter host antiviral response  
245 by antagonizing type I IFN production and signaling. CoV proteins have been characterized to exhibit  
246 innate immunosuppressive effects in cellular models. Below we will discuss them in three categories:  
247 structural, lineage-specific and non-structural proteins (nsps) (de Groot et al., 2013). Nsps of CoVs are  
248 involved in the assembly of the replicase complex for viral RNA synthesis (Sevajol et al., 2014). Certain  
249 nsps have also been reported to possess innate immunosuppressive effect that facilitates viral replication  
250 and propagation, although these proteins *per se* are not required for viral life cycle (Narayanan et al., 2008b;  
251 Lokugamage et al., 2015). Nsps of different CoVs are more or less evolutionarily conserved suggesting  
252 their functional significance, with the exception of nsp1 and nsp2, which are thought to contribute to  
253 virulence of certain CoVs (Neuman et al., 2014). Four structural proteins are found in CoVs, namely S,  
254 membrane (M), envelope (E) and N proteins. Structural proteins contribute the architecture for virion

255 assembly. Accessory proteins are lineage-specific with diverse behaviors in different CoVs but are not  
256 essential for viral replication and propagation (de Groot et al., 2013).

257

258 CoV nsps have shown suppressive effects in various immune pathways including type I IFN production  
259 and signaling. SARS-CoV and MERS-CoV nsp1 proteins have been shown to selectively induce  
260 degradation of host mRNA by inducing endonucleolytic cleavage while leaving viral RNAs intact (Huang  
261 et al., 2011; Lokugamage et al., 2015). In addition to the induction of endonucleolytic cleavage of host  
262 mRNA, general inhibition of host mRNA translation is achieved by binding of 40S subunit of ribosome  
263 with SARS-CoV nsp1 (Huang et al., 2011). Particularly, SARS-CoV nsp1 inhibits innate immune response  
264 by translational repression of IFN mRNA transcripts, hence altering IFN production and signaling  
265 (Narayanan et al., 2008a; Tanaka et al., 2012). MERS-CoV nsp1 has also been characterized to specifically  
266 induce endonucleolytic cleavage of nuclear transcribed mRNA while sparing cytoplasmic host mRNA and  
267 viral RNA (Lokugamage et al., 2015). This suggests a novel mechanism for evading host immune response.

268 CoV nsp3 protein has been characterized with a papain-like protease (PLpro) domain for enzymatic  
269 cleavage of pp1a and pp1ab as well as a PLP2 domain with deubiquitinating and deISGylating activity  
270 (Clementz et al., 2010; Mielech et al., 2014). MERS-CoV PLpro is able to antagonize IFN production  
271 induced by RIG-I and MDA5 as well as NF- $\kappa$ B activation (Mielech et al., 2014). MERS-CoV PLpro is  
272 catalytically more efficient (Báez-Santos et al., 2014) and its catalytic activity is indispensable for the  
273 suppressive effect on RIG-I, MDA5 and NF- $\kappa$ B (Mielech et al., 2014). In contrast, SARS-CoV PLpro does  
274 not require enzymatic activity for IFN antagonism (Clementz et al., 2010). HCoV-NL63 and SARS-CoV  
275 PLP2 transmembrane domain can also act as potent IFN antagonists to suppress IFN production induced  
276 by RIG-IN, a dominant active form of RIG-I (Clementz et al., 2010). In another view of direct inhibition  
277 of IFN induction, nsp3 with deubiquitinating and deISGylating activity may also influence the  
278 ubiquitination and ISGylation pattern and dynamics thus indirectly hindering innate immune response

279 against CoV infections (Clementz et al., 2010). For example, ISGylation and ubiquitination of IRF3  
280 required for optimal activation is probably altered by PLP domain of nsp3.

281 Apart from directly manipulating the signaling pathway involved in IFN production, several CoV nsps were  
282 identified to act on viral RNA to minimize IFN stimulation. N7-methylguanosine is the fundamental moiety  
283 of eukaryotic mRNA cap structure and 2'-O-methylation on this moiety is a representative host signature  
284 to avoid PRR activation as well as ISG action. Particularly, viral RNA with this modification evades  
285 recognition by MDA5 or IFIT family antiviral factors (Züst et al., 2011; Daffis et al., 2010). This is a  
286 common immunoevasive mechanism adopted by not only different CoVs but also other RNA viruses.  
287 Functional screening in yeasts suggested a novel function of SARS-CoV nsp14 as a guanine-N7-  
288 methyltransferase, the activity of which is required for viral replication and transcription (Chen et al., 2009).  
289 Another nsp of SARS-CoV, nsp16, also possesses 2'-O-methyltransferase activity (Menachery et al., 2014a;  
290 Menachery et al., 2014c). Structural modeling suggested that SARS-CoV nsp16 associates with nsp10 in  
291 1:1 ratio to form a complex of mature 2'-O-methyltransferase for viral cap methylation (Chen et al., 2011;  
292 Decroly et al., 2011). A short peptide derived from nsp10 conserved region has been shown to be a  
293 promising nsp16 antagonist which outcompetes native nsp10 to blunt 2'-O-methyltransferase activity and  
294 restrict viral replication (Wang et al., 2015). Plausibly, CoV nsps might execute their innate  
295 immunosuppressive roles by targeting type I IFN production and signaling. Further investigations are  
296 required to clarify whether and how far the sensing of CoV RNA and the induction of innate antiviral  
297 response are involved in the inhibitory activity of the nsp antagonists on CoV replication.

298

299 CoV structural proteins have been shown to inhibit IFN production and signaling at multiple levels. SARS-  
300 CoV N protein showed inhibitory effects on IFN production induced by Sendai virus and dsRNA analogue  
301 poly(I:C) but no inhibition could be observed when downstream signaling molecules of TLR and RLR  
302 pathway were overexpressed. Truncation mutant of N protein shows that the C-terminal domain is critical

303 for RNA-binding and IFN-antagonizing effect (Lu et al., 2011). This suggests SARS-CoV N may interfere  
304 with RNA recognition by host immune sensors such as RIG-I and MDA5 thus achieving suppressive role  
305 in IFN production. Other than N protein, SARS-CoV M protein has been characterized to potently down-  
306 regulate IFN production by impeding the formation of TRAF3·TANK·TBK1/IKK $\epsilon$  complex through the  
307 first transmembrane domain (Siu et al., 2009, 2014a). SARS-CoV M protein inhibits IFN production  
308 possibly through a sequestration model in which components of TRAF3·TANK·TBK1/IKK $\epsilon$  complex, an  
309 active complex for IRF3 phosphorylation, are sequestered to specific locations in the cell (Siu et al., 2009).  
310 SARS-CoV M protein therefore exerts its inhibitory effects by impeding the formation of  
311 TRAF3·TANK·TBK1/IKK $\epsilon$  complex but not by modulating the catalytic activity of the complex.

312 MERS-CoV M protein also exhibits IFN-antagonizing effects similar to its counterpart in SARS-CoV. In  
313 a previous study, MERS-CoV M is shown to impede IFN production by preventing IRF3 translocation into  
314 the nucleus (Yang et al., 2013). However, the detailed mechanism of inhibition remains unknown. Recently,  
315 our group has characterized the mode of inhibition of IFN production by MERS-CoV M. Consistently with  
316 previous report, we show that MERS-CoV M suppresses IFN production by preventing IRF3 activation.  
317 We showed that MERS-CoV M interacts with TRAF3 which impedes the recruitment of TBK1 to TRAF3  
318 complex. IRF3 activation and dimerization have also been hampered as a result. The inhibitory effect is at  
319 least in part accounted for by the N-terminal transmembrane domains. Despite of the similar behaviors,  
320 MERS-CoV M can only moderately suppress IFN expression when compared to SARS-CoV M.  
321 Interestingly, HCoV-HKU1 M protein does not exert any inhibitory effects on IFN production (Siu et al.,  
322 2014a), suggesting that the IFN-antagonizing activity of structural proteins is unique to each CoV but not  
323 universal. It will be of great interest to see whether this may correlate with the pathogenicity of different  
324 HCoVs.

325 Eight accessory proteins have been identified in SARS-CoV and five are found in MERS-CoV (Narayanan  
326 et al., 2008b). SARS-CoV genome encodes ORF3a, ORF3b, ORF6, ORF7a, ORF7b, ORF8a, ORF8b and  
327 ORF9b as accessory proteins (Narayanan et al., 2008b). SARS-CoV ORF3b and ORF6 have been found to

328 antagonize type I IFN production and signaling. Particularly, SARS-CoV ORF3b and ORF6 suppress IFN-  
329  $\beta$  production by perturbing IRF3 activation induced by Sendai virus infection. SARS-CoV ORF3b and  
330 ORF6 also suppress IFN- $\beta$ -induced activation of ISRE in ISG promoters (Kopecky-Bromberg et al., 2007),  
331 although they are not able to reduce the level of phosphorylation of STAT1, a transcription factor that  
332 activates ISRE activity once phosphorylated. However, SARS-CoV ORF6 has been shown to inhibit  
333 STAT1 translocation for ISRE activation (Kopecky-Bromberg et al., 2007). The findings suggest a mode  
334 of inhibition of IFN- $\beta$  signaling by SARS-CoV.

335 IFN antagonism of accessory proteins has also been observed in another deadly HCoV. MERS-CoV  
336 genome encodes ORF3, ORF4a, ORF4b, ORF5 and ORF8b (de Groot et al., 2013). Among the five  
337 accessory proteins, ORF4a, ORF4b and ORF5 show the ability to dampen IFN production (Yang et al.,  
338 2013). Suppression of IFN- $\beta$  promoter-driven luciferase activity has been observed in cells transfected with  
339 ORF4a, ORF4b and ORF5 plasmids. All these 3 accessory proteins are able to block IRF3 translocation to  
340 the nucleus to activate IFN promoter (Yang et al., 2013). MERS-CoV ORF4a shows an additional level of  
341 inhibition of innate immunity by intervening NF- $\kappa$ B activation. In another study, ORF4a has been shown  
342 as an antagonist of IFN production by inhibiting IRF3 translocation but has no effect on IFN signaling  
343 (Niemeyer et al., 2013). Our group demonstrated that MERS-CoV ORF4a interacts with PACT, a cellular  
344 dsRNA-binding protein that optimally activates RIG-I- and MDA5-induced type I IFN production, in an  
345 RNA-dependent manner (Siu et al., 2014c). This suggests that ORF4a may compete with RIG-I and MDA5  
346 for RNA, rendering the inactivation of RIG-I and MDA5. Direct interaction of ORF4a with PACT may  
347 also prevent interaction of PACT with RIG-I and MDA5, thus compromising PACT-dependent activation  
348 of RIG-I and MDA5 required for optimal induction of IFN production. Although we and others have  
349 observed the IFN-antagonizing activity of MERS-CoV ORF4b, different activity profiles and mechanisms  
350 have been suggested (Yang et al., 2013; Matthews et al., 2014). One recent report suggested that ORF4b  
351 directly interacts with and inhibits TBK1/IKK $\epsilon$  in the cytoplasm but might also perturb type I IFN  
352 production in the nucleus through an unknown mechanism (Yang et al., 2015).

353 Mouse hepatitis virus, another *betacoronavirus* closely related to HCoV-OC43 and HCoV-HKU1, encodes  
354 a lineage-specific accessory protein named ns2 with innate immunosuppressive property (Zhao et al., 2012).  
355 Biochemical assays indicate that ns2 protein has phosphodiesterase activity against 2', 5'-A, the product of  
356 OAS (Zhang et al., 2013). Thus, ns2 is a potent inhibitor of an IFN effector molecule and it might represent  
357 a new family of viral and cellular proteins with innate immunosuppressive activity (Zhang et al., 2013;  
358 Gusho et al., 2014). Whether distantly related proteins in HCoV-OC43 and HCoV-HKU1 might have  
359 similar activity remains to be determined. More importantly, it will be of interest to see whether SARS-  
360 CoV and MERS-CoV might encode proteins with similar enzymatic activity.

361 Multiple IFN antagonists have been identified and characterized in SARS-CoV and MERS-CoV. Some  
362 differences between these IFN-antagonizing viral proteins and their counterparts in other CoVs such as the  
363 parental bat viruses of MERS-CoV have also been noticed (Siu et al., 2014c). Existing evidence supports  
364 several important notions. First, although SARS-CoV and MERS-CoV share some features in common,  
365 they are distinct and use unique mechanisms for innate immune evasion (Perlman and Zhao, 2013). Second,  
366 both SARS-CoV and MERS-CoV are bat-origin CoVs that are well adapted in bats but newly emerge in  
367 human. This provides a golden opportunity for the study of CoV-host interaction, CoV adaptation as well  
368 as the arms race between host innate antiviral immunity and CoVs. Observing how the arms race between  
369 the host and SARS-CoV or MERS-CoV might evolve when the viruses become adapted to human will be  
370 most revealing and could provide important clues as to how a balance of power in this arms race might  
371 result in attenuation with increased transmissibility. Finally, studies on SARS-CoV and MERS-CoV have  
372 overturned existing concepts and derived new principles and thoughts to CoV biology. Particularly,  
373 mechanisms by which SARS-CoV and MERS-CoV evade innate immunity have attracted increasing  
374 attention. However, many key issues remain obscure. Particularly, better *in vivo* evidence should be  
375 obtained to clarify whether more potent inhibition of innate IFN production and signaling by SARS-CoV  
376 and MERS-CoV is a key determinant in virulence and disease severity.



377

378 **Conclusion**

379 CoVs have drawn a lot of interests in the light of the recent emergence of MERS-CoV. It remains to be  
380 understood whether the emerging deadly CoVs causing ARDS might ultimately be established and adapted  
381 in human resulting in significant attenuation of virulence. From the identification of the first two HCoVs,  
382 HCoV-229E and HCoV-OC43 in the mid-1960s, we learned that HCoV was able to cause only common  
383 cold. However, the outbreaks of SARS and MERS that have claimed hundreds of lives revealed the other  
384 extreme of CoV pathogenicity and raised new questions in CoV biology. So far no vaccines have been  
385 developed against SARS-CoV and MERS-CoV.

386 Infection with SARS-CoV and MERS-CoV has been accompanied with suppression of innate immune  
387 response, most notably with the suppression of type I IFN production and signaling pathways. As the first-  
388 line defense in the immune system, suppression of innate immune response by these CoVs has impeded the  
389 host ability to restrict infection, causing significant casualties. Although many reports have shed light on  
390 the molecular mechanism by which various CoV proteins antagonize type I IFN production and signaling,  
391 most of the studies were performed with overexpression experiments in cellular models. Future emphasis  
392 should be put on the characterization of knock-out viruses with which the function of a particular viral gene  
393 could be studied in a more physiologically relevant context. Infectious clones and replicons for SARS-CoV  
394 and MERS-CoV have been generated for this reverse genetic approach (Yount et al., 2003; Almazán et al.,  
395 2006, 2013, 2014; Scobey et al., 2013). IFN and cytokine profiles of deadly HCoVs such as SARS-CoV  
396 and MERS-CoV can be compared with HCoV-229E and HCoV-OC43 causing mild diseases. The pivotal  
397 significance of type I IFNs in innate immune activation and modulation has been discussed in this review.  
398 Suppression pattern of IFN may provide insights on the high pathogenicity of deadly HCoVs. The arms  
399 race between host innate antiviral response and emerging human CoVs might evolve after their introduction  
400 and establishment in human populations, with significant impact on virulence, transmissibility and disease

401 severity. Emerging human CoVs remain a potential threat to global public health. New knowledge about  
402 the host-CoV arms race will provide new ideas, targets and attenuated strains for the design and  
403 development of antivirals and vaccines for prevention and control of deadly CoV infections.

404 **Acknowledgments**

405 We thank Hinson Cheung, Kitty Fung, Edwin Kong and Sam Yuen for reading the manuscript critically.  
406 Coronavirus research in our laboratory was supported by Hong Kong Health and Medical Research Fund  
407 (13121032, 14130822 and HKM-15-M01) and Hong Kong Research Grants Council (HKU1/CRF/11G,  
408 C7011-15R and T11-707/15-R).

409 **Compliance with ethics guidelines**

410 The authors declare that they have no conflict of interest. This article does not contain any studies with  
411 human or animal subjects performed by any of the authors.

412 **References**

- 413 Almazán F, DeDiego ML, Galán C, Escors D, Álvarez E, Ortego J, Sola I, Zuniga S, Alonso S, Moreno  
414 JL, Nogales A, Capiscol C, Enjuanes L. 2006. Construction of a severe acute respiratory syndrome  
415 coronavirus infectious cDNA clone and a replicon to study coronavirus RNA synthesis. *J Virol*, 80:  
416 10900-10906.
- 417 Almazán F, DeDiego ML, Sola I, Zuñiga S, Nieto-Torres JL, Marquez-Jurado S, Andrés G, Enjuanes  
418 L. 2013. Engineering a replication-competent, propagation-defective Middle East respiratory syndrome  
419 coronavirus as a vaccine candidate. *mBio*, 4: 00650-13.
- 420 Almazán, F, Sola I, Zuñiga S, Marquez-Jurado S, Morales L, Becares M, Enjuanes, L. 2014.  
421 Coronavirus reverse genetic systems: infectious clones and replicons. *Virus Res*, 189: 262-270.
- 422 Báez-Santos YM, Mielech AM, Deng X, Baker S, Mesecar AD. 2014. Catalytic function and substrate  
423 specificity of the papain-like protease domain of nsp3 from the Middle East respiratory syndrome  
424 coronavirus. *J Virol*, 88: 12511–12527.
- 425 Berke IC, Yu X, Modis Y, Egelman EH. 2012. MDA5 assembles into a polar helical filament on  
426 dsRNA. *Proc Natl Acad Sci USA*, 109: 18437-18441.
- 427 Boehme KW, Compton T. 2004. Innate sensing of viruses by Toll-like receptors. *J Virol*, 78: 7867-  
428 7873.
- 429 Bosch BJ, van der Zee R, de Haan CAM, Rottier PJM. 2003. The coronavirus spike protein is a class  
430 I virus fusion protein: structural and functional characterization of the fusion core complex. *J Virol*,  
431 77: 8801-8811.

432 Bradburne AF, Bynoe ML, Tyrrell DA. 1967. Effects of a “new” human respiratory virus in  
433 volunteers. *Br Med J*, 3: 767–769.

434 Bruns AM, Leser GP, Lamb RA, Horvath CM. 2014. The innate immune sensor LGP2 activates  
435 antiviral signaling by regulating MDA5-RNA interaction and filament assembly. *Mol Cell*, 55: 771-  
436 781.

437 Burkard C, Verheije MH, Wicht O, van Kasteren SI, van Kuppeveld FJ, Haagmans BL, Pelkmans L,  
438 Rottier PJM, Bosch BJ, de Haan CAM. 2014. Coronavirus cell entry occurs through the endo-  
439 /lysosomal pathway in a proteolysis-dependent manner. *PLoS Pathog*, 10: e1004502.

440 Chan CP, Siu KL, Chin KT, Yuen KY, Zheng B, Jin DY. 2006. Modulation of the unfolded protein  
441 response by the severe acute respiratory syndrome coronavirus spike protein. *J Virol*, 80: 9279–9287.

442 Chan JF, To KK, Tse H, Jin DY, Yuen KY. 2013. Interspecies transmission and emergence of novel  
443 viruses: lessons from bats and birds. *Trends Microbiol*, 21: 544-555.

444 Chen Y, Cai H, Pan J, Xiang N, Tien P, Ahola T and Guo D. 2009. Functional screen reveals SARS  
445 coronavirus nonstructural protein nsp14 as a novel cap N7 methyltransferase. *Proceedings of the*  
446 *National Academy of Sciences of the United States of America*, 106: 3484–3489.

447 Chen Y, Su C, Ke M, Jin X, Xu L, Zhang Z, Wu A, Sun Y, Yang Z, Tien P, Ahola T, Liang Y, Liu X  
448 and Guo D. 2011. Biochemical and Structural Insights into the Mechanisms of SARS Coronavirus RNA  
449 Ribose 2'-O-Methylation by nsp16/nsp10 Protein Complex. *PLoS Pathog*, 7: e1002294.

450

- 451 Chen Y, Rajashankar KR, Yang Y, Agnihothram SS, Liu C, Lin YL, Baric RS, Li F. 2013. Crystal  
452 structure of the receptor-binding domain from newly emerged Middle East respiratory syndrome  
453 coronavirus. *J Virol*, 87: 10777–10783.
- 454 Cheng VCC, Lau SKP, Woo PCY, Yuen KY. 2007. Severe acute respiratory syndrome coronavirus as  
455 an agent of emerging and reemerging infection. *Clin Microbiol Rev*, 20: 660-694.
- 456 Clementz MA, Chen Z, Banach BS, Wang Y, Sun L, Ratia K, Baez-Santos YM, Wang J, Takayama  
457 J, Ghosh AK, Li K, Mesecar AD, Baker SC. 2010. Deubiquitinating and interferon antagonism  
458 activities of coronavirus papain-like proteases. *J Virol*, 84: 4619-4629.
- 459 Corman VM, Baldwin HJ, Tateno AF, Zerbinati RM, Annan A, Owusu M, Nkrumah EE, Maganga GD,  
460 Oppong S, Adu-Sarkodie Y, Vallo P, da Silva Filho LVR, Leroy EM, Thiel V, van der Hoek L, Poon  
461 LLM, Tschapka CD, Drexler JF. 2015. Evidence for an ancestral association of human coronavirus  
462 229E with bats. *J Virol*, 89: 11858-11870.
- 463 Daffis S, Szretter KJ, Schriewer J, Li J, Youn S, Errett J, Lin TY, Schneller S, Züst R, Dong H, Thiel  
464 V, Pierson TC, Muller RM, Gale MJ, Shi PY and Diamond MS. 2010. 2'-O methylation of the viral  
465 mRNA cap evades host restriction by IFIT family members. *Nature*, 468: 452–456.
- 466 Decroly E, Debarnot C, Ferron F, Bouvet M, Coutard B, Imbert I, Gluais L, Papageorgiou N, Sharff A,  
467 Bricogne G, Ortiz-Lombardia M, Lescar J and Canard, B. 2011. Crystal Structure and Functional  
468 Analysis of the SARS-Coronavirus RNA Cap 2'-O-Methyltransferase nsp10/nsp16 Complex. *PLoS*  
469 *Pathog*, 7: 1002059.
- 470 de Groot RJ, Baker SC, Baric RS, Brown CS, Drosten C, Enjuanes L, Fouchier RAM, Galiano M,  
471 Gorbalenya AE, Memish ZA, Perlman S, Poon LLM, Snijder EJ, Stephens GM, Woo PCY, Zaki AM,

- 472 Zambon M, Ziebuhr J. 2013. Middle East respiratory syndrome coronavirus (MERS-CoV):  
473 announcement of the coronavirus study group. *J Virol*, 87: 7790-7792.
- 474 Durai P, Batool M, Shah M, Choi S. 2015. Middle East respiratory syndrome coronavirus: transmission,  
475 virology and therapeutic targeting to aid in outbreak control. *Exp Mol Med*, 47: e181.
- 476 Falzarano D, de Wit E, Martellaro C, Callison J, Munster VJ, Feldmann H. 2013. Inhibition of novel  $\beta$   
477 coronavirus replication by a combination of interferon- $\alpha$ 2b and ribavirin. *Sci Rep*, 3: 1686.
- 478 Faure E, Poissy J, Goffard A, Fournier C, Kipnis E, Titecat M, Bortolotti P, Martinez L, Dubucquoi S,  
479 Dessein R, Gosset P, Mathieu D, Guery B. 2014. Distinct immune response in two MERS-CoV-  
480 infected patients: Can we go from bench to bedside? *PLoS ONE*, 9: e88716.
- 481 Ford E, Thanos D. 2010. The transcriptional code of human IFN- $\beta$  gene expression. *Biochim Biophys*  
482 *Acta*, 1799: 328-336.
- 483 Fouchier RAM, Hartwig NG, Bestebroer TM, Niemeyer B, de Jong JC, Simon JH, Osterhaus ADME.  
484 2004. A previously undescribed coronavirus associated with respiratory disease in humans. *Proc Natl*  
485 *Acad Sci USA*, 101: 6212–6216.
- 486 Frieman M, Heise M, Baric R. 2008. SARS coronavirus and innate immunity. *Virus Res*, 133: 101–  
487 112.
- 488 Frieman MB, Chen J, Morrison TE, Whitmore A, Funkhouser W, Ward JM, Lamirande EW, Roberts  
489 A, Heise M, Subbarao K, Baric RS. 2010. SARS-CoV pathogenesis is regulated by a STAT1 dependent  
490 but a type I, II and III interferon receptor independent mechanism. *PLoS Pathog*, 6: e1000849.
- 491 Fung TS, Huang M, Liu DX. 2014. Coronavirus-induced ER stress response and its involvement in  
492 regulation of coronavirus–host interactions. *Virus Res*, 194: 110-123.

- 493 Graham R, Donaldson EF, Baric RS. 2013. A decade after SARS: strategies for controlling emerging  
494 coronaviruses. *Nat Rev Microbiol*, 11: 836-848.
- 495 Gusho E, Zhang R, Jha BK, Thornbrough JM, Dong B, Gaughan C, Elliott R, Weiss SR, Silverman  
496 RH. 2014. Murine AKAP7 has a 2', 5'-phosphodiesterase domain that can complement an inactive  
497 murine coronavirus ns2 gene. *mBio*, 5: e01312-14.
- 498 Hamre D, Procknow JJ. 1966. A new virus isolated from the human respiratory tract. *Proc Soc Exp*  
499 *Biol Med*, 121: 190-193.
- 500 Hsu PD, Lander ES, Zhang F. 2014. Development and applications of CRISPR-Cas9 for genome  
501 engineering. *Cell*, 157: 1262-1278.
- 502 Huang C, Lokugamage KG, Rozovics JM, Narayanan K, Semler BL, Makino S. 2011. SARS  
503 coronavirus nsp1 protein induces template-dependent endonucleolytic cleavage of mRNAs: Viral  
504 mRNAs are resistant to nsp1-induced RNA cleavage. *PLoS Pathog*, 7: e1002433.
- 505 Huynh J, Li S, Yount B, Smith A, Sturges L, Olsen JC, Nagel J, Johnson JB, Gnanihram S, Gates JE,  
506 Frieman MB, Baric RS, Donaldson EF. 2012. Evidence supporting a zoonotic origin of human  
507 coronavirus strain NL63. *J Virol*, 86: 12816-12825.
- 508 Ivashkiv LB, Donlin LT. 2014. Regulation of type I interferon responses. *Nature Rev Immunol*, 14:  
509 36-49.
- 510 Jiang F, Ramanathan A, Miller MT, Tang GQ, Gale M, Patel SS, Marcotrigiano J. 2011. Structural  
511 basis of RNA recognition and activation by innate immune receptor RIG-I. *Nature*, 479: 423-427.
- 512 Kawai T, Akira S. 2010. The role of pattern-recognition receptors in innate immunity: update on Toll-  
513 like receptors. *Nat Immunol*, 11: 373-384.



- 514 Kell A, Stoddard M, Li H, Marcotrigiano J, Shaw GM, Gale M. 2015. Pathogen-associated molecular  
515 pattern recognition of hepatitis C virus transmitted/founder variants by RIG-I is dependent on U-core  
516 length. *J Virol*, 89: 11056-11068.
- 517 Kindler E, Jónsdóttir HR, Muth D, Hamming OJ, Hartmann R, Rodriguez R, Geffers R, Fouchier RAM,  
518 Drosten C, Muller MA, Dijkman R, Thiel V. 2013. Efficient replication of the novel human  
519 betacoronavirus EMC on primary human epithelium highlights its zoonotic potential. *mBio*, 4: e00611-  
520 12.
- 521 Kopecky-Bromberg SA, Martínez-Sobrido L, Frieman M, Baric RA, Palese P. 2007. Severe acute  
522 respiratory syndrome coronavirus open reading frame (ORF) 3b, ORF 6, and nucleocapsid proteins  
523 function as interferon antagonists. *J Virol*, 81: 548-557.
- 524 Kowalinski E, Lunardi T, McCarthy Andrew A, Luber J, Brunel J, Grigorov B, Gerlier D, Cusack S.  
525 2011. Structural basis for the activation of innate immune pattern-recognition receptor RIG-I by viral  
526 RNA. *Cell*, 147: 423-435.
- 527 Lau SKP, Woo PCY, Yip CCY, Tse H, Tsoi H, Cheng VCC, Lee P, Tang BSF, Cheung CHY, Lee RA,  
528 So LY, Lau YL, Chan KH, Yuen KY. 2006. Coronavirus HKU1 and other coronavirus infections in  
529 Hong Kong. *J Clin Microbiol*, 44: 2063–2071.
- 530 Levy DE, Garcia-Sastre A. 2001. The virus battles: IFN induction of the antiviral state and mechanisms  
531 of viral evasion. *Cytokine Growth Factor Rev*, 12: 143-156.
- 532 Li J, Liu Y, Zhang X. 2010. Murine coronavirus induces type I interferon in oligodendrocytes through  
533 recognition by RIG-I and MDA5. *J Virol*, 84: 6472-6482.

- 534 Li W, Moore MJ, Vasilieva N, Sui J, Wong SK, Berne MA, Somasundaran M, Sullivan JL, Luzuriaga  
535 K, Greenough TC, Choe H, Farzan M. 2003. Angiotensin-converting enzyme 2 is a functional receptor  
536 for the SARS coronavirus. *Nature*, 426: 450-454.
- 537 Liu S, Cai X, Wu J, Cong Q, Chen X, Li T, Du F, Ren J, Wu YT, Grishin NV, Chen ZJ. 2015a.  
538 Phosphorylation of innate immune adaptor proteins MAVS, STING, and TRIF induces IRF3 activation.  
539 *Science*, 347: 6227.
- 540 Liu G, Park HS, Pyo HM, Liu Q, Zhou Y. 2015b. Influenza A virus panhandle structure is directly  
541 involved in RIG-I activation and interferon induction. *J Virol*, 89: 6067-6079.
- 542 Lokugamage KG, Narayanan K, Nakagawa K, Terasaki K, Ramirez SI, Tseng CTK, Makino S. 2015.  
543 Middle East respiratory syndrome coronavirus nsp1 inhibits host gene expression by selectively  
544 targeting mRNAs transcribed in the nucleus while sparing mRNAs of cytoplasmic origin. *J Virol*, 89:  
545 10970-10981.
- 546 Loo YM, Gale M. 2011. Immune signaling by RIG-I-like receptors. *Immunity*, 34: 680-692.
- 547 Lu X, Pan J, Tao J, Guo D. 2011. SARS-CoV nucleocapsid protein antagonizes IFN- $\beta$  response by  
548 targeting initial step of IFN- $\beta$  induction pathway, and its C-terminal region is critical for the antagonism.  
549 *Virus Genes*, 42: 37-45.
- 550 Luo D, Ding SC, Vela A, Kohlway A, Lindenbach BD, Pyle AM. 2011. Structural insights into RNA  
551 recognition by RIG-I. *Cell*, 147: 409-422.
- 552 Ma F, Li B, Liu SY, Iyer SS, Yu Y, Wu A, Cheng G. 2015a. Positive feedback regulation of type I IFN  
553 production by the IFN-inducible DNA sensor cGAS. *J Immunol*, 194: 1545-1554.

- 554 Ma F, Li B, Yu Y, Iyer SS, Sun M, Cheng G. 2015b. Positive feedback regulation of type I interferon  
555 by the interferon-stimulated gene STING. *EMBO Rep*, 16: 202-212.
- 556 Matthews KL, Coleman CM, van der Meer Y, Snijder EJ, Frieman MB. 2014. The ORF4b-encoded  
557 accessory proteins of Middle East respiratory syndrome coronavirus and two related bat coronaviruses  
558 localize to the nucleus and inhibit innate immune signalling. *J Gen Virol*, 4:874-882.
- 559 Mazaleuskaya L, Veltrop R, Ikpeze N, Martin-Garcia J, Navas-Martin S. 2012. Protective role of Toll-  
560 like receptor 3-induced type I interferon in murine coronavirus infection of macrophages. *Viruses*, 4:  
561 901-923.
- 562 McIntosh K, Dees JH, Becker WB, Kapikian AZ, Chanock RM. 1967. Recovery in tracheal organ  
563 cultures of novel viruses from patients with respiratory disease. *Proc Natl Acad Sci USA*, 57: 933-940.
- 564 Memish ZA, Mishra N, Olival KJ, Fagbo SF, Kapoor V, Epstein JH, AlHakeem R, Durosinloun A,  
565 Asmari MA, Islam A, Kapoor A, Briese T, Daszak P, Al Rabeeah AA, Lipkin WI. 2013. Middle East  
566 respiratory syndrome coronavirus in bats, Saudi Arabia. *Emerg Infect Dis*, 19: 1819-1823.
- 567 Menachery VD, Debbink K, Baric RS. 2014a. Coronavirus non-structural protein 16: Evasion,  
568 attenuation, and possible treatments. *Virus Res*, 194: 191-199.
- 569 Menachery VD, Einfeld AJ, Schäfer A, Josset L, Sims AC, Proll S, Fan S, Li C, Neumann G, Tilton  
570 SC, Chang J, Gralinski LE, Long CG, Richard WCM, Weiss J, Matzke MM, Webb-Robertson BJ,  
571 Schepmoes AA, Shukla AK, Metz TO, Smith RD, Waters KM, Katze MG, Kawaoka Y, Baric RS.  
572 2014b. Pathogenic influenza viruses and coronaviruses utilize similar and contrasting approaches to  
573 control interferon-stimulated gene responses. *mBio*, 5: e01174-14.

- 574 Menachery VD, Yount, BL, Josset, L, Gralinski LE, Scobey T, Agnihothram S, Katze MG, Baric RS.  
575 2014c. Attenuation and restoration of severe acute respiratory syndrome coronavirus mutant lacking  
576 2'-O-methyltransferase activity. *J Virol*, 88: 4251-4264.
- 577 Mielech AM, Kilianski A, Baez-Santos YM, Mesecar AD, Baker SC. 2014. MERS-CoV papain-like  
578 protease has deISGylating and deubiquitinating activities. *Virology*, 450-451: 64-70.
- 579 Narayanan K, Huang C, Lokugamage K, Kamitani W, Ikegami T, Tseng CTK, Makino S. 2008a.  
580 Severe acute respiratory syndrome coronavirus nsp1 suppresses host gene expression, including that of  
581 type I interferon, in infected cells. *J Virol*, 82: 4471-4479.
- 582 Narayanan K, Huang C, Makino S. 2008b. SARS coronavirus accessory proteins. *Virus Res*, 133: 113-  
583 121.
- 584 Neuman BW, Chamberlain P, Bowden F, Joseph J. 2014. Atlas of coronavirus replicase structure. *Virus*  
585 *Res*, 194: 49-66.
- 586 Niemeyer D, Zillinger T, Muth D, Zielecki F, Horvath G, SulimanT, Barchet W, Weber F, Drosten C,  
587 Müller MA. 2013. Middle East respiratory syndrome coronavirus accessory protein 4a is a type I  
588 interferon antagonist. *J Virol*, 87: 12489-12495.
- 589 Omrani AS, Saad MM, Baig K, Bahloul A, Abdul-Matin M, Alaidaroos AY, Almakhlaifi GA, Albarrak  
590 MM, Memish ZA, Albarrak AM. 2014. Ribavirin and interferon alfa-2a for severe Middle East  
591 respiratory syndrome coronavirus infection: a retrospective cohort study. *Lancet Infect Dis*, 14: 1090-  
592 1095.
- 593 Peisley A, Wu B, Yao H, Walz T, Hur S. 2013. RIG-I forms signaling-competent filaments in an  
594 ATP-dependent, ubiquitin-independent manner. *Mol Cell*, 51: 573-583.

- 595 Pepin KM, Lass S, Pulliam JRC, Read AF, Lloyd-Smith JO. 2010. Identifying genetic markers of  
596 adaptation for surveillance of viral host jumps. *Nat Rev Microbiol*, 8: 802-813.
- 597 Perlman S, Zhao J. 2013. Human coronavirus EMC is not the same as severe acute respiratory  
598 syndrome coronavirus. *mBio*, 4: e00002-13.
- 599 Pyrc K, Berkhout B, van der Hoek L. 2007. The novel human coronaviruses NL63 and HKU1. *J*  
600 *Virology*, 81: 3051–3057.
- 601 Raj VS, Mou H, Smits SL, Dekkers DHW, Muller MA, Dijkman R, Muth D, Demmers JAA, Zaki A,  
602 Fouchier RAM, Thiel V, Drosten C, Rottire PJM, Osterhaus ADME, Bosch BJ, Haagmans BL. 2013.  
603 Dipeptidyl peptidase 4 is a functional receptor for the emerging human coronavirus-EMC. *Nature*, 495:  
604 251-254.
- 605 Rota PA, Oberste MS, Monroe SS, Nix WA, Campagnoli R, Icenogle JP, Peñaranda S, Bankamo B,  
606 Maher K, Chen MH, Ton SX, Tamin A, Lowe L, Frace M, DeRisi JL, Chen Q, Wang D, Erdman DD,  
607 Peret TCT, Burns C, Ksiazek TG, Rollin PE, Sanchez A, Liffick S, Holloway B, Limor J, McCaustland  
608 K, Olsen-Rasmussen M, Fouchier R, Gunther S, Osterhaus ADME, Drosten C, Pallansch MA,  
609 Anderson LJ, Bellini WJ. 2003. Characterization of a novel coronavirus associated with severe acute  
610 respiratory syndrome. *Science*, 300: 1394-1399.
- 611 Samuel CE. 1991. Antiviral actions of interferon interferon-regulated cellular proteins and their  
612 surprisingly selective antiviral activities. *Virology*, 183: 1-11.
- 613 Samuel CE. 2001. Antiviral actions of interferons. *Clin Microbiol Rev*, 14: 778–809.
- 614 Saito T, Owen DM, Jiang F, Marcotrigiano J, Gale M. 2008. Innate immunity induced by composition-  
615 dependent RIG-I recognition of Hepatitis C virus RNA. *Nature*, 454: 523-527.

- 616 Satoh T, Kato H, Kumagai Y, Yoneyama M, Sato S, Matsushita K, Tsujimura T, Fujuta T, Akira S,  
617 Takeuchi O. 2010. LGP2 is a positive regulator of RIG-I- and MDA5-mediated antiviral  
618 responses. *Proc Natl Acad Sci USA*, 107: 1512-1517.
- 619 Schneider WM, Chevillotte MD, Rice CM. 2014. Interferon-stimulated genes: a complex web of host  
620 defenses. *Annu Rev Immunol*, 32: 513–545.
- 621 Schoggins, JW, MacDuff DA, Imanaka N, Gainey MD, Shrestha B, Eitson JL, Mar KB, Richardson  
622 RB, Ratushny AV, Litvak V, Dabelic R, Manicassamy B, Aitchison JD, Aderem A, Elliott RM, García-  
623 Sastre A, Racaniello V, Snijder EJ, Yokoyama WM, Diamond MS, Virgin HW, Rice CM. 2014. Pan-  
624 viral specificity of IFN-induced genes reveals new roles for cGAS in innate immunity. *Nature*, 505:  
625 691-695.
- 626 Scobey T, Yount BL, Sims AC, Donaldson EF, Agnihothram SS, MenacheryVD, Graham RL,  
627 Swanstrom J, Bove PF, Kim JD, Grego S, Randell SH, Baric RS. 2013. Reverse genetics with a full-  
628 length infectious cDNA of the Middle East respiratory syndrome coronavirus. *Proc Natl Acad Sci*  
629 *USA*, 110: 16157-16162.
- 630 Sevajol M, Subissi L, Decroly E, Canard B, Imbert I. 2014. Insights into RNA synthesis, capping, and  
631 proofreading mechanisms of SARS-coronavirus. *Virus Res*, 194: 90-99.
- 632 Shirato K, Kawase M, Matsuyama S. 2013. Middle East respiratory syndrome coronavirus infection  
633 mediated by the transmembrane serine protease TMPRSS2. *J Virol*, 87: 12552-12561.
- 634 Siu KL, Kok KH, Ng MHJ, Poon VKM, Yuen, KY, Zheng BJ, Jin DY. 2009. Severe acute respiratory  
635 syndrome coronavirus m protein inhibits type I interferon production by impeding the formation of  
636 TRAF3·TANK·TBK1/IKK $\epsilon$  complex. *J Biol Chem*, 284: 16202-16209.

- 637 Siu KL, Chan CP, Kok KH, Woo PCY, Jin DY. 2014a. Suppression of innate antiviral response by  
638 severe acute respiratory syndrome coronavirus M protein is mediated through the first transmembrane  
639 domain. *Cell Mol Immunol*, 11: 141-149.
- 640 Siu KL, Chan CP, Kok KH, Woo PC, Jin DY. 2014b. Comparative analysis of the activation of unfolded  
641 protein response by spike proteins of severe acute respiratory syndrome coronavirus and human  
642 coronavirus HKU1. *Cell Biosci*, 4: 3.
- 643 Siu KL, Yeung ML, Kok KH, Yuen KS, Kew C, Lui PY, Chan CP, Tse H, Woo PCY, Yuen KY, Jin  
644 DY. 2014c. Middle East respiratory syndrome coronavirus 4a protein is a double-stranded RNA-  
645 binding protein that suppresses pact-induced activation of RIG-I and MDA5 in the innate antiviral  
646 response. *J Virol*, 88: 4866-4876.
- 647 Takaoka A, Hayakawa S, Yanai H, Stoiber D, Negishi H, Kikuchi H, Sasaki S, Imai K, Shibue T,  
648 Honda K, Taniguchi T. 2003. Integration of interferon- $\alpha/\beta$  signalling to p53 responses in tumour  
649 suppression and antiviral defence. *Nature*, 424: 516-523.
- 650 Tanaka T, Kamitani W, DeDiego ML, Enjuanes L, Matsuura Y. 2012. Severe acute respiratory  
651 syndrome coronavirus nsp1 facilitates efficient propagation in cells through a specific translational  
652 shutoff of host mRNA. *J Virol*, 86: 11128-11137.
- 653 Totura AL, Whitmore A, Agnihothram S, Schäfer A, Katze MG, HeiseMT, Baric RS. 2015. Toll-like  
654 receptor 3 signaling via TRIF contributes to a protective innate immune response to severe acute  
655 respiratory syndrome coronavirus infection. *mBio*, 6: 00638-15.
- 656 Tyrrell DAJ, Bynoe ML. 1965. Cultivation of a novel type of common-cold virus in organ cultures. *Br*  
657 *Med J*, 1: 1467-1470.

- 658 van der Hoek L, Pyrc K, Jebbink MF, Vermeulen-Oost W, Berkhout RJM, Wolthers KC, Wertheim-  
659 van Dillen PME, Kaandorp J, Spaargaren J, Berkhout B. 2004. Identification of a new human  
660 coronavirus. *Nat Med*, 10: 368-373.
- 661 Wang Y, Sun Y, Wu A, Xu S, Pan R, Zeng C, Jin X, Ge X, Shi Z, Ahola T, Chen Y and Guo D. 2015.  
662 Coronavirus nsp10/nsp16 methyltransferase can be targeted by nsp10-derived peptide *in vitro* and *in*  
663 *vivo* to reduce replication and pathogenesis. *J Virol*, 89: 8416-8427.
- 664 Weber M, Gawanbacht A, Habjan M, Rang A, Borner C, Schmidt AM, Veitinger S, Jacob R, Devignot  
665 S, Kochs G, Weber F. 2013. Incoming RNA virus nucleocapsids containing a 5'-triphosphorylated  
666 genome activate RIG-I and antiviral signaling. *Cell Host Microbe*, 13: 336-346.
- 667 Woo PCY, La, SKP, Chu C, Chan K, Tsoi H, Huang Y, Wong BHK, Poon RWS, Cai JJ, Luk WK,  
668 Poon LLM, Wong SSY, Guan Y, Peiris JSM, Yuen KY. 2005. Characterization and complete genome  
669 sequence of a novel coronavirus, coronavirus HKU1, from patients with pneumonia. *J Virol*, 79: 884-  
670 895.
- 671 Woo PCY, Lau SKP, Huang Y, Yuen KY. 2009. Coronavirus diversity, phylogeny and interspecies  
672 jumping. *Exp Biol Med (Maywood)*, 234: 1117-1127.
- 673 Woo PCY, Lau SKP, Lam CSF, Lau CCY, Tsang AKL, Lau JHN, Bai R, Teng JLL, Tsang CCC, Wang  
674 M, Zheng BJ, Chan KH, Yuen KY. 2012. Discovery of seven novel mammalian and avian  
675 coronaviruses in the genus deltacoronavirus supports bat coronaviruses as the gene source of  
676 alphacoronavirus and betacoronavirus and avian coronaviruses as the gene source of  
677 gammacoronavirus and deltacoronavirus. *J Virol*, 86: 3995-4008.
- 678 Wu B, Peisley A, Richards C, Yao H, Zeng X, Lin C, Chu F, Walz T, Hur S. 2013. Structural basis for  
679 dsRNA recognition, filament formation, and antiviral signal activation by MDA5. *Cell*, 152: 276-289.



- 680 Wu B, Peisley A, Tetrault D, Li Z, Egelman EH, Magor KE, Walz T, Penczek PA, Hur S. 2014.  
681 Molecular imprinting as a signal-activation mechanism of the viral RNA sensor RIG-I. *Mol Cell*, 55:  
682 511-523.
- 683 Xagorari A, Chlichlia K. 2008. Toll-like receptors and viruses: induction of innate antiviral immune  
684 responses. *Open Microbiol J*, 2: 49-59.
- 685 Yang Y, Zhang L, Geng H, Deng Y, Huang B, Guo Y, Zhao Z, Tan W. 2013. The structural and  
686 accessory proteins M, ORF 4a, ORF 4b, and ORF 5 of Middle East respiratory syndrome coronavirus  
687 (MERS-CoV) are potent interferon antagonists. *Protein Cell*, 4: 951-961.
- 688 Yang Y, Ye F, Zhu N, Wang W, Deng Y, Zhao Z, Tan W. 2015. Middle East respiratory syndrome  
689 coronavirus ORF4b protein inhibits type I interferon production through both cytoplasmic and nuclear  
690 targets. *Sci Rep*, 5: 17554.
- 691 Yeager CL, Ashmun RA, Williams RK, Cardellichio CB, Shapiro LH, Look AT, Holmes KV. 1992.  
692 Human aminopeptidase N is a receptor for human coronavirus 229E. *Nature*, 357: 420-422.
- 693 Yoneyama M, Kikuchi M, Natsukawa T, Shinobu N, Imaizumi T, Miyagishi M, Taira K, Akira S,  
694 Fujita T. 2004. The RNA helicase RIG-I has an essential function in double-stranded RNA-induced  
695 innate antiviral responses. *Nat Immunol*, 5: 730-737.
- 696 Yoneyama M, Kikuchi M, Matsumoto K, Imaizumi T, Miyagishi M, Taira K, Foy E, Loo YM, Gale M  
697 Jr, Akira S, Yonehara S, Kato A, Fujita T. 2005. Shared and unique functions of the DExD/H-box  
698 helicases RIG-I, MDA5, and LGP2 in antiviral innate immunity. *J Immunol*, 175: 2851-2858.
- 699 Yount B, Curtis KM, Fritz EA, Hensley LE, Jahrling PB, Prentice E, Denison MR, Geisbert TW, Baric  
700 RS. 2003. Reverse genetics with a full-length infectious cDNA of severe acute respiratory syndrome  
701 coronavirus. *Proc Natl Acad Sci USA*, 100: 12995-13000.

- 702 Yuen KS, Chan CP, Wong NHM, Ho CH, Ho TH, Lei T, Deng W, Tsao SW, Chen H, Kok KH, Jin  
703 DY. 2015. CRISPR/Cas9-mediated genome editing of Epstein-Barr virus in human cells. *J Gen Virol*,  
704 96: 626-636.
- 705 Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus AD, Fouchier RA. 2012. Isolation of a novel  
706 coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med*, 367: 1814-1820.
- 707 Zhang R, Jha BK, Ogden KM, Dong B, Zhao L, Elliott R, Patton JT, Silverman RH, Weiss SR. 2013.  
708 Homologous 2', 5'-phosphodiesterases from disparate RNA viruses antagonize antiviral innate  
709 immunity. *Proc Natl Acad Sci USA*, 110: 13114-13119.
- 710 Zhao L, Jha BK, Wu A, Elliott R, Ziebuhr J, Gorbalenya AE, Silverman RH, Weiss SR. 2012.  
711 Antagonism of the interferon-induced OAS-RNase L pathway by murine coronavirus ns2 protein is  
712 required for virus replication and liver pathology. *Cell Host Microbe*, 11: 607-616.
- 713 Zhong Y, Tan YW, Liu DX. 2012. Recent progress in studies of arterivirus- and coronavirus-host  
714 interactions. *Viruses*, 4: 980-1010.
- 715 Zornetzer GA, Frieman MB, Rosenzweig E, Korth MJ, Page C, Baric RS, Katze MG. 2010.  
716 Transcriptomic analysis reveals a mechanism for a profibrotic phenotype in STAT1 knockout mice  
717 during severe acute respiratory syndrome coronavirus infection. *J Virol*, 84: 11297-11309.
- 718 Züst R, Cervantes-Barragan L, Habjan M, Maier R, Neuman BW, Ziebuhr J, Szretter KJ, Baker SC,  
719 Barchet W, Diamond MS, Siddell SG, Ludewig B, Thiel V. 2011. Ribose 2'-O-methylation provides a  
720 molecular signature for the distinction of self and non-self mRNA dependent on the RNA sensor  
721 Mda5. *Nat Immunol*, 12: 137-143.
- 722

728 Figure 1. Innate immune response mediated against coronavirus infection and viral evasion mechanisms.

729 (Left) Upon CoV infection, viral genome ssRNA as well as dsRNA intermediate found in virus life cycle  
730 are exposed to host innate immune sensors, RIG-I/MDA5 in cytoplasm or Toll-like receptors TLR3/7/8 in  
731 endosome. Activation of these immune sensors initiates a downstream signaling cascade that leads to IFN-  
732  $\beta$  gene expression. RIG-I/MDA5 conveys signal through a mitochondrial adaptor MAVS while TLR signals  
733 through TRIF/MyD88. Both pathways share the common TRAF adaptor to activate transcription factors.  
734 TRAF3 serves as an adaptor which activates TANK·TBK1/IKK $\epsilon$  complex for IRF3 phosphorylation and  
735 subsequent dimerization, while TRAF6 is responsible for the activation of IKK complex which  
736 phosphorylates the canonical inhibitor of NF- $\kappa$ B (I $\kappa$ B). Activated transcription factors are translocated into  
737 the nucleus to drive IFN- $\beta$  expression. (Right) IFN- $\beta$  are secreted into extracellular space and bound to its  
738 cognate receptors IFNAR to activate downstream JAK-STAT signaling. Receptor-associated tyrosine  
739 kinases Jak1 and Tyk2 are brought to juxtaposition for self-phosphorylation and activation. STATs are  
740 recruited to and phosphorylated by the tyrosine kinases. Phosphorylated STAT1/2 with IRF9 forms a  
741 ternary complex ISGF3 which translocates into the nucleus and binds to ISRE in the promoter region  
742 upstream of ISG genes. ISG genes are expressed consequently to establish an antiviral state in cells. OAS  
743 is an example of ISG which produces 2', 5'-oligoadenylate (2', 5'-A) upon detection of dsRNA and activates  
744 RNase L to cleave viral RNA to yield more RLR ligand as a positive-feedback mechanism of IFN  
745 production. The CoV-encoded proteins shown in red are known to intervene the host innate immune  
746 signaling at various action points as evasion mechanisms to sustain viral replication and propagation. The  
747 action points at which viral proteins function marked with a question mark (?) represent controversial and  
748 inconclusive findings in the field or molecular mechanisms not well studied. MHV: mouse hepatitis virus.

