

Running head: Aging and Memory

A multi-method comparison of autobiographical memory impairments amongst younger and older adults

Tom J. Barry^{1,2}, James D. Gregory³, Jose M. Latorre^{4,5}, Laura Ros^{4,5}, Marta Nieto^{4,5}, Jorge J. Ricarte^{4,5}

¹Department of Psychology, The University of Hong Kong, Hong Kong, Hong Kong.

²Department of Psychology, The Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, United Kingdom.

³Department of Psychology, University of Bath, United Kingdom

⁴Department of Psychology, University of Castilla La Mancha, Albacete, Spain

⁵Applied Cognitive Research Unit, School of Medicine, Albacete, Spain

Acknowledgements: This study has been supported by the Regional Government of Castilla La Mancha [Consejería de Educación y Ciencia de Castilla La Mancha, grant PIII109-0274-8863], the Ministry of Science and Innovation [Ministerio de Ciencia e Innovación, grant PSI2010-20088] and Ministry of Economy and Competitiveness [Ministerio de Economía y Competitividad, grant DPI2016-80894-R].

Disclosures: The authors have no conflicts of interest to disclose.

Corresponding author:

Jorge J. Ricarte; Department Psychology UCLM, School of Medicine Albacete, Avda Almansa, 14, 02006 Albacete (Spain); jorgejavier.ricarte@uclm.es

Abstract

Objectives

Research indicates that, compared to younger adults, older adults have difficulty recalling memories of specific past events (those lasting less than 24 hours) and this difficulty is associated with depression. These studies are largely confined to a single measure of specific memory recall and there are conflicting findings when alternative measures are used. This investigation provides the first comparison of memory specificity between younger and older adults using several different measures.

Method

Older ($n = 105$) and younger ($n = 88$) adults completed the Autobiographical Memory Test (AMT), Autobiographical Memory Interview (AMI) and Sentence Completion for Events from the Past Test (SCEPT) and the number of specific memories was quantified for each measure. Participants also completed the Beck Depression Inventory Version II (BDI-II).

Results

Compared to younger adults, older adults recalled fewer specific memories in the AMT and more specific memories in the AMI. This latter effect was particularly pronounced for memories related to childhood. There was no group difference in responses in the SCEPT. There was no evidence of an association between memory specificity and depression for any of the measures.

Conclusion

Older adults have difficulty retrieving specific memories after cuing by nouns and adjectives, as in the AMT, but they have enhanced recall of specific memories after cuing by life periods, as in the AMI, and this is particularly true of memories related to childhood. Individual differences in memory specificity are not related to depression symptoms in healthy samples.

Keywords: Depression; Episodic Memory; Specificity; Overgeneral; Aging

Introduction

Autobiographical memory is a sub-system of episodic memory that pertains to personally experienced past events that are typically retrieved and constructed in response to environmental cues (e.g., a song we used to dance to) or due to conscious retrieval (Conway & Pleydell-Pearce, 2000). Compared to younger adults, older adults (aged over 50 years) have difficulty recalling memories of specific autobiographical events (Wilson & Gregory, 2018). In particular, older adults have difficulty recalling events from their past that lasted for less than 24 hours (e.g., when I attended my daughters' 20th birthday party) and instead tend to retrieve *general* memories: memories for events that occurred multiple times (e.g., my daughters' birthday parties) or which occurred over extended periods of time (e.g., when my daughter was travelling during her twenties). This is important because the ability to retrieve specific memories has been suggested to help older adults regulate negative moods (Carretero, Latorre, Fernández, Barry, & Ricarte, 2019) and difficulty retrieving specific memories has been associated with the presence and severity of depression (Farina, Barry, van Damme, van Hie, & Raes, 2019; Ono, Devilly, & Shum, 2015; Van Vreeswijk & De Wilde, 2004). Although autobiographical memory impairments and their association with depression in older adults has been widely researched (Wilson & Gregory, 2018), there are limitations to existing studies that must be overcome in order for us to better understand these problems and to inform the development of interventions to improve them (e.g., Leahy, Ridout, Mushtaq, & Holland, 2017).

The majority of studies in this area are confined to a single assessment method, the Autobiographical Memory Test (AMT; Williams & Broadbent, 1986). In a typical AMT, participants are presented with five negative and five positive cue words (e.g., sad, happy) and they are asked to retrieve a memory associated with each within 60 seconds. In studies with the AMT, older adults have shown problems recalling specific memories overall as well

as particular difficulty recalling specific memories cued by negative words (Latorre et al., 2013; Ros & Latorre, 2010; Serrano, Latorre, & Gatz, 2007). AMT research has also found that older adults with elevated depression symptoms (Serrano et al., 2007) or diagnoses of depression (Birch & Davidson, 2007; Ricarte et al., 2011) appear to have marked difficulty recalling specific memories compared to their low depression/diagnoses-free counterparts. However, there is little evidence that individual differences in depression symptoms are correlated with memory specificity amongst older adults (Serrano et al., 2007). In one study that used an adapted AMT that cued memories from different life-periods, participants with elevated depression symptoms showed difficulty recalling specific memories of events from childhood and adolescence compared to participants with lower depression levels (Gidron & Alon, 2007). It might be that the association between memory specificity and depression symptoms amongst older adults is most apparent when alternative cuing techniques to those used in the traditional AMT are used, and in particular those which probe early life periods. Depression is particularly prevalent amongst older adults over the age of 65 years who have a one in four chance of experiencing depression at any one time (Craig & Mindell, 2005). There is some evidence that the transition into retirement that typically occurs at this time can be a particularly high-risk time for the emergence of depression for some (Reitzes, Mutran, & Fernandez, 1996), but not all (Lindwall et al., 2017), older adults. Elucidating the factors that predict the presence and severity of depression symptoms amongst older adults is therefore critical.

Although the AMT has been invaluable in helping researchers elucidate the nature of reduced autobiographical memory specificity and its association with depression, it is merely one method for assessing memory specificity and the task itself has little relevance to the way that memories are cued or recalled in real life. In real life, memories are rarely cued using discrete nouns or adjectives that are free of any context and memories are typically not

demanded within a time limit. In addition, with one exception (Gidron & Alon, 2007), the AMT does not typically account for differences in specificity that might occur for memories from different points across the lifespan or how this might differ between younger and older adults.

One alternative to the AMT which accounts for this, the Autobiographical Memory Interview (AMI; Kopelman, Wilson, & Baddeley, 1990), instead cues memories related to times (childhood, adulthood and recently) and locations that are commonly experienced within a person's lifetime (e.g., when you were in primary school; when you were in, or visited, hospital) and participants are not given any time limit. To our knowledge, only two studies have used the AMI amongst older adults (De Medeiros, Mosby, Hanley, Pedraza, & Brandt, 2011; Gallassi, Di Sarro, Morreale, & Amore, 2006). Both of these studies involve interventions and have not examined group differences in memory specificity between older adults and any other control group (e.g., younger adults).

Another alternative to the AMT is the Sentence Completion for Events from the Past Test (SCEPT; Raes, Hermans, Williams, & Eelen, 2007). In the SCEPT participants are shown a series of sentence stems (e.g., I'll never forget when...) and they are asked to retrieve a memory that completes the sentence. The SCEPT was initially created because the AMT was seen as not sensitive enough to capture the problems with memory specificity that are present in non-clinical participants; and, that there was so little variation in specificity between such participants when it was measured using the AMT, that it appeared to be unrelated to other aspects of psychopathology (e.g., depression and rumination)(Raes et al., 2007). The SCEPT was thought to be more sensitive than the AMT given its focus on cuing memories with sentence stems that are more closely aligned with the ways in which memories are retrieved and discussed in regular discourse. Indeed, Raes et al. (2007) found that the SCEPT was more sensitive than the AMT in terms of its ability to measure the

association between specificity and depression in non-clinical samples. However, to our knowledge, no study has used the SCEPT to assess memory specificity amongst older adults.

In addition, to our knowledge no study has used multiple methods for assessing autobiographical memory specificity within a single investigation. It is unclear whether problems with memory specificity evident amongst older adults are consistent across cuing methods or whether they are confined to a particular method. Also, many previous studies are comparisons amongst older adults (e.g., Serrano et al., 2007) and so do not involve comparisons with younger groups of adults.

The present investigation overcomes these limitations by examining differences in autobiographical memory specificity between older adults (aged approximately between 50 and 80 years) and younger adults (aged approximately between 20 and 40 years) measured with three cuing procedures: the AMT, AMI and SCEPT, whilst also examining the contribution of individual differences in depression symptoms to memory recall. We also examined the extent to which each of the measures quantified the same underlying construct of autobiographical memory specificity and so whether performance in each tasks was related to one another.

Methods

Participants

196 participants were recruited to participate from different University academic programs of the region associated with the senior author where not only young but also older adults participate (e.g. summer schools, conferences, seminars, workshops, specific university program for those over 50 years old). Participation was voluntary and non-remunerated. Those participants with a possible psychiatric diagnoses according to screening with the Mini-International Neuropsychiatric Interview (Sheehan et al., 1998) were excluded from participation. In addition, in the group of older participants, the Mini-Mental Status

Examination (MMSE; Lobo et al., 1979) was used so that participants with cognitive impairment could be excluded. One participant did not complete any items of the measure of depression symptoms and two participants did not complete one of the autobiographical memory tasks. These participants were excluded from analyses. Data from 193 participants' (Older adults: $n = 105$, $\text{Range}_{age} = 53-80$, $M_{age} = 64.85$, $SD_{age} = 5.27$, $Females = 68$ (65%); Younger adults: $n = 88$, $\text{Range}_{age} = 19-36$, $M_{age} = 22.28$, $SD_{age} = 4.12$, $Females = 62$ (70%)) were analysed. Of these participants, one had six missing items on the measure of depression symptoms and seven had one missing item. These data were imputed using a person-level mean based on responses to the remaining items.

Measures

Beck Depression Inventory, second version (BDI-II)

The BDI-II is a self-report measure of depression severity (Beck, Steer, & Brown, 1996). Participants are presented with 21 items that refer to a range of depression symptoms and are asked to rate, on scales from 0 to 3, the intensity and frequency with which they experience each of these symptoms. A high score, therefore, reflects higher depression severity. In the present investigation, the BDI-II showed acceptable internal consistency ($\alpha = .75$).

Autobiographical memory

Sentence Completion for Events from the Past Test (SCEPT). The SCEPT (Raes et al., 2007) records autobiographical memories cued by 11 short uncompleted sentences (e.g., Why I think back to...), wherein participants complete the stem of the sentence with a memory. There was no time limit for participants' responses. Sentences describing unique events lasting less than 24 hours were categorized as specific memories (e.g. ...my wedding day); sentences describing repeated summaries of events were categorized as categoric memories (e.g...every time I saw my wife); sentences describing events lasting longer than 24 hours or periods of life were categorized as extended memories (e.g. ...when I was a child").

Sentences completed with the name of persons, animals, or things, repetitions, expressions of feelings not associated with an event or a period of time, as well as the omission of a response, were categorized as non-responses. Two independent researchers categorized responses. Inter-rater agreement calculated on 15% of the total obtained sentences was 92%.

Autobiographical Memory Test (AMT). The AMT comprised five positive (Happiness, Friendship, Energy, Smile, Excited) and five negative (e.g., Guilty, Failure, Delusion, Worry, Sadness, Illness) cue words presented orally in a fixed order, alternating between positive and negative cue words (Williams, Barnhofer, Crane, & Beck, 2005). Participants were asked to generate a specific memory that happened on a particular day one week ago or more.

Participants were told that they were not allowed to repeat memories. After 60-seconds the next cue word was presented. Responses were audio-recorded and transcribed. The same coding framework as was used for SCEPT responses was employed for AMT responses. A subsample of 15% of recalled memories were coded by two different examiners. The percentage of agreement in the codes given by raters was 85%.

Autobiographical Memory Interview (AMI). Although the AMI measure includes items that assess both episodic and semantic aspects of autobiographical memory, as the focus of the present investigation was on the former, only these nine items were included. This section of the AMI asks participants to retrieve a memory related to each of several times and locations related to three periods of their lives: childhood (e.g., a memory from primary school), adulthood (e.g., a memory from college or first job) and recent years (e.g., a memory from a hospital within the past five years). Items were administered orally by the experimenter without time restriction. When no answer was obtained or the answer was not related to the questions, participants were prompted once after which the next cue was given. Memories were coded using the same framework as the SCEPT and AMT and also 15% of responses were coded by two independent examiners with 86% agreement between raters.

Procedure

Study delivery and data handling procedures followed legal standards set out by the Clinical Research Ethics Committee of the local University Hospital after the study protocol was approved (reference: CEIC06/2016). After participants arrived at the laboratory, they were given a summary of information regarding the study and its aims and were informed that they could drop out of the study at any moment. Researchers confirmed that participants had understood the instructions and then informed consent was obtained. Tests were collected by seven graduate students trained in the delivery of the tests. The BDI was administered first, then the SCEPT and, in the second session a week later, the AMI was completed and then the AMT.

Statistical procedure

First, we clarified whether and to what extent the groups differed in terms of their average age, their depression symptoms and their proportions of males and females and educational levels (coded as 0 and 1 where 1 was university-level education). Where there was evidence that the groups differed in terms of their gender distribution or education levels, these variables were included in subsequent analyses.

Then, in accordance with recommendations regarding the benefits of robust statistical methods (Field & Wilcox, 2017), the present investigation tested for group differences between younger and older adults using robust linear models. Given that different measures had different numbers of cues, different models were tested for each of the task measures: SCEPT, AMT and AMI.

As the SCEPT has only a single cue type and so is scored using a single total score, two robust linear models (using the `lmrob()` function in the *robustbase* package in R (Maechler et al., 2019)) tested for group differences in the recall of specific autobiographical memories in the SCEPT. The first model examined age-group differences in recall whereas

the second model further examined the contribution of group differences in depression symptoms to any age-group differences. Main effects and interactions were evidenced by p values less than .05.

Two robust linear mixed models with maximum likelihood estimation (using the `rlmer()` function in the *robustlmm* package in R (Koller, 2016)) tested for group differences in the recall of specific memories following positive and negative cue words in the AMT. These models included a random effect for between-participant variation and included fixed effects for cue valence (positive vs. negative) and group, and the interaction between these effects. As in the SCEPT analysis, the second model examined the additional contribution of depression symptoms, and group differences in depression, to recall within the AMT. The `rlmer()` function does not return p values. As such, in this analysis and other analyses using this function, a main effect and interaction is evidenced by 95% CI that does not overlap with zero.

Two final robust linear mixed models (again using `rlmer()`) examined group differences in recall of specific memories in the AMI. These models were identical to those within the AMT analyses with the exception that the model included a fixed effect of time rather than cue valence. As time had three levels (childhood, adulthood, recently) this variable was modelled as dummy variables. Again, the second model examined the additional contribution of depression symptoms.

Where appropriate, main effects and interactions were followed up with comparisons of model estimated means using a Bonferroni correction.

Finally, total scores for the number of specific memories retrieved in each of the tasks were correlated with one another.

The analysis script and data are available on Open Science Framework (<https://osf.io/ky3c2/>).

Results

Participant characteristics

Younger adults were significantly younger than the older adults, $t(191) = -61.65$, $p < .001$, 95% CI[-43.93, -41.20], $d = 2.91$. There was no difference between the groups in their proportions of female participants, $X^2(1) = 0.706$, $p = 0.401$, $d = 0.12$, 95% CI[-0.16, 0.40].

Younger adults had a significantly larger proportion of participants that had obtained University-level education (98%) than older adults (41%), $X^2(1) = 69.63$, $p < .001$, $d = 1.50$, 95% CI[1.15, 1.86]. The younger adults also reported significantly lower depression symptoms ($M = 4.50$, $SD = 3.59$) compared to older adults ($M = 6.64$, $SD = 5.28$), $t(191) = -3.22$, $p = .001$, 95% CI[-3.45, -0.83], $d = 1.53$.

Group differences in autobiographical memory

See Table 1 for the summary statistics for the SCEPT, AMI and AMT.

Sentence Completion for Events from the Past Test (SCEPT)

In the first model, age group did not show a significant effect on recall within the SCEPT, $\beta = -0.299$, $SE = 0.388$, $p = .443$, 95% CI[-1.07, 0.47]. In the second model including BDI-II scores and its interaction with group, age group did not show a significant effect on recall, $\beta = -0.200$, $SE = 0.374$, $p = .593$, 95% CI[-0.94, 0.54]. Also, neither BDI-II scores, $\beta = -0.051$, $SE = 0.062$, $p = .408$, 95% CI[-0.17, 0.07], nor the interaction between BDI-II scores and age group, $\beta = 0.010$, $SE = 0.077$, $p = .901$, 95% CI[-0.14, 0.16], showed significant effects on recall. Education was not associated with recall in either the first model, $\beta = -0.501$, $SE = 0.437$, $p = .252$, 95% CI[-1.36, 0.36], or the second, $\beta = -0.492$, $SE = 0.433$, $p = .257$, 95% CI[-1.35, 0.36],

Autobiographical Memory Test (AMT)

In the first model, there was a main effect of group, $\beta = -1.032$, $SE = 0.200$, 95%CI[-1.42, -0.64]. The direction of this effect was such that younger adults recalled significantly more

specific memories than older adults. There was no main effect of valence, $\beta = -0.087$, $SE = 0.112$, 95% CI[-0.31, 0.13], or interaction between valence and group, $\beta = 0.004$, $SE = 0.152$, 95% CI[-0.29, 0.30]. Education was also not associated with recall, $\beta = 0.209$, $SE = 0.196$, 95% CI[-0.17, 0.59]

In the second model, there was a main effect of group, $\beta = -1.042$, $SE = 0.207$, 95%CI[-1.45, -0.64]. There was no main effect of valence, $\beta = -0.078$, $SE = 0.118$, 95% CI[-0.31, 0.15], or interaction between valence and group, $\beta = 0.006$, $SE = 0.158$, 95% CI[-0.30, 0.32]. Again, there was no main effects of BDI-II scores, $\beta = -0.002$, $SE = 0.034$, 95% CI[-0.07, 0.07], or interaction between BDI-II scores and group, $\beta = 0.017$, $SE = 0.041$, 95% CI[-0.06, 0.10], BDI-II scores and valence, $\beta = 0.010$, $SE = 0.032$, 95% CI[-0.05, 0.07], or a three-way interaction between BDI-II scores, group and cue valence, $\beta = -0.030$, $SE = 0.037$, 95% CI[-0.10, 0.04]. Education was not associated with recall, $\beta = 0.207$, $SE = 0.198$, 95% CI[-0.18, 0.59].

Autobiographical Memory Interview (AMI)

In the first model, there was a main effect of group, $\beta = 0.448$, $SE = 0.104$, 95% CI[0.24, 0.65]. However, contrary to the AMT analysis, the direction of this effect was such that, across cue types, younger adults recalled significantly fewer specific memories compared to older adults. There was also main effect for the difference in specificity between memories recalled from childhood and those recalled from adulthood, $\beta = 0.650$, $SE = 0.099$, 95% CI[0.45, 0.84], and this variable interacted with group, $\beta = -0.532$, $SE = 0.135$, 95% CI[-0.80, -0.27]. There was also a main effect for the difference in specificity between memories recalled from childhood and those recalled from recent years, $\beta = -0.313$, $SE = 0.099$, 95% CI[-0.51, -0.12], and this variable interacted significantly with group, $\beta = -0.371$, $SE = 0.135$, 95% CI[-0.63, -0.11]. There was also a main effect of education, $\beta = 0.159$, $SE = 0.073$, 95% CI[0.02, 0.30].

In the second model, examining the contribution of depression symptoms to recall, again there was a main effect of group, $\beta = 0.463$, $SE = 0.107$, 95% CI[0.25, 0.67]. The main effect for difference in recall from childhood to adulthood continued to be significant, $\beta = 0.675$, $SE = 0.105$, 95% CI[0.47, 0.88], and this effect interacted with group, $\beta = -0.575$, $SE = 0.140$, 95% CI[-0.85, -0.30]. Also, the main effect for difference in recall from childhood to recently continued to be significant, $\beta = -0.328$, $SE = 0.105$, 95% CI[-0.53, -0.12], and this effect interacted with group, $\beta = -0.342$, $SE = 0.140$, 95% CI[-0.62, -0.06]. There was also a main effect of education, $\beta = 0.162$, $SE = 0.072$, 95% CI[0.02, 0.30]. All of the main or interacting effects of depression symptoms showed confidence intervals that overlapped with zero.

After correcting for multiple comparisons with a Bonferroni correction, contrary to the AMT analyses, younger adults recalled significantly fewer specific memories from childhood than older adults, $b = -0.463$, $SE = 0.107$, $t(489) = -4.327$, $p < .001$, 95% CI[-0.78, -0.15]. There was no group difference for memories recalled from adulthood, $b = 0.112$, $SE = 0.107$, $t(489) = 1.045$, $p = 1.000$, 95% CI[-0.20, 0.43] or recently, $b = -0.120$, $SE = 0.107$, $t(489) = -1.122$, $p = 1.000$, 95% CI[-0.44, 0.20]. This change from a significant between-group effect in childhood to non-significant effects in adulthood and recently, was because younger adults recalled significantly more specific memories from adulthood than childhood, $b = -0.675$, $SE = 0.107$, $t(342)$, $p < .001$, 95% CI[-0.98, -0.37], whereas older adults did not, $b = -0.100$, $SE = 0.093$, $t(342) = -1.079$, $p = 1.000$, 95% CI[-0.37, 0.17]. In addition, both older adults, $b = 0.770$, $SE = 0.093$, $t(342) = 8.323$, $p < .001$, 95% CI[0.50, 1.04], and younger adults, $b = 1.002$, $SE = 0.105$, $t(342) = 9.586$, $p < .001$, 95% CI[0.69, 1.31], recalled significantly more specific memories from adulthood than recently.

Older adults recalled significantly more specific memories from childhood than younger adults did and both groups were able to recall significantly more specific memories

from adulthood than recently, with no difference between the groups for memories recalled from these timepoints.

Correlations between measures

The total number of specific memories retrieved in the AMT did not correlate with scores for the AMI, $r = -.100$, $p = .167$, 95% CI[-0.24, 0.04], or the SCEPT, $r = -.042$, $p = .558$, 95% CI[-0.18, 0.10]. However, AMI scores correlated significantly with SCEPT scores, $r = .327$, $p < .001$, 95% CI[0.20, 0.45].

Discussion

The present investigation provides an analysis of autobiographical memory specificity amongst older adults, relative to younger adults, by comparing performance in three measures of autobiographical memory. Compared to younger adults, older adults recalled significantly fewer specific memories cued by both positive and negative cue words in the AMT. However, when comparing performance between these groups in the AMI, relative to younger adults, older adults recalled significantly *more* specific memories. This group difference was also particularly pronounced for memories that were recalled from events in childhood. No group difference was evident in the third measure of autobiographical memory, the SCEPT.

That older adults had difficulty, compared to younger adults, recalling specific memories related to both positive and negative cues is consistent with the findings of a previous literature review (Wilson & Gregory, 2018). However, our analysis of the AMI did yield differences between cue types. In particular, older adults were better able than younger adults to recall specific memories related to childhood whereas both groups did not differ in the number of specific memories they retrieved for other time periods. Given that adults also showed a significant improvement in specificity for memories from adulthood relative to childhood, and that older adults showed no such improvement, the most likely explanation

for the group difference that was observed for memories cued from childhood is that it was due to worse recall amongst younger adults rather than an advantage amongst older adults. Indeed, previous results have shown that older adults remember more relevant life events from the end of childhood and beginning of adulthood than their younger counterparts because for younger people the most significant events in life are those things which are happening currently (e.g. marriage, stable work)(Rubin & Schulkind, 1997). Future investigations could examine this suggestion by asking participants to rate the importance of the recalled events within their lives. Future studies should also attempt to quantify the time that has elapsed since the events in the retrieved memory occurred as this may also influence the retrievability of these memories. It is of note that, one previous study using the AMI did not find significant differences between older and young participants in the total number of specific memories that were retrieved, although no information about differences by period was reported (Rathbone, Holmes, Murphy, & Ellis, 2015).

Given that no group difference was observed for two of the three cue types used in the AMI, it is perhaps not surprising that no group difference was observed in our third measure of memory specificity, the SCEPT. Nevertheless, the opposing findings evident in our AMT and AMI analyses are notable and in need of further research. It would be interesting for future research to codify the life periods of specific memories retrieved in the AMT. It may be that older adults who are more likely to retrieve memories from childhood are less likely to experience problems with memory specificity as measured using the AMT, than those who rely on more recent memories. In addition, research could explore the interaction between cue valence and the time period in a hybrid AMT/AMI task by asking participants for positive and negative memories from each of the three time periods referred to in the AMI (Gidron & Alon, 2007). Younger and older adults may differ in their abilities to retrieve positive and negative specific memories for each of the time points and this may contribute to

the broader group differences that were observed in the present investigation in the AMT and AMI measures. Indeed, there is evidence that older adults recall autobiographical memories with greater vividness and detail than do younger adults and that this effect may be moderated by memory valence (Comblain, D'Argembeau, & Van Der Linden, 2005; St. Jacques & Levine, 2007). This broad effect of memory vividness and detail might explain why older adults outperformed younger adults in the AMI. In addition to the suggestions regarding memory importance, future investigations could additionally assess other phenomenological properties such as vividness and detail to explore to what extent these qualities differ between groups and cue-types.

This is the first investigation to employ several different methods for measuring memory specificity within a single study amongst older adults. As all three measures employed here measure the same overarching construct, one might have expected performance in each of the measures to correlate with one another but this was not the case. There are important differences between the tasks that might explain these differences. In particular, the AMT cues memories using positive and negative nouns and adjectives; the AMI uses cues related to locations and time-periods from a person's past (e.g., an occasion when they were in hospital; or, an occasion from primary school); and, the SCEPT uses sentence stems from statements that typically precede instances of recollection (e.g., I'll never forget when...) from which participants must complete the sentence with a memory. The findings presented here suggest that older adults have particular difficulty retrieving specific memories when these memories are cued by valenced nouns and adjectives and when these cues are free from information about physical (e.g., a memory from when you were in hospital) or temporal (e.g., a memory from childhood) context as in the case of the AMI.

It is of note that specificity measured using the AMI and SCEPT correlated positively. It could be that these measures share similar cuing procedures and so they quantify similar

aspects of autobiographical memory specificity. Indeed, both measures ask for important, often life-defining memories (e.g., SCEPT: *I'll never forget when...*; AMI: *A memory from your first job.*) that are perhaps more accessible to older adults than the memories cued by the AMT. It may be that the AMT provides the most sensitive measure of specificity problems given that the cues that are used in this task are free from contextual information that might otherwise aid retrieval as in the AMI or SCEPT. However, it must also be noted that the tasks also differ in terms of whether a time limit was applied or not. Problems retrieving specific memories were seen amongst older adults only when a time limit was applied (AMT, 60 seconds) and in measures without a time limit, older adults either outperformed younger adults (AMI) or showed no difference from them (SCEPT). It may be that in the presence of a time limit and the absence of additional contextual information in the AMT cues, compared to the cues in the AMI or SCEPT, participants have to engage in lengthy and cognitive demanding generative retrieval that many participants fail to complete within the time limit. However, if the absence of a time limit caused the improvement in performance for older adult participants that was observed in the AMI then we would expect that the same would also be true for the SCEPT and also for all cue types within the AMI and not just for childhood. It could also be that time limits impose additional anxiety on participants as they struggle to meet the demands of the task given to them and that this anxiety influences their ability to retrieve memories (Tollenaar, Elzinga, Spinhoven, & Everaerd, 2009). Future research could quantify the amount of time it takes younger and older adults to retrieve specific memories in all tasks or use time limits of varying length and explore the effects of this on between-group differences. Both of these options would enable researchers to examine whether younger and older adults differ in the speed with which they retrieve specific memories and if, given more time in the AMT, older participants are better able to retrieve specific memories than if a time limit is imposed.

Although there was a group difference in self-reported depression symptoms between the groups, both groups were below BDI-II thresholds for likely depression diagnoses. Given the non-clinical nature of the present sample, it is not surprising that there was no association between memory specificity and self-reported depression symptoms in any of the analyses. This finding is similar to other studies using the AMT (Serrano et al., 2007) and is also consistent with research in younger adults (Farina et al., 2019; Van Vreeswijk & De Wilde, 2004). Any association between problems recalling specific memories and depression seems only to emerge when comparing participants with and without depression diagnoses (Farina et al., 2019).

Due to practical constraints, the tasks were not counterbalanced. It appears possible that group effects are evident in the AMT may be because older adults fatigued by the time they completed the AMT. However, as the mean age of older adults was approximately 65, it seems unlikely that these participants would experience significantly more fatigue than their younger counterparts after undergoing the same procedure. Also, participants were given as much time to rest between tasks as they wanted. Also, there were no time limits on the SCEPT or AMI, so in the event that a participant was fatigued they could have taken longer to complete the tasks if they wanted to. It seems unlikely that group differences would emerge in the AMI but not the SCEPT if these effects were due to fatigue in one of the groups.

Despite this, and despite the relative advantage that older adults seemed to have when retrieving specific memories related to childhood, that is not to say that interventions for improving memory specificity would not be worthwhile amongst older adults. Clearly, older adults show problems with retrieving specific memories, especially when the stimuli used to cue their memories are lacking in temporal or spatial contextual information. This suggestion aligns with findings from investigations with older adults with Alzheimer's disease where

there is evidence that people with Alzheimer's may show preserved autobiographical memory but that they rely heavily on visual cues and their associated imagery to aid in the retrieval of autobiographical memories (El Haj & Gallouj, 2019). This supports the need for memory therapeutics such as Memory Specificity Training (MeST)(Barry, Sze, & Raes, 2019; Raes, Williams, & Hermans, 2009) that assist older adults in retrieving rich, detailed memories based on positive and negative cues.

The present investigation therefore suggests that, although older adults show problems retrieving specific memories when they are cued by nouns and adjectives, their problems with memory retrieval seem to be confined to this form of cuing and in procedures in which they must retrieve a memory within a limited amount of time. When memories are cued by particular life periods, and when no constraint on retrieval time is given, older adults can even outperform younger adults. Memory therapeutics might capitalise on this relative advantage when developing memory therapeutic interventions.

Table 1. Means and standard deviations

	AMT			AMI				
	SCEPT	Positive	Negative	Total	Childhood	Adulthood	Recent	Total
YA	2.68 (1.57)	4.38 (1.03)	4.26 (1.03)	8.61 (1.85)	1.98 (0.98)	2.77 (0.52)	1.82 (0.44)	6.57 (1.24)
OA	3.06 (2.45)	3.22 (1.32)	3.14 (1.38)	6.35 (2.38)	2.46 (0.67)	2.58 (0.63)	1.84 (0.80)	6.88 (1.41)

Note. Means (and standard deviations) for responses on each of the measures of autobiographical memory. YA: Younger adult; OA: Older adult; SCEPT: Sentence Completion for Events from the Past Test; AMT: Autobiographical Memory Test; AMI: Autobiographical Memory Interview.

References

- Barry, T. J., Sze, W. Y., & Raes, F. (2019). A meta-analysis and systematic review of Memory Specificity Training (MeST) in the treatment of emotional disorders. *Behaviour Research and Therapy, 116*(November 2018), 36–51.
<https://doi.org/10.1016/j.brat.2019.02.001>
- Beck, A., Steer, R., & Brown, G. (1996). *Beck Depression Inventory-II*. San Antonio, TX: Psychological Corporation. <https://doi.org/10.1037/t00742-000>
- Birch, L. S., & Davidson, K. M. (2007). Specificity of autobiographical memory in depressed older adults and its relationship with working memory and IQ. *British Journal of Clinical Psychology, 46*(2), 175–186. <https://doi.org/10.1348/014466506X119944>
- Carretero, L. M., Latorre, J. M., Fernández, D., Barry, T. J., & Ricarte, J. J. (2019). Effects of positive personal and non-personal autobiographical stimuli on emotional regulation in older adults. *Aging Clinical and Experimental Research, 0*(0), 1–8.
<https://doi.org/10.1007/s40520-019-01147-0>
- Comblain, C., D'Argembeau, A., & Van Der Linden, M. (2005). Phenomenal characteristics of autobiographical memories for emotional and neutral events in older and younger adults. *Experimental Aging Research, 31*(2), 173–189.
<https://doi.org/10.1080/03610730590915010>
- Conway, M. A., & Pleydell-Pearce, C. W. (2000). The construction of autobiographical memories in the self-memory system. *Psychological Review, 107*(2), 261–288.
<https://doi.org/10.1037/0033-295X.107.2.261>
- Craig, R., & Mindell, J. (2005). The health of older people. In *Health Survey for England: Summary of key findings*. London: National Centre for Social Research.
- De Medeiros, K., Mosby, A., Hanley, K. B., Pedraza, M. S., & Brandt, J. (2011). A randomized clinical trial of a writing workshop intervention to improve autobiographical

- memory and well-being in older adults. *International Journal of Geriatric Psychiatry*, 26(8), 803–811. <https://doi.org/10.1002/gps.2605>
- El Haj, M., & Gallouj, K. (2019). Mental imagery and autobiographical memory in Alzheimers disease. *Neuropsychology*. <https://doi.org/10.1037/neu0000521>
- Farina, F. R., Barry, T. J., van Damme, I., van Hie, T., & Raes, F. (2019). Depression diagnoses, but not individual differences in depression symptoms, are associated with reduced autobiographical memory specificity. *British Journal of Clinical Psychology*, 58(2), 173–186. <https://doi.org/10.1111/bjc.12207>
- Field, A. P., & Wilcox, R. R. (2017). Robust statistical methods: A primer for clinical psychology and experimental psychopathology researchers. *Behaviour Research and Therapy*, 98, 19–38. <https://doi.org/10.1016/j.brat.2017.05.013>
- Gallassi, R., Di Sarro, R., Morreale, A., & Amore, M. (2006). Memory impairment in patients with late-onset major depression: The effect of antidepressant therapy. *Journal of Affective Disorders*, 91(2–3), 243–250. <https://doi.org/10.1016/j.jad.2006.01.018>
- Gidron, Y., & Alon, S. (2007). Autobiographical memory and depression in the later age: The bump is a turning point. *International Journal of Aging and Human Development*, 64(1), 1–11. <https://doi.org/10.2190/96L8-6J35-0297-M403>
- Koller, M. (2016). Robustlmm: An R package for Robust estimation of linear Mixed-Effects models. *Journal of Statistical Software*, 75(1), 1–24. <https://doi.org/10.18637/jss.v075.i06>
- Kopelman, M., Wilson, B., & Baddeley, A. (1990). The autobiographical memory interview. *Thames Valley Test Company, Suffolk, UK. Lavanex, P., Amaral, D.G.*
- Latorre, J. M., Ricarte, J. J., Serrano, J. P., Ros, L., Navarro, B., & Aguilar, M. J. (2013). Performance in autobiographical memory of older adults with depression symptoms. *Applied Cognitive Psychology*, 27(2), 167–172. <https://doi.org/10.1002/acp.2891>

- Leahy, F., Ridout, N., Mushtaq, F., & Holland, C. (2017). Improving specific autobiographical memory in older adults: impacts on mood, social problem solving, and functional limitations. *Aging, Neuropsychology, and Cognition*, *00*(00), 1–29. <https://doi.org/10.1080/13825585.2017.1365815>
- Lindwall, M., Berg, A. I., Bjälkebring, P., Buratti, S., Hansson, I., Hassing, L., ... Johansson, B. (2017). Psychological health in the retirement transition: Rationale and first findings in the HHealth, Ageing and Retirement Transitions in Sweden (HEARTS) study. *Frontiers in Psychology*, *8*(SEP). <https://doi.org/10.3389/fpsyg.2017.01634>
- Maechler, M., Rousseeuw, P., Croux, C., Todorov, V., Ruckstuhl, A., Salibian-Barrera, M., ... di Palma, M. A. (2019). robustbase: Basic robust statistics (R package version 0.93-5).
- Ono, M., Devilly, G. J., & Shum, D. H. K. (2015). A meta-analytic review of overgeneral memory: The role of trauma history, mood, and the presence of posttraumatic stress disorder. *Psychological Trauma: Theory, Research, Practice, and Policy*, *8*(2), 157–164. <https://doi.org/http://dx.doi.org/10.1037/tra0000027>
- Raes, F., Hermans, D., Williams, J. M. G., & Eelen, P. (2007). A sentence completion procedure as an alternative to the autobiographical memory test for assessing overgeneral memory in non-clinical populations. *Memory*, *15*(5), 495–507. <https://doi.org/10.1080/09658210701390982>
- Raes, F., Williams, J. M. G., & Hermans, D. (2009). Reducing cognitive vulnerability to depression: A preliminary investigation of MEMory Specificity Training (MEST) in inpatients with depressive symptomatology. *Journal of Behavior Therapy and Experimental Psychiatry*, *40*(1), 24–38. <https://doi.org/10.1016/j.jbtep.2008.03.001>
- Rathbone, C. J., Holmes, E. A., Murphy, S. E., & Ellis, J. A. (2015). Autobiographical memory and well-being in aging: The central role of semantic self-images.

Consciousness and Cognition, 33, 422–431.

<https://doi.org/10.1016/j.concog.2015.02.017>

Reitzes, D. C., Mutran, E. J., & Fernandez, M. E. (1996). Does retirement hurt well-being?

Factors influencing self-esteem and depression among retirees and workers.

Gerontologist, 36(5), 649–656. <https://doi.org/10.1093/geront/36.5.649>

Ricarte, J. J., Latorre, J. M., Ros, L., Navarro, B., Aguilar, M. J., & Serrano, J. P. (2011).

Overgeneral autobiographical memory effect in older depressed adults. *Aging and*

Mental Health, 15(8), 1028–1037. <https://doi.org/10.1080/13607863.2011.573468>

Ros, L., & Latorre, J. M. (2010). Gender and age differences in the recall of affective

autobiographical memories using the autobiographical memory test. *Personality and*

Individual Differences, 49(8), 950–954. <https://doi.org/10.1016/j.paid.2010.08.002>

Rubin, D. C., & Schulkind, M. D. (1997). The distribution of autobiographical memories

across the lifespan. *Memory and Cognition*, 25(6), 859–866.

<https://doi.org/10.3758/BF03211330>

Serrano, J. P., Latorre, J. M., & Gatz, M. (2007). Autobiographical memory in older adults

with and without depressive symptoms. *International Journal of Clinical and Health*

Psychology, 7(1), 41–57.

St. Jacques, P. L., & Levine, B. (2007). Ageing and autobiographical memory for emotional

and neutral events. *Memory*, 15(2), 129–144.

<https://doi.org/10.1080/09658210601119762>

Tollenaar, M. S., Elzinga, B. M., Spinhoven, P., & Everaerd, W. (2009). Autobiographical

memory after acute stress in healthy young men. *Memory*, 17(3), 301–310.

<https://doi.org/10.1080/09658210802665845>

Van Vreeswijk, M. F., & De Wilde, E. J. (2004). Autobiographical memory specificity,

psychopathology, depressed mood and the use of the Autobiographical Memory Test: A

meta-analysis. *Behaviour Research and Therapy*, 42(6), 731–743.

[https://doi.org/10.1016/S0005-7967\(03\)00194-3](https://doi.org/10.1016/S0005-7967(03)00194-3)

Williams, J. M. G., Barnhofer, T., Crane, C., & Beck, A. T. (2005). Problem solving deteriorates following mood challenge in formerly depressed patients with a history of suicidal ideation. *Journal of Abnormal Psychology*, 114(3), 421–431.

<https://doi.org/10.1037/0021-843X.114.3.421>

Williams, J. M. G., & Broadbent, K. (1986). Autobiographical memory in suicide attempters. *Journal of Abnormal Psychology*, 95, 144–149. [https://doi.org/10.1037/0021-](https://doi.org/10.1037/0021-843X.95.2.144)

843X.95.2.144

Wilson, F. C. L., & Gregory, J. D. (2018). Overgeneral autobiographical memory and depression in older adults: a systematic review. *Aging and Mental Health*. Taylor &

Francis. <https://doi.org/10.1080/13607863.2017.1326461>