



Homological branching law for $(\mathrm{GL}_{n+1}(F), \mathrm{GL}_n(F))$: projectivity and indecomposability

Kei Yuen Chan¹

Received: 20 December 2019 / Accepted: 1 January 2021 /
Published online: 15 February 2021
© The Author(s) 2021

Abstract Let F be a non-Archimedean local field. This paper studies homological properties of irreducible smooth representations restricted from $\mathrm{GL}_{n+1}(F)$ to $\mathrm{GL}_n(F)$. A main result shows that each Bernstein component of an irreducible smooth representation of $\mathrm{GL}_{n+1}(F)$ restricted to $\mathrm{GL}_n(F)$ is indecomposable. We also classify all irreducible representations which are projective when restricting from $\mathrm{GL}_{n+1}(F)$ to $\mathrm{GL}_n(F)$. A main tool of our study is a notion of left and right derivatives, extending some previous work joint with Gordan Savin. As a by-product, we also determine the branching law in the opposite direction.

1 Introduction

1.1

Let F be a non-Archimedean local field. Let $G_n = \mathrm{GL}_n(F)$. Let $\mathrm{Alg}(G_n)$ be the category of smooth representations of G_n . This paper is a sequel of [15] in studying homological properties of smooth representations of G_{n+1} restricted to G_n , which is originally motivated from the study of D. Prasad in his ICM proceeding [28]. In [15], we show that for generic representations π and π' of G_{n+1} and G_n respectively, the higher Ext-groups

$$\mathrm{Ext}_{G_n}^i(\pi, \pi') = 0, \text{ for } i \geq 1,$$

✉ Kei Yuen Chan
kychan@fudan.edu.cn

¹ Shanghai Center for Mathematical Sciences, Fudan University, Shanghai, China

which was previously conjectured in [28]. This result gives a hope that there is an explicit homological branching law, generalizing the multiplicity one theorem [2, 31] and the local Gan–Gross–Prasad conjectures [17, 18].

The main techniques in [15] are utilizing Hecke algebra structure developed in [13, 14] and simultaneously applying left and right Bernstein–Zelevinsky derivatives, based on the classical approach of using Bernstein–Zelevinsky filtration on representations of G_{n+1} restricted to G_n [27, 28]. We shall extend these methods further, in combination of other things, to obtain new results in this paper.

In [15], we showed that an essentially square-integrable representation π of G_{n+1} is projective when restricted to G_n . However, those representations do not account for all irreducible representations whose restriction is projective. The first goal of the paper is to classify such representations:

Theorem 1.1 (=Theorem 3.7) *Let π be an irreducible smooth representation of G_{n+1} . Then $\pi|_{G_n}$ is projective if and only if*

- (1) π is essentially square-integrable, or
- (2) $n + 1$ is even, and $\pi \cong \rho_1 \times \rho_2$ for some cuspidal representations ρ_1, ρ_2 of $G_{(n+1)/2}$.

There are also recent studies of the projectivity under restriction in other settings [1, 16, 22].

It is a well-known result that an irreducible representation of a reductive group G is projective (modulo center) if and only if it is supercuspidal, giving a tight connection between homological algebra and harmonic analysis. The classification theorem above for (G_{n+1}, G_n) suggests that one may still expect such connection in a relative setting. It would be interesting to see the interplay of our study with the harmonic analysis study in the relative Langlands program [30].

A main step in our classification is the following projectivity criteria:

Theorem 1.2 (=Theorem 3.3) *Let π be an irreducible smooth representation of G_{n+1} . Then $\pi|_{G_n}$ is projective if and only if the following two conditions hold:*

- (1) π is generic; and
- (2) $\text{Hom}_{G_n}(\pi|_{G_n}, \omega) = 0$ for any irreducible non-generic representation ω of G_n .

Theorem 1.2 turns the projectivity problem into a Hom-branching problem and is a consequence of two results: (1) the Euler–Poincaré pairing formula of D. Prasad [28] and (2) the Hecke algebra argument used in [15] by G. Savin and the author. Roughly speaking, (1) is used to show non-projectivity while (2) is used to show projectivity.

The second part of the paper studies indecomposability of a restricted representation. It is clear that an irreducible representation (except one-dimensional ones) restricted from G_{n+1} to G_n cannot be indecomposable as it has more than one non-zero Bernstein components. However, the Hecke algebra realization in [13, 15] of the projective representations in Theorem 1.1 immediately implies that each Bernstein component of those restricted representation is indecomposable. This is a motivation of our study in general case, and precisely we prove:

Theorem 1.3 (=Theorem 6.1) *Let π be an irreducible representation of G_{n+1} . Then for each Bernstein component τ of $\pi|_{G_n}$, any two non-zero G_n -submodules of τ have non-zero intersection.*

As a consequence, we have:

Corollary 1.4 *Let π be an irreducible smooth representation of G_{n+1} . Then any Bernstein component of $\pi|_{G_n}$ is indecomposable.*

In Sect. 6.2, we explain how to determine which Bernstein component of $\pi|_{G_n}$ is non-zero in terms of Zelevinsky segments, and hence Theorem 1.3 essentially parametrizes the indecomposable components of $\pi|_{G_n}$.

For a mirabolic subgroup M_n of G_{n+1} , it is known [35] that $\pi|_{M_n}$ is indecomposable for an irreducible representation π of G_{n+1} . The approach in [35] uses the Bernstein–Zelevinsky filtration of π to M_n and that the bottom piece of the filtration is irreducible. We prove that the Bernstein component of a bottom piece is indecomposable as a G_n -module, and then make use of left and right derivatives, developed and used to prove main results in [15]. The key fact is that left and right derivatives of an irreducible representation are asymmetric. We now make more precise the meaning of ‘asymmetric’. We say that an integer i is the level of an irreducible representation π of G_n if the left derivative $\pi^{(i)}$ is the highest derivative of π .

Theorem 1.5 (=Theorem 6.2) *Let π be an irreducible smooth representation of G_n . Let $v(g) = |\det(g)|_F$. Suppose i is not the level of π . Then $v^{1/2} \cdot \pi^{(i)}$ and $v^{-1/2} \cdot {}^{(i)}\pi$ have no isomorphic irreducible quotients whenever $v^{1/2} \cdot \pi^{(i)}$ and $v^{-1/2} \cdot {}^{(i)}\pi$ are non-zero. The statement also holds if one replaces quotients by submodules.*

We remark that $v^{1/2} \cdot \pi^{(i)}$ and $v^{-1/2} \cdot {}^{(i)}\pi$ are shifted derivatives in the sense of [4], which have been used in *loc. cit.* to study unitary representations of G_n .

As a by-product of Theorem 1.5, we give a complete answer to the branching law in another direction:

Theorem 1.6 (=Corollary 7.7) *Let π_1, π_2 be smooth irreducible representations of G_{n+1} and G_n respectively. Then*

$$\mathrm{Hom}_{G_n}(\pi_2, \pi_1|_{G_n}) \neq 0$$

if and only if both π_1 and π_2 are one-dimensional and $\pi_2 \cong \pi_1|_{G_n}$.

Computing the structure of a derivative of an arbitrary representation is a difficult question in general. Our approach is to approximate the information of derivatives of irreducible ones by some parabolically induced modules, whose derivatives can be computed via geometric lemma. On the other hand, the Speh representations behave more symmetrically for left and right derivatives, which motivates our proof to involve Speh representations.

Another key ingredient in proving Theorem 1.3 is a study on the submodule structure between left and right Bernstein–Zelevinsky inductions. We explain in Sect. 5 how the submodule structure of an induced module can be partly reflected from the module which induced from (see Proposition 5.6 for a precise statement). The study relies on the Hecke algebra structure of the Gelfand–Graev representation.

A related question to Theorem 1.3 is that the Bernstein–Zelevinsky induction functor preserves indecomposability at the level of each Bernstein component, which is shown in Theorem 8.3. This is in contrast with that the (usual) parabolic induction does not preserve indecomposability in general. In [12], we study some special situations of the parabolic induction that one can obtain certain indecomposability-preserving results, which have applications to the local nontempered Gan–Gross–Prasad conjectures [18].

1.2 Organization of the paper

Section 2 studies derivatives of generic representations, which simplifies some computations for Theorem 1.1. The results also give some guiding examples in the study of this paper and [15].

Section 3 firstly proves a criteria for an irreducible representation to be projective under restriction, and then apply this criteria to give a classification of such class of modules.

Section 4 discusses some results on Gelfand–Graev representations such as their indecomposability while Sect. 5 develops a theory of the submodule relation between left and right Bernstein–Zelevinsky inductions.

Section 6 proves a main result on the indecomposability of an irreducible representation under restriction, which uses results in Sect. 5 and the asymmetric property of derivatives proved in Sect. 7.

Section 8 proves that the Bernstein–Zelevinsky induction preserves indecomposability. This partly generalizes Sect. 5.

In the first appendix, we explain how an irreducible representation appears as the unique submodule of a product of some Speh representations. In the second appendix, we provide some preliminaries on module theory.

2 Bernstein–Zelevinsky derivatives of generic representations

2.1 Notations

Let $G_n = GL_n(F)$. All representations in this paper are smooth and over \mathbb{C} , and we usually omit the adjective ‘smooth’. Let $\text{Irr}(G_n)$ be the set of all irreducible representations of G_n and let $\text{Irr} = \sqcup_n \text{Irr}(G_n)$. For an admissible representation $\omega \in \text{Alg}(G_n)$, denote by $\text{JH}(\omega)$ the set of (isomorphism classes of) irreducible composition factors of ω . For $\pi \in \text{Alg}(G_n)$, let π^\vee be the smooth dual of π .

Let ρ be an (irreducible) cuspidal representation of G_l . Let $a, b \in \mathbb{C}$ with $b - a \in \mathbb{Z}_{\geq 0}$. We have a Zelevinsky segment $\Delta = [v^a \rho, v^b \rho]$, which we may simply call a segment. Denote $a(\Delta) = v^a \rho$ and $b(\Delta) = v^b \rho$. The relative length of Δ is defined as $b - a + 1$ and the absolute length of Δ is defined as $l(b - a + 1)$. We can truncate Δ from each side to obtain two segments of absolute length $r(b - a)$:

$${}^-\Delta = [v^{a+1} \rho, \dots, v^b \rho] \text{ and } \Delta^- = [v^a \rho, \dots, v^{b-1} \rho].$$

Moreover, if we perform the truncation k -times, the resulting segments will be denoted by ${}^{(kl)}\Delta$ and $\Delta^{(kl)}$. We remark that the convention here is different from the previous paper [15] for convenience later. If i is not an integer divisible by l , then we set $\Delta^{(i)}$ and ${}^{(i)}\Delta$ to be empty sets. We also denote $\Delta^\vee = [v^{-b} \rho^\vee, v^{-a} \rho^\vee]$. For a singleton segment $[\rho, \rho]$, we abbreviate as $[\rho]$. For $\pi \in \text{Alg}(G_l)$, define $n(\pi) = l$. For $c \in \mathbb{C}$ and a segment $\Delta = [v^a \rho, v^b \rho]$, define $v^c \Delta = [v^{a+c} \rho, v^{b+c} \rho]$.

For a Zelevinsky segment Δ , define $\langle \Delta \rangle$ and $\text{St}(\Delta)$ to be the (unique) irreducible submodule and quotient of $v^a \rho \times \dots \times v^b \rho$ respectively. We have

$$\text{St}(\Delta)^\vee \cong \text{St}(\Delta^\vee) \text{ and } \langle \Delta \rangle^\vee \cong \langle \Delta^\vee \rangle.$$

For two cuspidal representations ρ_1, ρ_2 of G_m , we say that ρ_1 precedes ρ_2 , denoted by $\rho_1 < \rho_2$, if $v^c \rho_1 \cong \rho_2$ for some $c \in \mathbb{Z}_{>0}$. We say two segments Δ and Δ' are *linked* if $\Delta \not\subset \Delta'$, $\Delta' \not\subset \Delta$ and $\Delta \cup \Delta'$ is still a segment. We say that a segment Δ *precedes* Δ' , denoted by $\Delta < \Delta'$, if $b(\Delta)$ precedes $b(\Delta')$; and Δ and Δ' are linked. If Δ does not precede Δ' , write $\Delta \not< \Delta'$.

A multisegment is a multiset of finite numbers of segments. Let Mult be the set of multisegments. Let $\mathfrak{m} = \{\Delta_1, \dots, \Delta_r\} \in \text{Mult}$. We relabel the segments in \mathfrak{m} such that for $i < j$, Δ_i does not precede Δ_j . The modules defined below are independent of the labeling (up to isomorphisms) [35]. Define $\zeta(\mathfrak{m}) = \langle \Delta_1 \rangle \times \dots \times \langle \Delta_r \rangle$. Denote by $\langle \mathfrak{m} \rangle$ the unique irreducible

submodule of $\zeta(\mathfrak{m})$.¹ Similarly, define $\lambda(\mathfrak{m}) = \text{St}(\Delta_1) \times \dots \times \text{St}(\Delta_r)$. Denote by $\text{St}(\mathfrak{m})$ the unique quotient of $\lambda(\mathfrak{m})$. Both notions $\langle \mathfrak{m} \rangle$ and $\text{St}(\mathfrak{m})$ give a classification of irreducible smooth representations of G_n ([35, Proposition 6.1], also see [24]) i.e. both the maps

$$\mathfrak{m} \mapsto \langle \mathfrak{m} \rangle, \quad \text{and} \quad \mathfrak{m} \mapsto \text{St}(\mathfrak{m})$$

determine bijections from Mult to Irr . For example, when $\mathfrak{m} = \{[\nu^{-1/2}], [\nu^{1/2}]\}$, $\langle \mathfrak{m} \rangle$ is the Steinberg representation and $\text{St}(\mathfrak{m})$ is the trivial representation. The two notions $\langle \mathfrak{m} \rangle$ and $\text{St}(\mathfrak{m})$ are related by the so-called Aubert-Zelevinsky duality, and Mœglin-Waldspurger algorithm.

We shall use the following results several times (see [35, Theorems 1.9, 4.2 and 9.7]): if two segments Δ, Δ' are not linked, then

$$\langle \Delta \rangle \times \langle \Delta' \rangle \cong \langle \Delta' \rangle \times \langle \Delta \rangle, \tag{2.1}$$

$$\text{St}(\Delta) \times \text{St}(\Delta') \cong \text{St}(\Delta') \times \text{St}(\Delta). \tag{2.2}$$

Let $\pi \in \text{Irr}(G_n)$. Then π is a subquotient of $\rho_1 \times \dots \times \rho_r$ for some cuspidal representations ρ_i of G_{n_i} . The multiset (ρ_1, \dots, ρ_r) , denoted by $\text{csupp}(\pi)$, is called the cuspidal support of π . We also set

$$\text{csupp}_{\mathbb{Z}}(\pi) = \{v^c \rho : \rho \in \text{csupp}(\pi), c \in \mathbb{Z}\},$$

which is regarded as a set (rather than a multiset).

2.2 Derivatives and Bernstein–Zelevinsky inductions

Let U_n (resp. U_n^-) be the group of unipotent upper (resp. lower) triangular matrices in G_n . For $i \leq n$, let P_i be the parabolic subgroup of G_n containing the block diagonal matrices $\text{diag}(g_1, g_2)$ ($g_1 \in G_i, g_2 \in G_{n-i}$) and the upper triangular matrices. Let $P_i = M_i N_i$ with the Levi M_i and the unipotent N_i . Let N_i^- be the opposite unipotent subgroup of N_i . Let $\nu : G_n \rightarrow \mathbb{C}$ given by $\nu(g) = |\det(g)|_F$. Let

$$R_{n-i} = \left\{ \begin{pmatrix} g & x \\ 0 & u \end{pmatrix} \in G_n : g \in \text{GL}_{n-i}(F), u \in U_i, x \in \text{Mat}_{n-i,i}(F) \right\}.$$

Let R_{n-i}^- be the transpose of R_{n-i} .

We shall use Ind for normalized induction and ind for normalized induction with compact support. Let ψ_i be a character on U_i given by $\psi_i(u) = \overline{\psi}(u_{1,2} +$

¹ In [35], the notion of preceding is defined for $a(\Delta)$ instead of $b(\Delta)$. Our notion of preceding gives the same classification due to (2.1).

$\dots + u_{i-1,i}$), where $\overline{\psi}$ is a nondegenerate character on F and $u_{k,k+1}$ is the value in the $(k, k + 1)$ -entry of u . For $\tau \in \text{Irr}(G_{n-i})$, we extend trivially the $G_{n-i} \times U_i$ -representation $\tau \boxtimes \psi_i$ to a R_{n-i} -representation. This defines functors from $\text{Alg}(G_{n-i})$ to $\text{Alg}(G_n)$ given by

$$\pi \mapsto \text{Ind}_{R_{n-i}}^{G_n} \pi \boxtimes \psi_i, \quad \text{and } \pi \mapsto \text{ind}_{R_{n-i}}^{G_n} \pi \boxtimes \psi_i,$$

both of which will be called (right) Bernstein–Zelevinsky inductions. Similarly, one has left Bernstein–Zelevinsky inductions by using R_{n-i}^- instead of R_{n-i} .

Let $\pi \in \text{Alg}(G_n)$. Following [15], define $\pi^{(i)}$ to be the left adjoint functor of $\text{Ind}_{R_{n-i}}^{G_n} \pi \boxtimes \psi_i$. Let $\theta_n : G_n \rightarrow G_n$ given by $\theta_n(g) = g^{-T}$, the inverse transpose of g . Define the left derivative

$${}^{(i)}\pi := \theta_{n-i}(\theta_n(\pi))^{(i)}, \tag{2.3}$$

which is left adjoint to $\text{Ind}_{R_{n-i}^-}^{G_n} \pi \boxtimes \psi'_i$, where $\psi'_i(u) = \psi_i(u^T)$ for $u \in U_i^-$. The level of an admissible representation π is the largest integer i^* such that $\pi^{(i^*)} \neq 0$ and $\pi^{(j)} = 0$ for all $j > i^*$. It follows from (2.3) that if i^* is the level of π , then ${}^{(i^*)}\pi \neq 0$ and ${}^{(j)}\pi = 0$ for all $j > i^*$. When i^* is the level for π , we shall call $\pi^{(i^*)}$ and ${}^{(i^*)}\pi$ to be the highest right and left derivatives of π respectively, where we usually drop the term of left and right if no confusion.

We now define the shifted derivatives as follow: for any i ,

$$\pi^{[i]} = \nu^{1/2} \cdot \pi^{(i)}, \quad \text{and } [i]\pi = \nu^{-1/2} \cdot {}^{(i)}\pi.$$

For the details of Bernstein–Zelevinsky filtrations, see [15].

The following result [7, Corollary 4.6] will be used several times, which is a consequence of the geometric lemma: for $\pi_k \in \text{Alg}(G_{n_k})$ ($k = 1, \dots, r$), for any i ,

$$(\pi_1 \times \dots \times \pi_r)^{(i)}$$

admits a filtration whose successive quotients isomorphic to

$$(\pi_1)^{(i_1)} \times \dots \times (\pi_r)^{(i_r)}, \tag{2.4}$$

where i_1, \dots, i_r run for all integers satisfying $i_1 + \dots + i_r = i$.

2.3 On computing derivatives

Let $\Pi_i = \text{ind}_{U_i}^{G_i} \psi_i$ be the Gelfand–Graev representation. Using inductions by stages, we have that

$$\text{ind}_{R_{n-i}}^{G_n} \pi \boxtimes \psi_i \cong \pi \times \text{ind}_{U_i}^{G_i} \psi_i = \pi \times \Pi_i.$$

Let $\dot{w}_0 \in G_n$ whose antidiagonal entries are 1 and other entries are 0. By the left translation of \dot{w}_0 on $\text{ind}_{R_{n-i}}^{G_n} \pi \boxtimes \psi_i$, we have that

$$\text{ind}_{R_{n-i}}^{G_n} \pi \boxtimes \psi_i \cong \left(\text{ind}_{U_i}^{G_i} \psi_i \right) \times \pi = \Pi_i \times \pi.$$

Since the right derivative is left adjoint to $\text{Ind}_{R_{n-i}}^{G_n}$, we consequently have:

$$\pi^{(i)} \cong (\pi_{N_i})_{U_i, \psi_i},$$

where U_i is regarded as the subgroup $G_{n-i} \times G_i$ via the embedding $g \mapsto \text{diag}(1, g)$. We shall use the later expression when computing derivatives in Sect. 3.2. We also have an analogous expression for left derivatives.

We shall often use the following lemma:

Lemma 2.1 *Let $\pi \in \text{Alg}(G_{n+1})$ and let π' be an admissible representation of G_n . Suppose there exists i such that the following conditions hold:*

$$\text{Hom}_{G_{n+1-i}}(\pi^{[i]}, {}^{(i-1)}\pi') \neq 0;$$

and

$$\text{Ext}_{G_{n+1-j}}^k(\pi^{[j]}, {}^{(j-1)}\pi') = 0$$

for all $j = 1, \dots, i - 1$ and all k . Then $\text{Hom}_{G_n}(\pi, \pi') \neq 0$.

Proof The Bernstein–Zelevinsky filtration of π gives that there exists

$$0 \subset \pi_n \subset \dots \subset \pi_0 = \pi$$

such that $\pi_{i-1}/\pi_i \cong \text{ind}_{R_{n-i+1}}^{G_n} \pi^{[i]} \boxtimes \psi_i$. Now, by the second adjointness property of derivatives [15, Lemma 2.2], we have

$$\text{Ext}_{G_n}^k(\text{ind}_{R_{n-j+1}}^{G_n} \pi^{[j]} \boxtimes \psi_{j-1}, \pi') \cong \text{Ext}_{G_{n-j+1}}^k(\pi^{[j]}, {}^{(j-1)}\pi') = 0$$

for all k and j . Now a long exact sequence argument gives that

$$\dim \text{Hom}_{G_n}(\pi, \pi') \geq \dim \text{Hom}_{G_n}(\text{ind}_{R_{n+1-i}}^{G_n} \pi^{[i]} \boxtimes \psi_{i-1}, \pi') \tag{2.5}$$

$$= \dim \text{Hom}_{G_{n+1-i}}(\pi^{[i]}, {}^{(i-1)}\pi') \neq 0. \tag{2.6}$$

□

2.4 Subrepresentation of a standard representation

Lemma 2.2 *Let $\mathfrak{m} \in \text{Mult}$. Suppose all segments in \mathfrak{m} are singletons i.e. of relative length 1. Then $\lambda(\mathfrak{m})$ has unique irreducible submodule and quotient. Moreover, the unique irreducible submodule is generic.*

Proof By definitions, $\lambda(\mathfrak{m}) = \zeta(\mathfrak{m})$ and hence has unique submodule and quotient. Since all segments in \mathfrak{m} are singletons, the submodule is generic [35]. □

It is known (see [21]²) that $\lambda(\mathfrak{m})$ always has a generic representation as the unique submodule. We shall prove a slightly stronger statement, using Zelevinsky theory:

Proposition 2.3 *Let $\mathfrak{m} \in \text{Mult}$. Then $\lambda(\mathfrak{m})$ can be embedded to $\lambda(\mathfrak{m}')$ for some $\mathfrak{m}' \in \text{Mult}$ whose segments are singletons. In particular, $\lambda(\mathfrak{m})$ has a unique irreducible submodule and moreover, the submodule is generic.*

Proof Let ρ be a cuspidal representation in \mathfrak{m} such that for any cuspidal representation ρ' in \mathfrak{m} , $\rho \not\cong \rho'$. Let Δ be a segment in \mathfrak{m} with the shortest relative length among all segments Δ' in \mathfrak{m} with $b(\Delta') \cong \rho$.

By definition of $\lambda(\mathfrak{m})$, we have that

$$\lambda(\mathfrak{m}) \cong \text{St}(\Delta) \times \lambda(\mathfrak{m} \setminus \{\Delta\}).$$

On the other hand, we have that

$$\text{St}(\Delta) \hookrightarrow b(\Delta) \times \text{St}(\Delta^-).$$

Thus we have that

$$\lambda(\mathfrak{m}) \hookrightarrow b(\Delta) \times \text{St}(\Delta^-) \times \lambda(\mathfrak{m} \setminus \{\Delta\}) \tag{2.7}$$

$$\cong b(\Delta) \times \lambda(\mathfrak{m} \setminus \{\Delta\} + \Delta^-) \tag{2.8}$$

² The author would like to thank D. Prasad for mentioning this reference in a discussion.

We now explain the last isomorphism (2.8), and it suffices to show

$$\lambda(\mathfrak{m} \setminus \{\Delta\} + \Delta^-) \cong \text{St}(\Delta^-) \times \lambda(\mathfrak{m} \setminus \{\Delta\}).$$

To this end, we write $\mathfrak{m} = \{\Delta, \Delta_1, \dots, \Delta_k, \Delta_{k+1}, \dots, \Delta_s\}$ such that

$$b(\Delta) \cong b(\Delta_1) \cong \dots \cong b(\Delta_k)$$

and $b(\Delta_j) \not\cong b(\Delta)$ for $j \geq k + 1$. Then we have that

$$\begin{aligned} \text{St}(\Delta^-) \times \lambda(\mathfrak{m} \setminus \{\Delta\}) &\cong \text{St}(\Delta^-) \times \text{St}(\Delta_1) \times \dots \times \text{St}(\Delta_s) \\ &\cong \text{St}(\Delta_1) \times \dots \times \text{St}(\Delta_k) \times \text{St}(\Delta^-) \\ &\quad \times \text{St}(\Delta_{k+1}) \times \dots \times \text{St}(\Delta_s) \\ &\cong \lambda(\mathfrak{m} \setminus \{\Delta\} + \Delta^-) \end{aligned}$$

where the second isomorphism follows from applying (2.2) few times. Here we need to use our choice of Δ , which guarantees that Δ^- and Δ_j are unlinked for $j = 1, \dots, k$.

Now $\lambda(\mathfrak{m} \setminus \{\Delta\} + \Delta^-)$ embeds to $\lambda(\mathfrak{m}')$ by induction for some $\mathfrak{m}' \in \text{Mult}$ with all segments to be singletons. Thus this gives that $b(\Delta) \times \lambda(\mathfrak{m} \setminus \{\Delta\} + \Delta^-)$ embeds to $b(\Delta) \times \lambda(\mathfrak{m}') \cong \lambda(\mathfrak{m}' + b(\Delta))$, and so does $\lambda(\mathfrak{m})$ by (2.7).

The second assertion follows from Lemma 2.2. □

2.5 Derivatives of generic representations

Recall that the socle (resp. cosocle) of an admissible representation π of G_n is the maximal semisimple submodule (resp. quotient) of π .

Lemma 2.4 *Let $\pi \in \text{Irr}(G_n)$. Then the cosocle of $\pi^{(i)}$ (resp. ${}^{(i)}\pi$) is isomorphic to the socle of $\pi^{(i)}$ (resp. ${}^{(i)}\pi$).*

Proof It follows from that for an irreducible G_n -representation π ,

$${}^{(i)}\pi \cong \theta_{n-i}(\theta_n(\pi)^{(i)}) \cong \theta_{n-i}((\pi^\vee)^{(i)}) \cong \theta_{n-i}(({}^{(i)}\pi)^\vee)$$

and the fact [6] that $\theta_{n-i}(\tau) \cong \tau^\vee$ for any irreducible G_{n-i} -representation τ . The last isomorphism follows from [15, Lemma 2.2]. □

Proposition 2.5 *Let $\pi \in \text{Irr}(G_{n+1})$. Both socle and cosocle of $\pi^{(i)}$ (and ${}^{(i)}\pi$) are multiplicity-free.*

Proof Let π_0 be an irreducible quotient of $\pi^{(i)}$. Let π_1 be a cuspidal representation of G_{i-1} which is not in $\text{cupp}_{\mathbb{Z}}(\pi_0)$. Then, by comparing cuspidal supports,

$$\text{Ext}_{G_{n+1-k}}^j(\pi^{[k], (k-1)}(\pi_0 \times \pi_1)) = 0$$

for all j and $k < i$. With a long exact sequence argument using Bernstein–Zelevinsky filtration (similar to the proof of Lemma 2.1), we have that

$$\dim \text{Hom}_{G_n}(\pi, \pi_0 \times \pi_1) \geq \dim \text{Hom}_{G_{n+1-i}}(\pi^{[i], (i-1)}(\pi_0 \times \pi_1))$$

Now one applies the geometric lemma to obtain a filtration on $\pi^{(i-1)}(\pi_0 \times \pi_1)$, and then by comparing cuspidal supports, the only possible layer that can contribute the above non-zero Hom is π_0 . Hence,

$$\dim \text{Hom}_{G_n}(\pi, \pi_0 \times \pi_1) \geq \dim \text{Hom}_{G_{n+1-i}}(\pi^{[i]}, \pi_0).$$

The first dimension is at most one by [2] and so is the second dimension. This implies the cosocle statement by Lemma 2.1 and the socle statement follows from Lemma 2.4. □

A representation π of G_n is called *generic* if $\pi^{(n)} \neq 0$. The Zelevinsky classification of irreducible generic representations is in [35], that is, $\text{St}(\mathfrak{m})$ is generic if and only if any two segments in \mathfrak{m} are unlinked. With Proposition 2.5, the following result essentially gives a combinatorial description on the socle and cosocle of the derivatives of a generic representation.

Corollary 2.6 *Let $\pi \in \text{Irr}(G_{n+1})$ be generic. Then any simple quotient and submodule of $\pi^{(i)}$ (resp. ${}^{(i)}\pi$) is generic.*

Proof By Lemma 2.4, it suffices to prove the statement for quotient. Let $\mathfrak{m} = \{\Delta_1, \dots, \Delta_r\}$ be the Zelevinsky segment \mathfrak{m} such that

$$\pi \cong \text{St}(\mathfrak{m}) = \lambda(\mathfrak{m}) = \text{St}(\Delta_1) \times \dots \times \text{St}(\Delta_r).$$

Since any two segments in \mathfrak{m} are unlinked, we can label in any order and so we shall assume that for $i < j$, $b(\Delta_j)$ does not precede $b(\Delta_i)$. Then geometric lemma (2.4) produces a filtration on $\pi^{(i)}$ whose successive subquotient is isomorphic to

$$\text{St}^{(i_1)} \Delta_1 \times \dots \times \text{St}^{(i_r)} \Delta_r,$$

where $i_1 + \dots + i_r = i$. The last module is isomorphic to $\lambda(\mathfrak{m}')^\vee$, where

$$\mathfrak{m}' = \left\{ ({}^{(i_1)} \Delta_1)^\vee, \dots, ({}^{(i_r)} \Delta_r)^\vee \right\}.$$

If π' is a simple quotient of $\pi^{(i)}$, then π' is a simple quotient of one successive subquotient in the filtration, or in other words, π' is a simple submodule of $\lambda(\mathfrak{m}')$ for a multisegment \mathfrak{m}' . Now the result follows from Proposition 2.3. \square

Remark 2.7 One can formulate the corresponding statement of Proposition 2.5 for affine Hecke algebra level by using the sign projective module in [13]. Then it might be interesting to ask for an analogue result for affine Hecke algebras over fields of positive characteristics.

Here we give a consequence to branching law (c.f. [12, 18, 19]):

Corollary 2.8 *Let $\pi \in \text{Irr}(G_{n+1})$ be generic. Let $\pi' \in \text{Irr}(G_n)$ and let $\mathfrak{m} \in \text{Mult}$ with $\pi' \cong \langle \mathfrak{m} \rangle$. If $\text{Hom}_{G_n}(\pi, \langle \mathfrak{m} \rangle) \neq 0$, then each segment in \mathfrak{m} has relative length at most 2.*

Proof Write $\mathfrak{m} = \{\Delta_1, \dots, \Delta_r\}$ such that Δ_i does not precede Δ_j if $i < j$. Let $\pi_0 = \langle \mathfrak{m} \rangle$. By using the Bernstein–Zelevinsky filtration, $\text{Hom}_{G_n}(\pi, \pi_0) \neq 0$ implies that

$$\text{Hom}_{G_{n+1-i}}(\pi_1^{[i]}, {}^{(i-1)}\pi_0) \neq 0$$

for some $i \geq 1$ [15] (c.f. Lemma 2.1). Hence Corollary 2.6 implies that ${}^{(i-1)}\pi_0$ is generic for some $i \geq 1$. Since

$$\pi_0 \hookrightarrow \zeta(\mathfrak{m})$$

we have ${}^{(i-1)}\pi_0 \hookrightarrow {}^{(i-1)}\zeta(\mathfrak{m})$ and so ${}^{(i-1)}\zeta(\mathfrak{m})$ has a generic composition factor. On the other hand, geometric lemma gives that ${}^{(i-1)}\pi_0$ admits a filtration whose successive quotients are isomorphic to ${}^{(i_1)}\langle \Delta_1 \rangle \times \dots \times {}^{(i_r)}\langle \Delta_r \rangle$ where i_1, \dots, i_r run for all sums equal to $i - 1$. Then at least one such quotient is non-degenerate and so in that quotient, all ${}^{(i_k)}\langle \Delta_k \rangle$ are cuspidal. Following from the derivatives on $\langle \Delta \rangle$, Δ can have at most of relative length 2. \square

Another consequence is on the indecomposability of derivatives of generic representations.

Corollary 2.9 *Let $\pi \in \text{Irr}(G_{n+1})$ be generic. Then the projections of $\pi^{(i)}$ and ${}^{(i)}\pi$ to any cuspidal support component have unique simple quotient and submodule. In particular, the projections of $\pi^{(i)}$ and ${}^{(i)}\pi$ to any cuspidal support component are indecomposable.*

Proof For a fixed cuspidal support, there is a unique (up to isomorphism) irreducible smooth generic representation. Now the result follows from Proposition 2.5 and Corollary 2.6. \square

The analogous statement is not true in general if one replaces an irreducible generic representation by an arbitrary irreducible representation. We give an example below.

Example 2.10 Let $\mathfrak{m} = \{[1, \nu], [\nu^{-1}, 1], [1], [\nu^{-1}, \nu]\}$. We note that

$$\langle \mathfrak{m} \rangle = \langle [1, \nu], [\nu^{-1}, 1] \rangle \times \langle [1] \rangle \times \langle [\nu^{-1}, \nu] \rangle.$$

We compute the components of $\langle \mathfrak{m} \rangle^{(2)}$, at the cuspidal support $\{\nu^{-1}, \nu^{-1}, 1, 1, 1, \nu\}$. In this case, there are four composition factors whose Zelevinsky multisegments are:

$$\begin{aligned} \mathfrak{m}_1 &= \{[1], [\nu^{-1}], [1], [\nu^{-1}, \nu]\}, & \mathfrak{m}_2 &= \{[1, \nu], [\nu^{-1}], [1], [\nu^{-1}, 1]\} \\ \mathfrak{m}_3 &= \{[1, \nu], [\nu^{-1}, 1], [\nu^{-1}, 1]\} \text{ (multiplicity 2)} \end{aligned}$$

Note that the socle and cosocle coincide by Lemma 2.4. Note $\langle \mathfrak{m}_1 \rangle$ and $\langle \mathfrak{m}_3 \rangle$ are in socle (and so cosocle). $\langle \mathfrak{m}_2 \rangle$ cannot be a submodule or quotient. Thus the only possible structure is two indecomposable modules. One of them has composition factors of $\langle \mathfrak{m}_3 \rangle$ (with multiplicity 2) and $\langle \mathfrak{m}_2 \rangle$. Another one is a simple module isomorphic to $\langle \mathfrak{m}_1 \rangle$.

3 Projectivity

3.1 Projectivity criteria

We need the following formula of D. Prasad:

Theorem 3.1 [28] *Let π_1 and π_2 be admissible representations of $GL_{n+1}(F)$ and $GL_n(F)$ respectively. Then*

$$\sum_{i \in \mathbb{Z}} (-1)^i \dim \text{Ext}_{G_n}^i(\pi_1, \pi_2) = \dim \text{Wh}(\pi_1) \cdot \dim \text{Wh}(\pi_2),$$

where $\text{Wh}(\pi_1) = \pi_1^{(n+1)}$ and $\text{Wh}(\pi_2) = \pi_2^{(n)}$.

Lemma 3.2 *Let $\pi \in \text{Irr}(G_{n+1})$. If $\pi^{(i)}$ has a non-generic irreducible submodule or quotient for some i , then there exists a non-generic representation π' of G_n such that $\text{Hom}_{G_n}(\pi, \pi') \neq 0$. The statement still holds if we replace $\pi^{(i)}$ by ${}^{(i)}\pi$.*

Proof By Lemma 2.4, it suffices just to consider that $\pi^{(i)}$ has a non-generic irreducible quotient, say λ . Now let

$$\pi' = (\nu^{1/2}\lambda) \times \tau,$$

where τ is a cuspidal representation such that τ is not an unramified twist of a cuspidal representation appearing in any segment in \mathfrak{m} . Here \mathfrak{m} is a multisegment with $\pi \cong \langle \mathfrak{m} \rangle$. Now

$$\text{Ext}_{G_{n+1-j}}^k(\pi^{[j]}, {}^{(j-1)}\pi') = 0$$

for $j < n(\tau)$ and any k since τ is in $\text{csupp}({}^{(j-1)}\pi')$ whenever it is nonzero while $\tau \notin \text{csupp}(\pi^{[j]})$. Moreover, ${}^{(n(\tau)-1)}\pi'$ has a simple quotient isomorphic to $\nu^{1/2}\lambda$. This checks the Hom and Ext conditions in Lemma 2.1 and hence proves the lemma. The proof for ${}^{(i)}\pi$ is almost identical with switching left and right derivatives in suitable places. □

Theorem 3.3 *Let $\pi \in \text{Irr}(G_{n+1})$. Then the following conditions are equivalent:*

- (1) $\pi|_{G_n}$ is projective
- (2) π is generic and any irreducible quotient of $\pi|_{G_n}$ is generic.

Proof For (2) implying (1), it is proved in [15]. We now consider π is projective. All higher Exts vanish and so $\text{EP}(\pi, \pi') = \dim \text{Hom}_{G_n}(\pi, \pi')$ for any irreducible π' of G_n . If π' is an irreducible quotient of π , then $\text{EP}(\pi, \pi') \neq 0$ and hence π is generic by Theorem 3.1. But Theorem 3.1 also implies π' is generic. This proves (1) implying (2). □

3.2 Classification

Definition 3.4 We say that an irreducible representation π of G_{n+1} is *relatively projective* if either one of the following conditions holds:

- (i) π is essentially square-integrable;
- (ii) π is isomorphic to $\pi_1 \times \pi_2$ for some irreducible cuspidal representations π_1, π_2 of $G_{(n+1)/2}$ with $\pi_1 \not\cong \nu^{\pm 1}\pi_2$.

In particular, a relatively-projective representation is generic. The condition $\pi_1 \not\cong \nu^{\pm 1}\pi_2$ is in fact automatic from π being irreducible.

We can formulate the conditions (i) and (ii) combinatorially as follows. Let $\pi \cong \text{St}(\mathfrak{m})$ for a multisegment $\mathfrak{m} = \{\Delta_1, \dots, \Delta_r\}$. Then (i) is equivalent to $r = 1$; and (ii) is equivalent to that $r = 2$, and Δ_1 and Δ_2 are not linked, and the relative lengths of Δ_1 and Δ_2 are both 1.

Lemma 3.5 *Let $\pi \in \text{Irr}(G_{n+1})$ be not relatively projective. Then there exists an irreducible non-generic representation π' of G_n such that $\text{Hom}_{G_n}(\pi, \pi') \neq 0$.*

Proof It suffices to construct an irreducible non-generic representation π' satisfying the Hom and Ext properties in Lemma 2.1.

Let $\mathfrak{m} = \{\Delta_1, \dots, \Delta_r\}$ be a multisegment such that $\pi \cong \text{St}(\mathfrak{m})$. We divide into few cases.

Case 1 $r \geq 3$; or when $r = 2$, each segment has relative length at least 2; or when $r = 2$, $\Delta_1 \cap \Delta_2 = \emptyset$. We choose a segment Δ' in \mathfrak{m} with the shortest absolute length. Now we choose a maximal segment Δ in \mathfrak{m} with the property that $\Delta' \subset \Delta$. By genericity, $v^{-1}a(\Delta) \notin \Delta_k$ for any $\Delta_k \in \mathfrak{m}$. Let

$$\mathfrak{m}' = \{v^{1/2}\Delta, [v^{-1/2}a(\Delta)], [\tau]\},$$

where τ is a cuspidal representation so that $\text{St}(\mathfrak{m}')$ is a representation of G_n and τ is not an unramified twist of any cuspidal representation appearing in a segment of \mathfrak{m} . To make sense of the construction, it needs the choices and the assumptions on this case. Let $k = n(a(\Delta)), l = n(\tau)$. Let

$$\pi' = \text{St}(\mathfrak{m}').$$

Now as for $i < k + l + 1$, either $v^{-1/2}a(\Delta)$ or τ appears in the cuspidal support of ${}^{(i-1)}\pi$, but not in that of $v^{1/2}\pi^{(i)}$,

$$\text{Ext}_{G_{n+1-i}}^j(v^{1/2} \cdot \pi^{(i)}, {}^{(i-1)}\pi') = 0$$

and all j . By Corollary 2.6, $v^{1/2} \cdot \pi^{(k+l+1)}$ has a simple generic quotient isomorphic to $v^{1/2}\text{St}(-\Delta)$. On the other hand, a submodule structure of ${}^{(k+l)}\pi'$ can be computed as follows:

$$0 \neq \text{Hom}_{G_n}(\lambda(\mathfrak{m}'), \text{St}(\mathfrak{m}')) \tag{3.9}$$

$$\cong \text{Hom}_{G_{n-k-l} \times G_{k+l}}(\text{St}(v^{1/2} \cdot -\Delta) \boxtimes (v^{-1/2}a(\Delta) \times \tau), \text{St}(\mathfrak{m}')_{N_{k+l}^-}), \tag{3.10}$$

Here the non-zerosness comes from the fact that $\text{St}(\mathfrak{m}')$ is the unique quotient of $\lambda(\mathfrak{m}')$, and the isomorphism follows from Frobenius reciprocity. Since taking the derivative is an exact functor, we have that $\text{St}(v^{1/2} \cdot -\Delta)$ is a subrepresentation of ${}^{(k+l)}\text{St}(\mathfrak{m}')$ (see Sect. 2.3). Thus we have

$$\text{Hom}_{G_{n-k-l}}(\pi^{[k+l+1]}, {}^{(k+l)}\pi') \neq 0.$$

Case 2 $r = 2$ with $\Delta_1 \cap \Delta_2 \neq \emptyset$ and one segment having relative length 1 (and not both having relative length 1 by the definition of relatively-projective type). By switching the labeling on segments if necessary, we assume that $\Delta_1 \subset \Delta_2$. Let p and let l be the absolute and relative length of Δ_2 respectively. Let

$$\mathfrak{m}' = \{[v^{1/2}a(\Delta_1)], [v^{3/2-l}a(\Delta_1)], v^{-1/2}a(\Delta_1)], [\tau]\}, \quad \pi' = \text{St}(\mathfrak{m}'),$$

where τ is a cuspidal representation of G_k (here k is possibly zero) so that $\text{St}(\mathfrak{m}')$ is a G_n -representation. Note that π' is non-generic. By Corollary 2.6 and geometric lemma, a simple quotient $v^{1/2} \cdot \pi^{(p)}$ is isomorphic to $v^{1/2}a(\Delta_1)$. Similar computation as in (3.9) gives that a simple module of ${}^{(p-1)}\pi'$ is isomorphic to $v^{1/2}a(\Delta_1)$. This implies the non-vanishing Hom between those two G_{n+1-p} -representations.

We now prove the vanishing Ext-groups in order to apply Lemma 2.1. Now applying the Bernstein–Zelevinsky derivatives ($j = 1, \dots, p - 1$), unless $a(\Delta_1) \cong b(\Delta_2)$, we have that $v^{3/2-l}a(\Delta_1)$ is a cuspidal support for ${}^{(j-1)}\pi'$ whenever ${}^{(j-1)}\pi'$ is nonzero and is not a cuspidal support for $v^{1/2}\pi^{(j)}$. It remains to consider $a(\Delta_1) \cong b(\Delta_2)$. We can similarly consider the cuspidal support for $v^{-1/2}a(\Delta_1)$ and $v^{1/2}a(\Delta_1)$ to make conclusion. \square

Lemma 3.6 *Let $\pi \in \text{Irr}(G_{n+1})$. If π is (generic) relatively projective, then $\pi|_{G_n}$ is projective.*

Proof When π is essentially square-integrable, it is proved in [15]. We now assume that π is in the case (2) of Definition 3.4. It is equivalent to prove the condition (2) in Theorem 3.3. Let $\pi' \in \text{Irr}(G_n)$ with $\text{Hom}_{G_n}(\pi, \pi') \neq 0$. We have to show that π' is generic. Note that the only non-zero derivative of $\pi^{(i)}$ can occur when $i = n + 1$ and $\frac{n+1}{2}$.

Case 1 $\text{Hom}_{G_{(n+1)/2}}(\pi^{[(n+1)/2]}, {}^{((n-1)/2)}\pi') \neq 0$. For (1), let

$$\pi = \rho_1 \times \rho_2$$

for some cuspidal representations ρ_1, ρ_2 of $G_{(n+1)/2}$ with $\rho_1 \not\cong v^{\pm 1}\rho_2$, and

$$\mathfrak{m}' = \{\Delta'_1, \dots, \Delta'_s\} \quad \text{for } \pi' \cong \text{St}(\mathfrak{m}').$$

By a simple count on dimensions, we must have $\Delta'_k \cong v^{1/2}\rho_1$ or $\cong v^{1/2}\rho_2$ for some k . Using dimensions again, we have for $l \neq k$, $[\rho_1]$ and $[\rho_2]$ are unlinked to Δ'_l and so

$$\pi' \cong (v^{1/2} \cdot \rho_r) \times \text{St}(\mathfrak{m}' \setminus \{\Delta'_k\}),$$

for $r = 1$ or 2 . Then

$${}^{((n-1)/2)}\pi' \cong v^{1/2} \cdot \rho_r,$$

which implies

$${}^{((n-1)/2)}\text{St}(\mathfrak{m}' \setminus \{\Delta'_k\}) \neq 0.$$

Thus $\text{St}(\mathfrak{m}' \setminus \{\Delta'_k\})$ is generic and so is π' .

Case 2 $\text{Hom}_{G_{(n+1)/2}}(\pi^{[(n+1)/2]}, {}^{(n-1)/2}\pi') = 0$. We must have

$$\text{Hom}_{G_{n+1-i}}(\pi^{[n+1]}, {}^{(n)}\pi') \neq 0$$

and so ${}^{(n)}\pi' \neq 0$. Hence π' is generic. □

We now achieve the classification of irreducible representations which are projective when restricted from G_{n+1} to G_n :

Theorem 3.7 *Let $\pi \in \text{Irr}(G_{n+1})$. Then $\pi|_{G_n}$ is projective if and only if π is relatively projective in Definition 3.4.*

Proof The if direction is proved in Lemma 3.6. The only if direction follows from Lemma 3.5 and Theorem 3.1. □

One advantage for such classification is that those restricted representations admit a more explicit realization as shown in [15]:

Theorem 3.8 *Let $\pi, \pi' \in \text{Irr}(G_{n+1})$. If π and π' are relatively projective, then $\pi|_{G_n} \cong \pi'|_{G_n}$. In particular, $\pi|_{G_n}$ is isomorphic to the Gelfand–Graev representation $\text{ind}_{U_n}^{G_n} \psi_n$.*

Proof This follows from that π and π' are projective in $\text{Alg}(G_n)$ and [15]. □

4 Gelfand–Graev representations and affine Hecke algebras

Several insights come from the affine Hecke algebra realization of Gelfand–Graev representations. We shall first recall those results.

4.1 Affine Hecke algebras

Definition 4.1 The affine Hecke algebra $\mathcal{H}_l(q)$ of type A is an associative algebra over \mathbb{C} generated by $\theta_1, \dots, \theta_l$ and T_w ($w \in S_l$) satisfying the relations:

- (1) $\theta_i \theta_j = \theta_j \theta_i$;
- (2) $T_{s_k} \theta_k - \theta_{k+1} T_{s_k} = (q - 1)\theta_k$, where q is a certain prime power and s_k is the transposition between the numbers k and $k + 1$;
- (3) $T_{s_k} \theta_i = \theta_i T_{s_k}$, where $i \neq k, k + 1$
- (4) $(T_{s_k} - q)(T_{s_k} + 1) = 0$;
- (5) $T_{s_k} T_{s_{k+1}} T_{s_k} = T_{s_{k+1}} T_{s_k} T_{s_{k+1}}$.

Let $\mathcal{A}_l(q)$ be the (commutative) subalgebra generated by $\theta_1, \dots, \theta_l$. Let $\mathcal{H}_{W,l}(q)$ be the subalgebra generated by $T_{s_1}, \dots, T_{s_{l-1}}$. Let sgn be the 1-dimensional $\mathcal{H}_{W,l}(q)$ -module characterized by T_{s_k} acting by -1 .

It is known from [25, Proposition 3.11] that the center \mathcal{Z}_l of $\mathcal{H}_l(q)$ has a basis $\left\{ z_M = \sum_{w \in S_l} \theta_1^{i_{w(1)}} \dots \theta_l^{i_{w(l)}} \right\}_M$, where $M = (i_1, \dots, i_l)$ runs for all l -tuples in \mathbb{Z}^l/S_l .

Bernstein decomposition asserts that

$$\mathfrak{R}(G_n) \cong \prod_{\mathfrak{s} \in \mathfrak{B}(G_n)} \mathfrak{R}_{\mathfrak{s}}(G_n),$$

where $\mathfrak{R}(G_n)$ is the category of smooth G_n -representations, $\mathfrak{B}(G_n)$ is the set of inertial equivalence classes of G_n and $\mathfrak{R}_{\mathfrak{s}}(G_n)$ is the full subcategory of $\mathfrak{R}(G_n)$ associated to \mathfrak{s} (see [10]). For a smooth representation π of G_n , define $\pi_{\mathfrak{s}}$ to be the projection of π to the component $\mathfrak{R}_{\mathfrak{s}}(G_n)$.

For each $\mathfrak{s} \in \mathfrak{B}(G_n)$, [9] and [10] associate with a compact group $K_{\mathfrak{s}}$ and a finite-dimensional representation τ of $K_{\mathfrak{s}}$, such that the convolution algebra

$$\mathcal{H}(K_{\mathfrak{s}}, \tau) := \{ f : G_n \rightarrow \text{End}(\tau^{\vee}) : f(k_1 g k_2) = \tau^{\vee}(k_1) \circ f(g) \circ \tau^{\vee}(k_2) \text{ for } k_1, k_2 \in K_{\mathfrak{s}} \}$$

is isomorphic to the product $\mathcal{H}_{n_1}(q_1) \otimes \dots \otimes \mathcal{H}_{n_r}(q_r)$ of affine Hecke algebra of type A , denoted by $\mathcal{H}_{\mathfrak{s}}$. For a smooth representation π of G_n , the algebra $\mathcal{H}(K_{\mathfrak{s}}, \tau)$ acts naturally on the space $\text{Hom}_{K_{\mathfrak{s}}}(\tau, \pi) \cong (\tau^{\vee} \otimes \pi)^{K_{\mathfrak{s}}}$. This defines an equivalence of categories:

$$\mathfrak{R}_{\mathfrak{s}}(G_n) \cong \text{category of } \mathcal{H}_{\mathfrak{s}}\text{-modules} . \tag{4.11}$$

By abuse notation, we shall identify $\text{Hom}_{K_{\mathfrak{s}}}(\tau, \pi)$ with $\pi_{\mathfrak{s}}$ under (4.11) and consider $\pi_{\mathfrak{s}}$ as $\mathcal{H}_{\mathfrak{s}}$ -module.

Let $\mathcal{A}_{\mathfrak{s}} = \mathcal{A}_{n_1}(q_1) \otimes \dots \otimes \mathcal{A}_{n_r}(q_r)$. Let $\mathcal{H}_{W, \mathfrak{s}} = \mathcal{H}_{W, n_1}(q_1) \otimes \dots \otimes \mathcal{H}_{W, n_r}(q_r)$. Let

$$\text{sgn}_{\mathfrak{s}} = \text{sgn} \boxtimes \dots \boxtimes \text{sgn}$$

as an $\mathcal{H}_{W, \mathfrak{s}}$ -module. Note that in [13], we proved when \mathfrak{s} is a simply type, but the generalization to all types follows from [10] and a simple generalization of [13, Theorem 2.1]. The center of $\mathcal{H}_{\mathfrak{s}}$ is equal to $\mathcal{Z}_1 \otimes \dots \otimes \mathcal{Z}_r$, where each \mathcal{Z}_k is the center of $\mathcal{H}_{n_k}(q_k)$.

Recall that $\Pi_n = \text{ind}_{U_n}^{G_n} \psi_n$. We may simply write Π for Π_n if there is no confusion.

Theorem 4.2 [13] *For any $\mathfrak{s} \in \mathfrak{B}(G_n)$, the Bernstein component of the Gelfand–Graev representation $\Pi_{\mathfrak{s}}$ is isomorphic to $\mathcal{H}_{\mathfrak{s}} \otimes_{\mathcal{H}_{W, \mathfrak{s}}} \text{sgn}_{\mathfrak{s}}$.*

Using Theorem 4.2, we have that $\Pi_{\mathfrak{s}}$ is isomorphic to $\mathcal{A}_{\mathfrak{s}}$, as $\mathcal{A}_{\mathfrak{s}}$ -module. This observation has the following consequence:

Lemma 4.3 *For any $\mathfrak{s} \in \mathfrak{B}(G_n)$, any two non-zero submodules of $\Pi_{\mathfrak{s}}$ have non-zero intersections. In particular, $\Pi_{\mathfrak{s}}$ is indecomposable.*

We remark that for general quas-split group G , the indecomposability of $\Pi_{\mathfrak{s}}$ can be deduced from results of [8], which show that endomorphism algebra of $\Pi_{\mathfrak{s}}$ is isomorphic to the Bernstein center at the corresponding (indecomposable) Bernstein block $\mathfrak{R}_{\mathfrak{s}}(G)$.

We shall also need:

Lemma 4.4 *Fix $\mathfrak{s} \in \mathfrak{B}(G_n)$. Let \mathcal{J} be a maximal ideal in $\mathcal{Z}_{\mathfrak{s}}$ that annihilates an irreducible module in $\mathfrak{R}_{\mathfrak{s}}(G_n)$. Let π be a non-zero submodule of $\Pi_{\mathfrak{s}}$. Let $\widehat{\pi}$ be the \mathcal{J} -adic completion of π . Then $\widehat{\pi} \neq 0$.*

Proof Note that $\mathcal{A}_{\mathfrak{s}}$ is an integral domain and is finitely-generated over $\mathcal{Z}_{\mathfrak{s}}$. Hence, by Krull’s intersection theorem, $\bigcap_k \mathcal{J}^k \mathcal{A}_{\mathfrak{s}} = 0$. Thus $\bigcap_k \mathcal{J}^k \pi = 0$. This implies that $\pi / \mathcal{J}^k \pi \neq 0$ for some k . Hence $\widehat{\pi} \neq 0$. \square

4.2 Jacquet functors on Gelfand–Graev representations

Lemma 4.5 *Let $P = LN$ be the parabolic subgroup containing upper triangular matrices and block-diagonal matrices $\text{diag}(g_1, \dots, g_r)$ with $g_k \in G_{i_k}$, where $i_1 + \dots + i_r = n$. Then $(\Pi_n)_N \cong \Pi_{i_1} \boxtimes \dots \boxtimes \Pi_{i_r}$.*

Proof Let w be a permutation matrix in G_n . Then $w(N) \cap U_n$ contains a unipotent subgroup $\{I_n + tu_{k,k+1} : t \in F\}$ for some k if and only if $w(N) \not\subset U_n^-$. Here $u_{k,k+1}$ is a matrix with $(k, k + 1)$ -entry 1 and other entries 0. For any such w , it gives that PwB is the same unique open orbit in G_n . Now the geometric lemma in [7, Theorem 5.2] gives the lemma. \square

5 Submodule structure of Bernstein–Zelevinsky layers

5.1 Inertial equivalence classes

We give more discussions on inertial equivalence classes, relating the parabolic induction. See, for example, [10]. An inertial equivalence class \mathfrak{s} of G_n can be represented by a pair $[G_{m_1} \times \dots \times G_{m_r}, \rho_1 \boxtimes \dots \boxtimes \rho_r]$, where $m_1 + \dots + m_r = n$ and each ρ_k is a cuspidal G_{m_k} -representation. Two pairs

$$[G_{m_1} \times \dots \times G_{m_r}, \rho_1 \boxtimes \dots \boxtimes \rho_r], [G_{m'_1} \times \dots \times G_{m'_s}, \rho'_1 \boxtimes \dots \boxtimes \rho'_s]$$

represent the same inertial equivalence class if and only if $r = s$ and there exists a permutation $\sigma \in S_r$ such that

$$G_{m_1} = G_{m'_{\sigma(1)}}, \dots, G_{m_r} = G_{m'_{\sigma(r)}}$$

and

$$\rho_1 \cong \chi_1 \otimes \rho'_{\sigma(1)}, \dots, \rho_r \cong \chi_r \otimes \rho'_{\sigma(r)}$$

for some unramified character χ_k of G_{m_k} ($k = 1, \dots, r$).

Let

$$\begin{aligned} \mathfrak{s}_1 &= [G_{m_1} \times \dots \times G_{m_r}, \rho_1 \boxtimes \dots \boxtimes \rho_r] \in \mathfrak{B}(G_{n_1}), \\ \mathfrak{s}_2 &= [G_{m'_1} \times \dots \times G_{m'_s}, \rho'_1 \boxtimes \dots \boxtimes \rho'_s] \in \mathfrak{B}(G_{n_2}). \end{aligned}$$

Then $\pi_1 \times \pi_2$ lies in $\mathfrak{R}_{\mathfrak{s}}(G_{n_1+n_2})$, where

$$\mathfrak{s} = [G_{m_1} \times \dots \times G_{m_r} \times G_{m'_1} \times \dots \times G_{m'_s}, \rho_1 \boxtimes \dots \boxtimes \rho_r \boxtimes \rho'_1 \boxtimes \dots \boxtimes \rho'_s]$$

From this, one deduces the following lemma:

Lemma 5.1 *Let $\pi_1 \in \mathfrak{R}_{\mathfrak{s}}(G_{n_1})$ for some $\mathfrak{s} \in \mathfrak{B}(G_{n_1})$ and let $\pi_2 \in \text{Alg}(G_{n_2})$. Fix $\mathfrak{t} \in \mathfrak{B}(G_{n_1+n_2})$ with $(\pi_1 \times \pi_2)_{\mathfrak{t}} \neq 0$. There exists a unique $\mathfrak{s}' \in \mathfrak{B}(G_{n_1+n_2})$ such that $\pi_1 \times (\pi_2)_{\mathfrak{s}'} \cong (\pi_1 \times \pi_2)_{\mathfrak{t}}$.*

5.2 Bernstein center

The Bernstein center of a category \mathfrak{R} is defined as the endomorphism ring of the identity functor in \mathfrak{R} . Denote by \mathfrak{Z}_n the Bernstein center of $\mathfrak{R}(G_n)$. For $\mathfrak{s} \in \mathfrak{B}(G_n)$, denote by $\mathfrak{Z}_{\mathfrak{s}}$ the Bernstein center of $\mathfrak{R}_{\mathfrak{s}}(G_n)$, which is a finitely-generated commutative algebra [3, Theorem 2.13]. Explicitly, for

$$\mathfrak{s} = [G_{n_1} \times \dots \times G_{n_k}, \rho_1 \boxtimes \dots \boxtimes \rho_k] \in \mathfrak{B}(G_n),$$

set, the Bernstein variety, to be

$$\mathfrak{X}_{\mathfrak{s}} = (\mathbb{C}^{\times})^k / W_{\mathfrak{s}},$$

where each copy of \mathbb{C}^{\times} comes from the group of unramified characters of G_{n_p} and $W_{\mathfrak{s}}$ is the subgroup of $N_G(M)/M$, where $M = G_{n_1} \times \dots \times G_{n_k}$, which stabilizes $\rho_1 \boxtimes \dots \boxtimes \rho_k$, and $W_{\mathfrak{s}}$ permutes the factors in $(\mathbb{C}^{\times})^k$ via its action on M . Here $N_G(M)$ is the normalizer of M in G . According to [3],

$$\mathfrak{Z}_{\mathfrak{s}} \cong \mathbb{C}[\mathfrak{X}_{\mathfrak{s}}],$$

and we also have

$$\mathfrak{Z}_n \cong \prod_{\mathfrak{s} \in \mathfrak{B}(G_n)} \mathfrak{Z}_{\mathfrak{s}}.$$

It follows from [3, Corollaire 3.4] that for any maximal ideal \mathcal{J} of \mathfrak{Z}_n and any finitely-generated π of G_n , $\pi/\mathcal{J}(\pi)$ is admissible. For more properties of the Bernstein center, see, e.g. [3, 5, 8, 20].

For a maximal ideal \mathcal{J} in \mathfrak{Z}_n and $\pi \in \text{Alg}(G_n)$, define $\widehat{\pi}$ to be the \mathcal{J} -adic completion of π i.e. the inverse limit

$$\widehat{\pi} = \lim_i \pi/(\mathcal{J}^i \pi),$$

which has a natural G -module structure from π . The notion $\widehat{\pi}$ depends on \mathcal{J} , and it should be clear from the context. Note that \mathcal{J} comes from some maximal ideal in $\mathfrak{Z}_{\mathfrak{s}}$ for some unique $\mathfrak{s} \in \mathfrak{B}(G_n)$ and this gives that

$$\widehat{\pi} \in \mathfrak{R}_{\mathfrak{s}}(G_n)$$

Lemma 5.2 *Let $\omega \in \text{Alg}(G_{n_1})$ be admissible with $n_1 \neq 0$. Fix $\tau \in \text{Irr}(G_{n_1+n_2})$ such that $\text{csupp}(\omega) \cap \text{csupp}(\tau) = \emptyset$. Let $\pi = \omega \times \Pi_{n_2}$ and $\pi' = \Pi_{n_2} \times \omega$. Let \mathcal{J} be the maximal ideal in $\mathfrak{Z}_{n_1+n_2}$ which annihilates τ . Then*

- (1) $\widehat{\pi} = 0$; and
- (2) $\widehat{\pi}' = 0$.

Proof We only prove (1) and a proof for (2) is similar. Each Bernstein component of Π is finitely generated [8]. Thus $\pi/\mathcal{J}(\pi)$ is admissible [3] and is annihilated by \mathcal{J} . If $\pi/\mathcal{J}(\pi)$ is non-zero, then π has a subquotient with irreducible composition factors isomorphic to τ' with $\text{csupp}(\tau') = \text{csupp}(\tau)$ [3, Proposition 2.11, Theorem 2.13]. However, using Jacquet functor, one shows that the cuspidal support of any irreducible subquotients of $\omega \times \Pi_{n_2}$ has non-zero intersection with $\text{csupp}(\omega)$. This gives a contradiction. Hence $\pi/\mathcal{J}(\pi) = 0$ and so $\widehat{\pi} = 0$. □

5.3 Intersection properties of Bernstein–Zelevinsky layers

The main tool of this section is the exactness of Jacquet functor.

Lemma 5.3 *For $i \neq j$, let ω, τ be an admissible G_{n-i} -representation and G_{n-j} -representation respectively. There is no non-zero isomorphic submodule of $\text{ind}_{R_{n-i}}^{G_n} \omega \boxtimes \psi_i$ and $\text{ind}_{R_{n-j}}^{G_n} \tau \boxtimes \psi_j$.*

Proof We only prove for $i > j$ and the case for $j < i$ is similar. Suppose there exists a non-zero isomorphic submodule λ of $\omega \times \Pi_i$ and $\Pi_j \times \tau$. By Frobenius reciprocity on $\lambda \hookrightarrow \omega \times \Pi_i$,

$$\lambda_{N_i} \neq 0. \tag{5.12}$$

We have that $(\text{ind}_{R_{n-i}}^{G_n} \omega \otimes \psi_i)_{N_i}$ admits a filtration, from geometric lemma [7, Theorem 5.2], that successive quotients take the form

$$(\omega' \times \Pi_k) \boxtimes (\omega'' \times \Pi_{i-k}). \tag{5.13}$$

Similarly, $(\text{ind}_{R_{n-i}}^{G_n} \tau \boxtimes \psi_j)_{N_i}$ admits a filtration that successive quotients take the form

$$(\Pi_l \times \tau') \boxtimes (\Pi_{j-l} \times \tau''), \tag{5.14}$$

where τ' is an admissible G_{n-i-l} -representation and τ'' is an admissible G_{i-j+l} representation. Since Jacquet functor is exact, it reduces, by Lemma 10.3 and (5.12), to see that there is no submodule for

$$(\Pi_l \times \tau') \boxtimes (\Pi_{j-l} \times \tau''), \quad \text{and} \quad (\omega' \times \Pi_k) \boxtimes (\omega'' \times \Pi_{i-k}).$$

for all k, l .

In the case that $k = 0$ and $l = j$; or $k = n - i$ and $l = 0$, a certain element in the Bernstein center annihilates ω'' or ω' , but not on Π_i or Π_{n-i} respectively (by [8]). Now to prove other cases, we notice that either $k > l$ or $i - k > j - l$ (which happens when $l \geq k$). In the first case, we apply $N_k \subset G_{n-i}$ on the first factor, and in the second case, apply N_{i-k} on the second factor. Then repeat the process and the process terminates as each process the value on i decreases. \square

A special case of Lemma 5.3 is that $i = 0$ and $j = n \geq 1$. In such case, Lemma 5.3 says that the Gelfand–Graev representation Π_n does not admit an irreducible submodule. We also remark that the last fact also holds for a more general quasi-split connected reductive group with a non-compact center,³ which can be deduced from a combination of Ext-duality (see [11, 26, 29]) and the projectivity of a Gelfand–Graev representation (see [13]).

For convenience, an admissible representation π of G_n is said to be uniform if all its composition factors have the same cuspidal support. We shall set $\text{csupp}(\pi) = \text{csupp}(\pi')$ for any irreducible $\pi' \in \text{JH}(\pi)$. It is well-known that any admissible representation can be written as direct sum of uniform representations. For a uniform representation π of G_n , we associate the maximal ideal \mathcal{J}_π in \mathfrak{Z}_n such that some power of \mathcal{J}_π annihilates π .

Lemma 5.4 *Let $\omega \in \text{Alg}(G_{n-i})$ be uniform. Let $\tau \in \text{Irr}(G_i)$ such that $\text{csupp}(\tau) \cap \text{csupp}(\omega) = \emptyset$. Set $\mathcal{I} = \mathcal{J}_\omega \otimes 1 + 1 \otimes \mathcal{J}_\tau$ in $\mathfrak{Z}_{n-i} \otimes \mathfrak{Z}_i$. Then, as $G_{n-i} \times G_i$ -representations,*

$$(\widehat{\omega \times \Pi_i})_{N_i} = \omega \boxtimes \widehat{\Pi_i}.$$

³ The author would like to thank the referee for pointing out that.

(Here $(\widehat{\omega \times \Pi_i})_{N_i}$ is the \mathcal{I} -adic completion for the $G_{n-i} \times G_i$ -representation $(\omega \times \Pi_i)_{N_i}$. The $\widehat{\Pi_i}$ is the \mathcal{J}_τ -adic completion of Π_i .)

Proof We have observed that $(\omega \times \Pi_i)_{N_i}$ admits a filtration with successive quotients in (5.13). We now apply the \mathcal{I} -adic completion on each subquotients in the form of (5.13). By using Lemma 5.2 and the condition that $\text{csupp}(\tau) \cap \text{csupp}(\omega) = \emptyset$, the only successive quotient which is non-zero after taking the \mathcal{I} -adic completion is the factor $\omega \boxtimes \Pi_i$. Since \mathcal{I} -adic completion is exact on finitely-generated $G_{n-i} \times G_i$ -representations, we obtain that

$$(\widehat{\omega \times \Pi_i})_{N_i} \cong \omega \boxtimes \widehat{\Pi_i}.$$

We make two remarks on the above proof:

- In order to get finitely-generated modules, one has to consider each Bernstein component of $\omega \times \Pi_i$ and those successive quotients.
- To show the exactness on the \mathcal{I} -adic completion, one can pass to the Hecke algebra so that a finitely-generated $G_{n-i} \times G_i$ -representation will give rise a corresponding Hecke algebra module which is finitely-generated over $\mathfrak{Z}_{n-i} \otimes \mathfrak{Z}_i$. □

Lemma 5.5 *Keep using the notation in the previous lemma. We also have that*

$$(\widehat{\Pi_i \times \omega})_{N_i} \cong \omega \boxtimes \widehat{\Pi_i}.$$

Proof The proof is the same as that of Lemma 5.4 except that we use the filtration (5.14) instead of (5.13). □

Proposition 5.6 *Let $\omega_1, \omega_2 \in \text{Alg}(G_{n-i})$ be admissible and non-zero. If $\text{ind}_{R_{n-i}}^{G_n} \omega_1 \boxtimes \psi_i$ and $\text{ind}_{R_{n-i}}^{G_n} \omega_2 \boxtimes \psi_i$ have isomorphic non-zero submodules, then ω_1 and ω_2 have isomorphic non-zero submodules.*

Proof Let $0 \neq \pi \in \text{Alg}(G_n)$ such that

$$\pi \xhookrightarrow{\iota_1} \omega_1 \times \Pi_i, \quad \text{and} \quad \pi \xhookrightarrow{\iota_2} \Pi_i \times \omega_2.$$

We now write ω_1 as direct sum of uniform representations $\lambda_1, \dots, \lambda_k$. Then $\omega_1 \times \Pi_i$ admits a filtration whose successive quotients of the form $\lambda_r \times \Pi_i$. By Lemma 10.2, we have that

$$\pi' \hookrightarrow \lambda_r \times \Pi_i \tag{5.15}$$

for some r and some non-zero submodule π' of π . It suffices to show that λ_r and ω_2 share an isomorphic irreducible submodule. To this end, now set $\mathcal{I} = \mathcal{J}_{\lambda_r} \otimes 1 + 1 \otimes \mathcal{J}_\tau$, where $\tau \in \text{Irr}(G_i)$ satisfies $\text{csupp}(\lambda_r) \cap \text{csupp}(\tau) = \emptyset$.

Frobenius reciprocity gives a non-zero map from π'_{N_i} to $\lambda_r \boxtimes \Pi_i$ and so the image is a non-zero submodule of $\lambda_r \boxtimes \Pi_i$, which is still non-zero after taking \mathcal{I} -adic completion by Lemma 4.4. As taking \mathcal{I} -adic completion is exact, we have that $\widehat{\pi'_{N_i}} \neq 0$.

Now taking exact functors on $\pi' \hookrightarrow \Pi_i \times \omega_2$, we obtain:

$$0 \neq \widehat{\pi'_{N_i}} \hookrightarrow (\widehat{\Pi_i \times \omega_2})_{N_i} \tag{5.16}$$

Now we regard $\widehat{\pi'_{N_i}}$ as G_{n-i} -submodule via the embedding $g \mapsto \text{diag}(g, I_i)$. Using (5.15) and Lemma 5.4, any G_{n-i} -submodule of $\widehat{\pi'_{N_i}}$ is a submodule of λ_r . Similarly, using (5.16) and Lemma 5.5, any G_{n-i} -submodule of $\widehat{\pi'_{N_i}}$ is a submodule of ω_2 . This concludes that λ_r and ω_2 share isomorphic irreducible G_{n-i} -submodules. \square

5.4 Strong indecomposability

We first prove a preparation lemma.

Lemma 5.7 *Let π_1, π_2, π be in $\text{Alg}(G_n)$ such that $\pi_1 \hookrightarrow \pi$ and $\pi_2 \hookrightarrow \pi$. Let $N = N_i$ for some i . Then $(\pi_1 \cap \pi_2)_N \cong (\pi_1)_N \cap (\pi_2)_N$. Here the later intersection is taken in π_N .*

Proof We have the natural projection $p : \pi_1 \cap \pi_2 \rightarrow \pi_N$ as linear spaces. Since the image of the projection lies in both $(\pi_1)_N$ and $(\pi_2)_N$, the projection factors through the embedding $(\pi_1)_N \cap (\pi_2)_N$ to π_N . Now taking the Jacquet functor on p gives an isomorphism from $(\pi_1 \cap \pi_2)_N$ onto $(\pi_1 \cap \pi_2)_N \subset \pi_N$. Thus the map from $(\pi_1)_N \cap (\pi_2)_N$ to $(\pi_1 \cap \pi_2)_N$ is also surjective. \square

We shall prove a weak version on the strong indecomposability of the Bernstein–Zelevinsky induction. The stronger version of Lemma 5.8 is to only assume the embedding on either left or right Bernstein–Zelevinsky induction. Proving such statement requires more work and we will not do it here. Some special cases can be more easily achieved by using Theorem 6.1 below. In Sect. 8, we shall prove a variation, which says that the Bernstein–Zelevinsky induction preserves indecomposability (but not strong indecomposability).

Lemma 5.8 *Let $\omega \in \text{Irr}(G_{n-i})$. Let $\mathfrak{s} \in \mathfrak{B}(G_n)$. Suppose there exists non-zero $\pi_1, \pi_2 \in \text{Alg}(G_n)$ with the following embeddings:*

$$\pi_1 \hookrightarrow (\omega \times \Pi_i)_{\mathfrak{s}}, \quad \pi_2 \hookrightarrow (\omega \times \Pi_i)_{\mathfrak{s}}$$

and

$$\pi_1 \hookrightarrow (\Pi_i \times \omega)_{\mathfrak{s}}, \quad \pi_2 \hookrightarrow (\Pi_i \times \omega)_{\mathfrak{s}}.$$

Then the images of the embeddings of π_1 and π_2 in $(\omega \times \Pi_i)_\mathfrak{s}$ have non-zero intersections. Similarly, the images of the embeddings of π_1 and π_2 in $(\Pi_i \times \omega)_\mathfrak{s}$ have non-zero intersections.

Proof We shall prove the latter statement. The former statement can be proved similarly. We shall identify π_k ($k = 1, 2$) with the image of the embedding of π_k in $(\Pi_i \times \omega)_\mathfrak{s}$.

Let $\tau \in \text{Irr}(G_i)$ with $\text{csupp}(\tau) \cap \text{csupp}(\omega) = \emptyset$. Set $\mathcal{I} = \mathcal{J}_\omega \otimes 1 + 1 \otimes \mathcal{J}_\tau$. Using a similar argument to the one in the proof of Proposition 5.6 (which uses Frobenius reciprocity and Lemma 4.4), we have that

$$\widehat{(\pi_1)_{N_i}} \neq 0 \tag{5.17}$$

(We recall that we need the embedding from π_1 to $\omega \times \Pi_i$ to prove the non-zero part.)

On the other hand, as observed in Proposition 5.6, among all the successive quotients in the filtration of $(\Pi_i \times \omega)_{N_i}$ of the form (5.15), the only one that does not vanish after taking \mathcal{I} -adic completion is $\omega \boxtimes \Pi_i$.

Moreover, the successive quotient $\omega \boxtimes \Pi_i$ lies in the bottom layer of $(\Pi_i \times \omega)_{N_i}$ [7, Theorem 5.2]. Thus, the non-vanishing (5.17) implies that $(\pi_1)_{N_i} \cap (\omega \boxtimes \Pi_i) \neq 0$. Moreover, the Bernstein component \mathfrak{s} uniquely determines a $\mathfrak{t} \in \mathfrak{B}(G_i)$ such that $(\pi_1)_{N_i} \cap (\omega \boxtimes (\Pi_i)_\mathfrak{t}) \neq 0$ (c.f. Lemma 5.1).

Similarly, we have that $(\pi_2)_{N_i} \cap (\omega \boxtimes (\Pi_i)_\mathfrak{t}) \neq 0$. Thus since $\omega \boxtimes (\Pi_i)_\mathfrak{t}$ is strongly indecomposable $G_{n-i} \times G_i$ -representation (by Lemma 4.3 and ω is irreducible), we have that

$$(\pi_1)_{N_i} \cap (\pi_2)_{N_i} \neq 0.$$

By Lemma 5.7, $(\pi_1 \cap \pi_2)_{N_i} \neq 0$ and so $\pi_1 \cap \pi_2 \neq 0$. □

6 Indecomposability of restricted representations

6.1 Indecomposability of restriction

We now prove our main result:

Theorem 6.1 *Let $\pi \in \text{Irr}(G_{n+1})$. Then for each $\mathfrak{s} \in \mathfrak{A}(G_n)$, $\pi_\mathfrak{s}$ is strongly indecomposable whenever it is nonzero i.e. for any two non-zero submodules τ, τ' of $\pi_\mathfrak{s}$, $\tau \cap \tau' \neq 0$.*

Proof There exists a Bernstein–Zelevinsky G_n -filtration on π with

$$\pi_n \subset \pi_{n-1} \subset \dots \subset \pi_1 \subset \pi_0 = \pi \tag{6.18}$$

such that

$$\pi_i/\pi_{i+1} \cong \text{ind}_{R_{n-i}}^{G_n} \pi^{[i+1]} \boxtimes \psi_i. \tag{6.19}$$

We also have a G_n -filtration on π with

$${}_n\pi \subset {}_{n-1}\pi \subset \dots \subset {}_1\pi \subset {}_0\pi = \pi \tag{6.20}$$

such that

$${}_i\pi/{}_{i+1}\pi \cong \text{ind}_{R_{n-i}^-}^{G_n} {}^{[i+1]}\pi \boxtimes \psi_i. \tag{6.21}$$

Let i^* be the level of π . For $\mathfrak{s} \in \mathfrak{B}(G_n)$ such that $\pi_{\mathfrak{s}} \neq 0$, we also have $(\pi_{i^*-1})_{\mathfrak{s}} \neq 0$ (see Sect. 6.2 for the detail). For notation simplicity, we set $\tau = (\pi_{i^*-1})_{\mathfrak{s}}$ and $\tau' = ({}_{i^*-1}\pi)_{\mathfrak{s}}$, both regarded as subspaces of π .

Let ω and γ be two non-zero submodules of $\pi_{\mathfrak{s}}$.

Claim $\omega \cap \tau \neq 0$.

Proof of the Claim Suppose not. Then, the natural projection gives an injection

$$\omega \hookrightarrow \pi_{\mathfrak{s}}/\tau \tag{6.22}$$

By Lemma 5.3, there is no isomorphic submodules between $\pi_{\mathfrak{s}}/\tau$ and τ' . This implies $\omega \cap \tau' = 0$. Hence, we also have an injection:

$$\omega \hookrightarrow \pi_{\mathfrak{s}}/\tau' \tag{6.23}$$

By (6.18), (6.20), (6.22), (6.23) and Lemma 10.3, there exists a G_n -representation which is isomorphic to submodules of

$$(\pi^{[j]} \times \Pi_{j-1})_{\mathfrak{s}}, \quad \text{and} \quad (\Pi_{k-1} \times {}^{[k]}\pi)_{\mathfrak{s}}$$

for some $j, k < i^*$. By Lemma 5.3 again, we must have $j = k$. However, Proposition 5.6 contradicts to the following Theorem 6.2 below (whose proof does not depend on this result). This proves the claim.

Since ω is an arbitrary submodule of π , we also have $\gamma \cap \tau \neq 0$. Now we refine our G_n -modules and set:

$$0 \neq \omega' = \omega \cap \tau \subset \pi_{\mathfrak{s}}, \quad 0 \neq \gamma' = \gamma \cap \tau \subset \pi_{\mathfrak{s}}.$$

Now using similar argument as above, we also have that

$$\omega' \cap \tau' \neq 0, \quad \gamma' \cap \tau' \neq 0.$$

Now we further refine our G_n -modules and set

$$\omega'' = \omega' \cap \tau' \subset \pi_{\mathfrak{s}}, \quad \gamma'' = \gamma' \cap \tau' \subset \pi_{\mathfrak{s}}.$$

Now ω'' and γ'' have desired embeddings as in Lemma 5.8. Hence Lemma 5.8 implies that $\omega'' \cap \gamma'' \neq 0$ and so we have $\omega \cap \gamma \neq 0$ as desired. \square

We remark that one can use Theorem 6.1 to show that for $\omega \in \text{Irr}(G_m)$ and for i large enough (e.g. $i \geq m$), $(\omega \times \Pi_i)_{\mathfrak{s}}$ (resp. $(\Pi_i \times \omega)_{\mathfrak{s}}$) are strongly indecomposable for any $\mathfrak{s} \in \mathfrak{B}(G_{m+i})$ (c.f. Lemma 5.8). This is done by realizing $\omega \times \Pi_i$ (resp. $\Pi_i \times \omega$) as the bottom layer of the Bernstein–Zelevinsky filtration of some irreducible module π of G_{m+i+1} and then embed the submodules of $\omega \times \Pi_i$ (resp. $\Pi_i \times \omega$) to π .

Theorem 6.2 *Let $\pi \in \text{Irr}(G_{n+1})$. If i is not the level of π , then $\pi^{[i]}$ and ${}^{[i]}\pi$ do not have an isomorphic irreducible quotient, and also do not have an isomorphic irreducible submodule whenever the two derivatives are not zero.*

The proof of Theorem 6.2 will be carried out in Sect. 7. Note that the converse of the above theorem is also true, which follows directly from the well-known highest derivative due to Zelevinsky [35, Theorem 8.1].

6.2 Non-zero Bernstein components

Let $\pi \in \text{Irr}(G_{n+1})$. In order to give an explicit parametrization of indecomposable components of $\pi|_{G_n}$, we also have to determine when $\pi_{\mathfrak{s}} \neq 0$ for $\mathfrak{s} \in \mathfrak{B}(G_n)$. Indeed this can be done as follows. Write $\pi \cong \langle \mathfrak{m} \rangle$ for a multisegment $\mathfrak{m} = \{\Delta_1, \dots, \Delta_k\}$. Let π' be the (right) highest derivative of π . Then we obtain a multiset

$$\text{csupp}(\pi') = (\rho_1, \dots, \rho_p).$$

This multiset determines a cuspidal $G_{k_1} \times \dots \times G_{k_p}$ -representation $\rho_1 \boxtimes \dots \boxtimes \rho_p$. Now we pick positive integers k_{p+1}, \dots, k_q such that $k_1 + \dots + k_q = n$, and pick cuspidal representations $\rho_{p+1}, \dots, \rho_q$ of $G_{k_{p+1}}, \dots, G_{k_q}$ respectively. Then for the inertial equivalence class

$$\mathfrak{s} = [G_{k_1} \times \dots \times G_{k_q}, \rho_1 \boxtimes \dots \boxtimes \rho_q],$$

we have that $\pi_{\mathfrak{s}} \neq 0$, which follows from that the bottom Bernstein–Zelevinsky layer $(\pi^{[i^*]} \times \Pi_{i^*-1})_{\mathfrak{s}} \subset \pi_{\mathfrak{s}}$ is non-zero. Here i^* is the level of π .

Indeed for any $\mathfrak{s} \in \mathfrak{B}(G_n)$ with $\pi_{\mathfrak{s}} \neq 0$, \mathfrak{s} arises in the above way. To see this, we need the following lemma:

Lemma 6.3 *Let $\pi \in \text{Irr}(G_{n+1})$. Let i^* be the level of π . Then, for any $i \leq i^*$ with $\pi^{[i]} \neq 0$ and for any $\omega \in \text{JH}(\pi^{[i]})$, $\text{csupp}(\pi^{[i^*]}) \subset \text{csupp}(\omega)$ (counting multiplicities).*

Proof Let $\mathfrak{m} \in \text{Mult}$ such that $\pi \cong \langle \mathfrak{m} \rangle$. Now by definition, we have that

$$\langle \mathfrak{m} \rangle \hookrightarrow \zeta(\mathfrak{m})$$

and so $\langle \mathfrak{m} \rangle^{[i]} \hookrightarrow \zeta(\mathfrak{m})^{[i]}$.

For any segment Δ , set $\Delta^{[0]} = \nu^{1/2} \Delta$ and set $\Delta^{[-1]} = \nu^{1/2} \Delta^-$. We also set $\mathfrak{m} = \{\Delta_1, \dots, \Delta_k\}$ and set

$$\mathcal{M} = \{ \{ \Delta_1^\#, \dots, \Delta_k^\# \} : \# = [0], [-1] \}.$$

Now the geometric lemma gives a filtration on $\zeta(\mathfrak{m})^{[i]}$ whose successive quotients are isomorphic to $\zeta(\mathfrak{n})$ for some $\mathfrak{n} \in \mathcal{M}$ (see Lemma 7.3 below). Since $\langle \mathfrak{m} \rangle^{[i]} \hookrightarrow \zeta(\mathfrak{m})^{[i]}$ as discussed before, any $\omega \in \text{JH}(\langle \mathfrak{m} \rangle^{[i]})$ is a composition factor of $\zeta(\mathfrak{n})$ for some $\mathfrak{n} \in \mathcal{M}$. Hence, $\text{csupp}(\omega) = \cup_{\Delta \in \mathfrak{n}} \Delta$ (counting multiplicities).

On the other hand, $\pi^{[i^*]} = \langle \{ \Delta_1^{[-1]}, \dots, \Delta_k^{[-1]} \} \rangle$. Hence $\text{csupp}(\pi^{[i^*]}) \subset \text{csupp}(\omega)$ for any $\omega \in \text{JH}(\pi^{[i]})$. □

Now we go back to consider that $\mathfrak{s} \in \mathfrak{B}(G_n)$ with $\pi_{\mathfrak{s}} \neq 0$. By the Bernstein–Zelevinsky filtration,

$$\pi_{\mathfrak{s}} \neq 0 \Rightarrow (\pi^{[i]} \times \Pi_{i-1})_{\mathfrak{s}} \neq 0$$

for some $i \leq i^*$. Now, as a similar manner to what we did for the bottom layer above, we could determine (abstractly) all possible $\mathfrak{s}' \in \mathfrak{B}(G_n)$ with $(\pi^{[i]} \times \Pi_{i-1})_{\mathfrak{s}'} \neq 0$. Then by Lemma 6.3, one sees that,

$$(\pi^{[i]} \times \Pi_{i-1})_{\mathfrak{s}} \neq 0 \Rightarrow (\pi^{[i^*]} \boxtimes \Pi_{i^*-1})_{\mathfrak{s}} \neq 0.$$

Remark 6.4 We use notations in Sect. 5.2. Fix $\mathfrak{s} \in \mathfrak{B}(G_n)$. Let $\pi \in \text{Irr}(G_{n+1})$ with $\pi_{\mathfrak{s}} \neq 0$. We consider the set of points $\sigma \in \mathfrak{X}_{\mathfrak{s}}$ such that the corresponding maximal ideal \mathcal{J}^σ satisfies $\pi/\mathcal{J}^\sigma \pi \neq 0$. The above discussion with Lemma 4.4 gives a description on such set, and in particular, such set forms an irreducible closed subvariety in $\mathfrak{X}_{\mathfrak{s}}$.

Indeed, for any submodule π' of $\pi|_{G_n}$, the analogous set for π' defines precisely the same variety as $\pi|_{G_n}$. This can be seen by refining to a submodule ω of π' that embeds to the bottom layer $\pi^{[i^*]} \times \Pi_{i^*-1}$ (as what we saw in the

proof of Theorem 6.1). Then one shows, as in the proof of Proposition 5.6, that $\omega_{N_i^*_{-1}}/\mathcal{I}\omega_{N_i^*_{-1}} \neq 0$ for any ideal \mathcal{I} of the form $\mathcal{J}_{\pi^{[i^*]}} \otimes 1 + 1 \otimes \mathcal{J}_{\tau}$. (Here $\tau \in \text{Irr}(G_{i^*_{-1}})$ without any conditions.)

6.3 Indecomposability of Zelevinsky induced modules

Let $\mathfrak{m} \in \text{Mult}$. In [7], it is shown that the restriction of $\zeta(\mathfrak{m})$ to the mirabolic subgroup is strongly indecomposable. One may expect the following conjecture, which is a stronger statement of Theorem 6.1:

Conjecture 6.5 *Let $\mathfrak{m} \in \text{Mult}$ with sum of absolute lengths of all its segments equal to $n + 1$. Then any Bernstein component of $\zeta(\mathfrak{m})|_{G_n}$ is strongly indecomposable.*

A variation of the above conjecture is to replace $\zeta(\mathfrak{m})$ in Conjecture 6.5 by $\lambda(\mathfrak{m})$. However, it is not true in general for the smooth dual $\zeta(\mathfrak{m})^\vee|_{G_n}$. An example is $\mathfrak{m} = \{[v^{-1/2}], [v^{1/2}]\}$. In this case, the short exact sequence:

$$0 \rightarrow \langle \Delta \rangle \rightarrow \zeta(\mathfrak{m})^\vee|_{G_1} \rightarrow \text{St}(\Delta) \rightarrow 0,$$

where $\Delta = [v^{-1/2}, v^{1/2}]$, gives a split sequence since $\text{St}(\Delta)|_{G_1}$ is projective (see Theorem 3.7). Hence the Iwahori component of $\zeta(\mathfrak{m})^\vee|_{G_1}$ is not indecomposable.

7 Asymmetric property of left and right derivatives

We are going to prove Theorem 6.2 in this section. The idea lies in two simple cases. The first one is a generic representation. Since an irreducible generic representation is isomorphic to $\lambda(\mathfrak{m}) \cong \text{St}(\mathfrak{m})$ for $\mathfrak{m} \in \text{Mult}$ (with the property that any two segments in \mathfrak{m} are unlinked), a simple counting on cuspidal supports of derivatives can show Theorem 6.2 for that case. The second one is an irreducible representation whose Zelevinsky multisegment has all segments with relative length strictly greater than 1. In such case, one can narrow down the possibility of irreducible submodules of the derivatives via the embedding $\langle \mathfrak{m} \rangle^{(i)} \hookrightarrow \zeta(\mathfrak{m})^{(i)}$ and ${}^{(i)}\langle \mathfrak{m} \rangle \hookrightarrow {}^{(i)}\zeta(\mathfrak{m})$, and use geometric lemma to compute the possible submodules of derivatives of $\zeta(\mathfrak{m})^{(i)}$ and ${}^{(i)}\zeta(\mathfrak{m})$. The combination of these two cases seems to require some extra work. The strategy is to use Speh representations, which can be viewed as a generalization of generalized Steinberg representations, and then apply Lemma 9.4 to obtain information on submodules.

7.1 Union–intersection operation

Let $\mathfrak{m} = \{\Delta_1, \dots, \Delta_r\}$. For two segments Δ and Δ' in \mathfrak{m} which are linked, the process of replacing Δ and Δ' by $\Delta \cap \Delta'$ and $\Delta \cup \Delta'$ is called the union–intersection process. It follows from [35, Chapter 7] that the Zelevinsky multisegment of any irreducible composition factor in

$$\langle \Delta_1 \rangle \times \dots \times \langle \Delta_r \rangle$$

can be obtained by a chain of intersection-union process. For a positive integer l , define $N(\mathfrak{m}, l)$ to be the number of segments in \mathfrak{m} with relative length l .

Lemma 7.1 *Let $k \geq 1$. Let $\mathfrak{m}_0, \mathfrak{m}_1, \dots, \mathfrak{m}_k \in \text{Mult}$ such that for $i = 1, \dots, k$, each \mathfrak{m}_i is obtained from \mathfrak{m}_{i-1} by one-step of union–intersection operation. Suppose Δ_i, Δ'_i are two segments in \mathfrak{m}_{i-1} involved in the union–intersection operation to obtain \mathfrak{m}_i . Let l_i be the relative length of $\Delta_i \cup \Delta'_i$. Then there exists l such that $l \geq l_i$ for all i , and*

$$N(\mathfrak{m}_k, l) > N(\mathfrak{m}_0, l)$$

and, for any $l' > l$,

$$N(\mathfrak{m}_k, l') \geq N(\mathfrak{m}_0, l').$$

Proof We shall prove inductively on k . The basic case that $k = 1$ can be proved by a similar argument that will be used to prove the inductive case, and so we omit the details. We assume the targeted statement is true for $X \geq 1$ number of union–intersection operations. Let \mathfrak{m}_{X+1} be a new multisegment obtained from \mathfrak{m}_0 by $X + 1$ number of union–intersection operations. Then we have a multisegment \mathfrak{m}_X obtained from \mathfrak{m}_0 by X number of union–intersection operations and \mathfrak{m}_{X+1} is obtained from \mathfrak{m}_X by one union–intersection operation.

By inductive hypothesis, we can find a positive number l_X such that

$$N(\mathfrak{m}_X, l_X) > N(\mathfrak{m}_0, l_X),$$

and for any $l' > l_X$,

$$N(\mathfrak{m}_X, l') \geq N(\mathfrak{m}_0, l').$$

Now let Δ_i, Δ_j be the segments in \mathfrak{m}_X involved in the union–intersection operation to obtain \mathfrak{m}_{X+1} . In particular, Δ_i and Δ_j are linked. Let l_0 be the relative length of $\Delta_i \cup \Delta_j$. If $l_0 \geq l_X$, set $l = l_0$, and otherwise set $l = l_X$. Now it is straightforward to check that such l satisfies the required properties. \square

7.2 Speh multisegments

Definition 7.2 Let Δ be a segment. Let

$$m(m, \Delta) = \left\{ v^{-(m-1)/2} \Delta, v^{1-(m-1)/2} \Delta, \dots, v^{(m-1)/2} \Delta \right\}.$$

We shall call $m(m, \Delta)$ to be a Speh multisegment. Define

$$u(m, \Delta) = \langle m(m, \Delta) \rangle,$$

which will be called a Speh representation. In the literature, it is sometimes called an essentially Speh representation, reflecting that it is not necessarily unitary. Denote by $L(m(m, \Delta))$ the relative length of Δ .

We similarly define

$$u_r(m, i, \Delta) = \langle v^{-(m-1)/2} \Delta^-, \dots, v^{-(m-2i+1)/2} \Delta^-, v^{-(m-2i-1)/2} \Delta, \dots, v^{(m-1)/2} \Delta \rangle,$$

and

$$u_l(m, i, \Delta) = \langle v^{-(m-1)/2} \Delta, \dots, v^{(m-2i-1)/2} \Delta, v^{(m-2i+1)/2} (-\Delta), \dots, v^{(m-1)/2} (-\Delta) \rangle.$$

Let $l = n(\rho)$. It follows from [23,33] (also see [13]) that

$$u(m, \Delta)^{(li)} \cong u_r(m, i, \Delta), \tag{7.24}$$

and $u(m, \Delta)^{(k)}$ is zero if l does not divide k . Applying (2.3), we have that

$$^{(li)}u(m, \Delta) \cong u_l(m, i, \Delta), \tag{7.25}$$

and $^{(k)}u(m, \Delta) = 0$ if l does not divide k .

7.3 Notations for multisegments

For a multisegment $m = \{\Delta_1, \dots, \Delta_r\}$, define

$$m^{(i_1, \dots, i_r)} = \left\{ \Delta_1^{(i_1)}, \dots, \Delta_r^{(i_r)} \right\}, \quad m^{(i)} = \left\{ m^{(i_1, \dots, i_r)} : i_1 + \dots + i_r = i \right\},$$

$$^{(i_1, \dots, i_r)}m = \left\{ ^{(i_1)}\Delta_1, \dots, ^{(i_r)}\Delta_r \right\}, \quad ^{(i)}m = \left\{ ^{(i_1, \dots, i_r)}m : i_1 + \dots + i_r = i \right\}.$$

We shall need the following lemma:

Lemma 7.3 *Let $\mathfrak{m} \in \text{Mult}$. Then $\zeta(\mathfrak{m})^{(i)}$ (resp. ${}^{(i)}\zeta(\mathfrak{m})$) admits a filtration whose successive subquotients are isomorphic to $\zeta(\mathfrak{n})$ for some $\mathfrak{n} \in \mathfrak{m}^{(i)}$ (resp. $\mathfrak{n} \in {}^{(i)}\mathfrak{m}$).*

Proof Write $\mathfrak{m} = \{\Delta_1, \dots, \Delta_r\}$. We shall assume that for $i < j$, $b(\Delta_i) \not\leq b(\Delta_j)$ and if $b(\Delta_i) \cong b(\Delta_j)$, then $a(\Delta_j) < a(\Delta_i)$. By definition,

$$\zeta(\mathfrak{m}) \cong \langle \Delta_1 \rangle \times \dots \times \langle \Delta_r \rangle.$$

By (2.4), we have that $\zeta(\mathfrak{m})^{(i)}$ admits a filtration whose successive quotients are:

$$\langle \Delta_1 \rangle^{(i_1)} \times \dots \times \langle \Delta_r \rangle^{(i_r)}$$

for $i_1 + \dots + i_r = i$. Now the lemma follows from that the product $\langle \Delta_1 \rangle^{(i_1)} \times \dots \times \langle \Delta_r \rangle^{(i_r)}$ is isomorphic to $\zeta(\mathfrak{n})$ for $\mathfrak{n} = \{\Delta_1^{(i_1)}, \dots, \Delta_r^{(i_r)}\}$. We remark that in order to check the isomorphism, we have to use (2.1) and our choice of labelling for \mathfrak{m} . □

7.4 Proof of Theorem 6.2

By Lemma 2.6, it suffices to prove the statement for submodules of the derivatives.

Let \mathfrak{m} be the Zelevinsky multisegment with $\pi \cong \langle \mathfrak{m} \rangle$. We shall assume that any cuspidal representation in each segment of \mathfrak{m} is an unramified twist of a fixed cuspidal representation ρ , i.e.

$$\text{csupp}(\langle \mathfrak{m} \rangle) \subset \{v^c \rho : c \in \mathbb{C}\}.$$

We shall prove Theorem 6.2 for such π . The general case follows from this by writing an irreducible representation as a product of irreducible representations of such specific form.

Step 1: First approximation using Lemma 7.3 Let π' be a common isomorphic irreducible quotient of $\pi^{[i]}$ and ${}^{[i]}\pi$. (Here we assume that $\pi^{[i]}$ and ${}^{[i]}\pi$ are non-zero.) Recall that we have that

$$\pi \hookrightarrow \zeta(\mathfrak{m}).$$

Since taking derivatives is an exact functor, $v^{1/2} \cdot \pi^{(i)}$ embeds to $v^{1/2} \cdot \zeta(\mathfrak{m})^{(i)}$ and so does π' .

By Lemma 7.3, there is a filtration on $\zeta(\mathfrak{m})^{(i)}$ given by $\zeta(\mathfrak{m}^{(i_1, \dots, i_r)})$ for $i_1 + \dots + i_r = i$, where $i_k = 0$ or $n(\rho)$. Then π' is isomorphic to the unique sub-

module of $v^{1/2}\zeta(\mathfrak{m}^{(i'_1, \dots, i'_r)})$ for some (i'_1, \dots, i'_r) and so $\pi' \cong v^{1/2}\langle \mathfrak{m}^{(i'_1, \dots, i'_r)} \rangle$. Similarly, π' is isomorphic to $v^{-1/2}\langle \mathfrak{m}^{(j'_1, \dots, j'_r)} \rangle$ for some $j'_1 + \dots + j'_r = i$.

Suppose i is not the level of π . Then there exists at least one $i'_k = 0$ and at least one $j'_k = 0$. Among all those segment Δ_k with either $i'_k = 0$ or $j'_k = 0$, we shall choose Δ_{k^*} to have the largest relative (and absolute) length. Denote the relative length of Δ_{k^*} by L .

Step 2: Second approximation using Lemma 9.4

We write \mathfrak{m} as the sum of Speh multisegments

$$\mathfrak{m} = \mathfrak{m}'_1 + \dots + \mathfrak{m}'_s \tag{7.26}$$

satisfying properties in Proposition 9.3.

Let

$$\mathfrak{m}_1, \dots, \mathfrak{m}_r$$

be all the Speh multisegments appearing in the sum (7.26) such that $L(\mathfrak{m}_k) = L$. For each \mathfrak{m}_k , write $\mathfrak{m}_k = \mathfrak{m}(\mathfrak{m}_k, \Delta_k)$ and define $b(\mathfrak{m}_k) = v^{(m_k-1)/2}b(\Delta_k)$. We shall label \mathfrak{m}_k in the way that $b(\mathfrak{m}_k) \neq b(\mathfrak{m}_l)$ for $k < l$. Furthermore, the labelling satisfies the properties that

- (\diamond) for any \mathfrak{m}_p and $p < q$, $\mathfrak{m}_p + \Delta$ is not a Speh multisegment for any $\Delta \in \mathfrak{m}_q$.
- ($\diamond\diamond$) if $\mathfrak{m}_p \cap \mathfrak{m}_q \neq \emptyset$ and $p \leq q$, then $\mathfrak{m}_q \subset \mathfrak{m}_p$.

Let \mathfrak{n}_1 be the collection of all segments Δ' in $\mathfrak{m} \setminus (\mathfrak{m}_1 + \dots + \mathfrak{m}_r)$ which satisfies either (1) $b(\mathfrak{m}_1) < b(\Delta')$ or (2) $b(\mathfrak{m}_1) \cong b(\Delta')$. Define inductively that \mathfrak{n}_k is the collection of all segments Δ' in $\mathfrak{m} \setminus (\mathfrak{m}_1 + \dots + \mathfrak{m}_r + \mathfrak{n}_1 + \dots + \mathfrak{n}_{k-1})$ that satisfies the property that either (1) $b(\mathfrak{m}_k) < b(\Delta')$, or (2) $b(\mathfrak{m}_k) \cong b(\Delta')$. (It is possible that some \mathfrak{n}_k is empty.)

By Lemma 9.4, we have a series of embeddings:

$$\begin{aligned} \langle \mathfrak{m} \rangle &\hookrightarrow \zeta(\mathfrak{n}_1) \times \langle \mathfrak{m}_1 \rangle \times \dots \times \zeta(\mathfrak{n}_r) \times \langle \mathfrak{m}_r \rangle \times \zeta(\mathfrak{n}_{r+1}) \\ &\hookrightarrow \dots \\ &\hookrightarrow \zeta(\mathfrak{n}_1) \times \langle \mathfrak{m}_1 \rangle \times \zeta(\mathfrak{n}_2) \times \langle \mathfrak{m}_2 \rangle \times \zeta(\mathfrak{n}_3 + \dots + \mathfrak{n}_r + \mathfrak{m}_3 + \mathfrak{m}_r + \mathfrak{n}_{r+1}) \\ &\hookrightarrow \zeta(\mathfrak{n}_1) \times \langle \mathfrak{m}_1 \rangle \times \zeta(\mathfrak{n}_2 + \dots + \mathfrak{n}_r + \mathfrak{m}_2 + \mathfrak{m}_r + \mathfrak{n}_{r+1}) \\ &\hookrightarrow \zeta(\mathfrak{n}_1 + \dots + \mathfrak{n}_{r+1} + \mathfrak{m}_1 + \dots + \mathfrak{m}_r) = \zeta(\mathfrak{m}) \end{aligned}$$

For simplicity, define, for $k \geq 0$,

$$\lambda_k = \zeta(\mathfrak{n}_1) \times \langle \mathfrak{m}_1 \rangle \times \dots \times \zeta(\mathfrak{n}_k) \times \langle \mathfrak{m}_k \rangle \times \zeta(\mathfrak{n}_{k+1} + \dots + \mathfrak{n}_{r+1} + \mathfrak{m}_{k+1} + \dots + \mathfrak{m}_r)$$

Step 3: Approximation on right derivatives Then for each k , we again have an embedding:

$$\pi' \hookrightarrow \nu^{1/2} \cdot \pi^{(i)} \hookrightarrow \nu^{1/2} \cdot \lambda_k^{(i)}.$$

As λ_k is a product of representations, we again have a filtration on $\lambda_k^{(i)}$ by (2.4). This gives that π' embeds to a successive quotient of the filtration:

$$\pi' \hookrightarrow \nu^{1/2} \cdot (\zeta(\mathfrak{n}_1)^{(p_1^k)} \times \langle \mathfrak{m}_1 \rangle^{(q_1^k)} \times \dots \times \zeta(\mathfrak{n}_k)^{(p_k^k)} \times \langle \mathfrak{m}_k \rangle^{(q_k^k)} \times \zeta(\mathfrak{o}_{k+1})^{(s^k)})$$

with $p_1^k + \dots + p_k^k + q_1^k + \dots + q_k^k + s^k = i$,

$$\mathfrak{o}_{k+1} = \mathfrak{n}_{k+1} + \dots + \mathfrak{n}_{r+1} + \mathfrak{m}_{k+1} + \dots + \mathfrak{m}_r.$$

Lemma 7.4 (1) $i_k = 0$ for some Δ_k in \mathfrak{m} with relative length L (see the choice of L in Step 1).

(2) Following above notations, there exists a $k' \geq 1$ such that at least one of $q_1^{k'}$ is not equal to the level of $\langle \mathfrak{m}_l \rangle$.

Remark 7.5 Similarly, we have $j_k = 0$ for some Δ_k in \mathfrak{m} with relative length L .

Proof We first assume (1) to prove (2). It suffices to show that when $k' = r$, at least one of $q_1^{k'}$ is not equal to the level of $\langle \mathfrak{m}_l \rangle$. Suppose not. Let $\langle \mathfrak{m}_i \rangle^-$ be the highest derivative of $\langle \mathfrak{m}_i \rangle$. Then we obtain an embedding:

$$\pi' \hookrightarrow \nu^{1/2} \cdot (\zeta(\mathfrak{n}_1)^{(p_1^k)} \times \langle \mathfrak{m}_1 \rangle^- \times \dots \times \zeta(\mathfrak{n}_r)^{(p_r^k)} \times \langle \mathfrak{m}_r \rangle^- \times \zeta(\mathfrak{n}_{r+1})^{(s_r)}).$$

Now set \mathfrak{m}_i^- to be the multisegment such that $\langle \mathfrak{m}_i^- \rangle \cong \langle \mathfrak{m}_i \rangle^-$. By definitions, \mathfrak{m}_i^- is still a Speh multisegment. Now, by Lemma 7.3, we have that

$$\pi' \hookrightarrow \nu^{1/2} \cdot (\zeta(\tilde{\mathfrak{n}}_1) \times \langle \mathfrak{m}_1^- \rangle \times \dots \times \zeta(\tilde{\mathfrak{n}}_r) \times \langle \mathfrak{m}_r^- \rangle \times \zeta(\tilde{\mathfrak{n}}_{r+1})),$$

where $\tilde{\mathfrak{n}}_a \in \mathfrak{n}_a^{(p_a^k)}$. From the construction of \mathfrak{n}_a , we can check that those $\tilde{\mathfrak{n}}_i$ satisfies the conditions in Lemma 9.4. Hence, by Lemma 9.4,

$$\pi' \cong \nu^{1/2} \cdot (\tilde{\mathfrak{n}}_1 + \mathfrak{m}_1^- + \dots + \tilde{\mathfrak{n}}_k + \mathfrak{m}_k^- + \tilde{\mathfrak{n}}_{k+1}).$$

Now we write $\mathfrak{n}_a = \{ \Delta_{a,1}, \dots, \Delta_{a,r(a)} \}$ for each a . Then we have that

$$\tilde{\mathfrak{n}}_a = \left\{ \Delta_{a,1}^{(p_{a,1})}, \dots, \Delta_{a,r(a)}^{(p_{a,r(a)})} \right\},$$

where each $p_{a,k} = 0$ or $n(\rho)$.

Claim For any segment $\Delta_{a,k}$ with relative length at least $L + 1$, $p_{a,k} = n(\rho)$.

Proof of Claim Recall that we also have $\pi' \cong v^{1/2} \cdot \langle m^{(i_1, \dots, i_r)} \rangle$ in Step 1. Hence,

$$m^{(i_1, \dots, i_r)} = \tilde{n}_1 + m_1^- + \dots + \tilde{n}_k + m_k^- + \tilde{n}_{k+1}. \tag{7.27}$$

Let L^{**} be the largest relative length among all the relative length of segments in m . If $L^{**} = L$, then there is nothing to prove (for the claim). We assume $L^{**} > L$. In such case, there is no segment in $m^{(i_1, \dots, i_r)}$ has relative length L^{**} . This implies that all segments $\Delta_{a,k}$ with relative length L^{**} have $p_{a,k} = n(\rho)$. With the use of (7.27) and

$$m = n_1 + m_1^- + \dots + n_k + m_k^- + n_{k+1},$$

we can proceed to the length $L^{**} - 1$ in a similar fashion. Inductively, we obtain the claim.

Now we go back to the proof of the lemma. By using the claim (and the definitions of m_i),

$$N(\tilde{n}_1 + m_1^- + \dots + \tilde{n}_k + m_k^- + \tilde{n}_{k+1}, L) = N(m, L + 1).$$

On the other hand,

$$N(m^{(i_1, \dots, i_r)}, L) \geq N(m, L + 1) + 1$$

by using the hypothesis in the lemma. However, $m = \tilde{n}_1 + m_1^- + \dots + \tilde{n}_k + m_k^- + \tilde{n}_{k+1}$ and this gives a contradiction. This proves (2) modulo (1).

It remains to prove (1). From our choice of L , either

$$N({}^{(j_1, \dots, j_r)}m, L) \geq N(m, L + 1) + 1$$

or

$$N(m^{(i_1, \dots, i_r)}, L) \geq N(m, L + 1) + 1.$$

however, since $v^{1/2} \cdot m^{(i_1, \dots, i_r)} \cong v^{-1/2} \cdot {}^{(j_1, \dots, j_r)}m$, we have that

$$N(m^{(i_1, \dots, i_r)}, L) = N({}^{(j_1, \dots, j_r)}m, L).$$

Thus, we must have that $N(m^{(i_1, \dots, i_r)}, L) \geq N(m, L + 1) + 1$. But this forces (1). \square

Now let \bar{k} be the smallest number such that at least one of $q_l^{\bar{k}}$ is not equal to the level of $\langle m_l \rangle$. Now we shall denote such l by l^* , and so $\bar{k} \geq l^*$.

We use similar strategy to further consider the filtrations on each n_a by geometric lemma. For that we write, for each a ,

$$n_a = \{ \Delta_{a,1}, \dots, \Delta_{a,r(a)} \}$$

and

$$\mathfrak{o}_{\bar{k}+1} = \left\{ \Delta_{\bar{k}+1,1}, \dots, \Delta_{\bar{k}+1,r(\bar{k}+1)} \right\}.$$

Here $r(a)$ is an index counting the number of segments in \mathfrak{n}_a depending on a .

Then we again have an embedding

$$\pi' \hookrightarrow \nu^{1/2} \cdot (\zeta(\tilde{\mathfrak{n}}_1) \times \langle \mathfrak{m}_1 \rangle^{(q_1^{\bar{k}})} \times \dots \times \zeta(\tilde{\mathfrak{n}}_{\bar{k}}) \times \langle \mathfrak{m}_{\bar{k}} \rangle^{(q_{\bar{k}}^{\bar{k}})} \times \zeta(\mathfrak{o}_{\bar{k}+1})^{(s_{\bar{k}})}),$$

where, for $a = 1, \dots, \bar{k}$,

$$\tilde{\mathfrak{n}}_a = \left\{ \Delta_{a,1}^{(p_{a,1})}, \dots, \Delta_{a,r(a)}^{(p_{a,r(a)})} \right\} \in \mathfrak{n}_a^{(p_a^{\bar{k}})}$$

with $p_{a,1} + \dots + p_{a,r(a)} = p_a^{\bar{k}}$ and each $p_{a,b} = 0$ or $n(\rho)$, and

$$\tilde{\mathfrak{o}}_{\bar{k}+1} = \left\{ \Delta_{\bar{k}+1,1}^{(p_{\bar{k}+1,1}^{\bar{k}})}, \dots, \Delta_{\bar{k}+1,r(\bar{k}+1)}^{(p_{\bar{k}+1,r(\bar{k}+1)}^{\bar{k}})} \right\} \in \mathfrak{o}_{\bar{k}+1}^{(s_{\bar{k}})}$$

with $p_{\bar{k}+1,1} + \dots + p_{\bar{k}+1,r(\bar{k}+1)} = s_{\bar{k}}$, with each $p_{\bar{k}+1,b} = 0$ or $n(\rho)$.

Step 4: Computing some indexes $p_{a,b}$ and approximating the number of special segments by union–intersection operations We claim (*) that if $\Delta_{a,b}$ has a relative length at least $L + 1$, then $p_{a,b} = n(\rho)$. This is indeed similar to the proof of Lemma 7.4 and the main difference is that we do not have an analogous form of (7.27) (since we cannot apply Lemma 9.4). Instead, we can obtain this from Lemma 7.1 (and its proof of Lemma 7.1).

Now from our choice of \bar{k} , we have that $\langle \mathfrak{m}_{l^*} \rangle^{(q_{l^*}^{\bar{k}})}$ is not a Speh representation. We can write

$$\mathfrak{m}_{l^*} = \left\{ \nu^{-x+1} \Delta^*, \dots, \nu^{-1} \Delta^*, \Delta^* \right\}$$

for a certain Δ^* with relative length L and some x . By (\diamond), $\nu \Delta^* \notin \mathfrak{m}_l$ for any $l \geq l^*$ from our labelling on \mathfrak{m}_l . Rephrasing the statement, we get the following statement:

$$(**) \nu^{1/2} \Delta^* \notin \nu^{-1/2} \mathfrak{m}_l \text{ for any } l \geq l^*.$$

Now with (*), we have that π' is a composition factor of $\zeta(\mathfrak{m}''')$, where \mathfrak{m}''' is some multisegment containing all the segments Δ^- (counting multiplicities) with Δ in \mathfrak{m} that has relative length at least $L + 1$ and containing an additional segment $\nu^{1/2} \Delta^*$ (from $\langle \mathfrak{m}_{l^*} \rangle^{(q_{l^*}^{\bar{k}})}$ by Sect. 7.2), and we shall call the former segments (i.e. the segment in the form of Δ^-) to be *special* for convenience.

We can apply the intersection-union process to obtain the Zelevinsky multisegment for π' from \mathfrak{m}''' . However in each step of the process, any one of the two segments involved in the intersection-union cannot be special. Otherwise, by Lemma 7.1, there exists $l \geq L + 1$ such that the number of segments in the resulting multisegment with

relative length l is more than the number of segments in $\mathfrak{m}^{(i_1, \dots, i_r)}$. (Here we also used that $L(\mathfrak{m}^{(i_1, \dots, i_r)}, l) = L(\mathfrak{m}''', l)$ for any $l \geq L + 1$.) Hence we obtain the following:

(***) *the number of segments Δ in $v^{1/2} \cdot \mathfrak{m}^{(i_1, \dots, i_r)}$ such that $v^{1/2} \Delta^* \subset \Delta$ is at least equal to one plus the number special segments in \mathfrak{m}''' satisfying the same property*

Step 5: Approximation on left derivatives Now we come to the final part of the proof. We now consider $v^{-1/2} \cdot {}^{(i)}\pi$. Following the strategy for right derivatives, we have that for each k

$$\pi' \hookrightarrow v^{-1/2} \cdot {}^{(i)}\pi \hookrightarrow v^{-1/2} \cdot {}^{(i)}\lambda_k.$$

This gives that π'

$$\pi' \hookrightarrow v^{-1/2} \cdot ({}^{(u_1^k)}\zeta(\mathfrak{n}_1) \times ({}^{(v_1^k)}\langle \mathfrak{m}_1 \rangle \times \dots \times ({}^{(u_k^k)}\zeta(\mathfrak{n}_k) \times ({}^{(v_k^k)}\zeta(\mathfrak{m}_k) \times ({}^{(w_k^k)}\zeta(\mathfrak{o}_{k+1})))$$

with $u_1^k + \dots + u_k^k + v_1^k + \dots + v_k^k + w^k = i$. Let \tilde{k} be the smallest integer such that at least one of $v_l^{\tilde{k}}$ is not equal to the level of $\langle \mathfrak{m}_l \rangle$. (For the existence of \tilde{k} , see Lemma 7.4.) We shall denote such l by \tilde{l} .

Step 6: the case that $\tilde{k} \geq \bar{k}$ and determining the number of special segments

We firstly consider the case that $\tilde{k} \geq \bar{k}$. We abbreviate $v_p = v_p^{\tilde{k}-1}$. Similar to right derivatives, we have that

$$\pi' \hookrightarrow v^{-1/2} \cdot \zeta(\widehat{\mathfrak{n}}_1) \times ({}^{(v_1)}\langle \mathfrak{m}_1 \rangle \times \dots \times \zeta(\widehat{\mathfrak{n}}_{\tilde{k}-1}) \times ({}^{(v_{\tilde{k}-1})}\langle \mathfrak{m}_{\tilde{k}-1} \rangle \times ({}^{(w_{\tilde{k}-1})}\zeta(\widehat{\mathfrak{o}}_{\tilde{k}})),$$

where

$$\widehat{\mathfrak{n}}_a = \left\{ ({}^{(u_{a,1})}\Delta_{a,1}, \dots, ({}^{(u_{a,r(a)})}\Delta_{a,r(a)}) \right\}.$$

with $u_{a,1} + \dots + u_{a,r(a)} = u_a^{\tilde{k}-1}$ and each $u_{a,b} = 0$ or $n(\rho)$. Since we assume that $\tilde{k} \geq \bar{k}$, we have that $({}^{(v_l)}\langle \mathfrak{m}_l \rangle)$ is a highest derivative for any $l \leq \tilde{k} - 1$ and so is a Speh representation, and we can apply Lemma 9.4(1). Hence the unique subrepresentation of

$$v^{-1/2} \cdot \zeta(\widehat{\mathfrak{n}}_1) \times ({}^{(v_1)}\langle \mathfrak{m}_1 \rangle \times \dots \times \zeta(\widehat{\mathfrak{n}}_{\tilde{k}-1}) \times ({}^{(v_{\tilde{k}-1})}\langle \mathfrak{m}_{\tilde{k}-1} \rangle \times \zeta(\widehat{\mathfrak{o}}_{\tilde{k}})$$

is isomorphic to

$$v^{-1/2} \cdot (\widehat{\mathfrak{n}}_1 + \widehat{\mathfrak{m}}_1 + \dots + \widehat{\mathfrak{n}}_{\tilde{k}-1} + \widehat{\mathfrak{m}}_{\tilde{k}-1} + \widehat{\mathfrak{o}}_{\tilde{k}}), \tag{7.28}$$

where $\langle \widehat{\mathfrak{m}}_l \rangle = ({}^{(v_l)}\langle \mathfrak{m}_l \rangle)$. Similar to (*) for right derivatives (but the proof could be easier here), we obtain the analogous statement for those $\widehat{\mathfrak{n}}_a$. Now if

$$\Delta \in v^{-1/2}(\widehat{\mathfrak{n}}_1 + \widehat{\mathfrak{m}}_1 + \dots + \widehat{\mathfrak{n}}_{\tilde{k}-1} + \widehat{\mathfrak{m}}_{\tilde{k}-1} + \widehat{\mathfrak{o}}_{\tilde{k}})$$

such that $\Delta = \nu^{1/2} \Delta^*$, then we must have that $\Delta = \nu^{-1/2} \cdot -\Delta_0$ or $\nu^{-1/2} \cdot \Delta_0$ for some segment Δ_0 in m . For the latter case to happen, we must have that $\Delta_0 \in m_l \subset o_{\bar{k}}$ with $l \geq \bar{k} \geq l^*$, but, by (**), the possibility $\Delta = \nu^{-1/2} \Delta_0$ cannot happen. Thus we must have that Δ is a special in the same sense as the discussion in right derivatives. This concludes the following:

(****) *The number of segments Δ in $\nu^{-1/2}(\widehat{n}_1 + \widehat{m}_1 + \dots + \widetilde{n}_{\bar{k}-1} + \widehat{m}_{\bar{k}-1} + \widehat{o}_{\bar{k}})$ with the property that $\nu^{1/2} \Delta^* \subset \Delta$ is equal to the number of special segments satisfying the same property.*

Now the above statement contradicts to (***) since both Zelevinsky multisegments give an irreducible representation isomorphic to π' .

Remark 7.6 The isomorphism (7.28) is a key to obtain the equality in (****), in contrast with the inequality in (***) .

Step 7: the case that $\bar{k} > \widetilde{k}$ Now the way to get contradiction in the case $\bar{k} \geq \widetilde{k}$ is similar by interchanging the role of left and right derivatives. We remark that to prove the analogue of (**), one uses (\diamond). And to obtain the similar isomorphism as (7.28), one needs to use Lemma 9.4(2). We can argue similarly to get an analogue of (***) and (****). Hence the only possibility that $\nu^{1/2} \cdot \pi^{(i)}$ and so $\nu^{-1/2} \cdot {}^{(i)}\pi$ have an isomorphic irreducible quotient only if i is the level for π . This completes the proof of Theorem 6.2.

7.5 Branching law in opposite direction

Here is another consequence of the asymmetric property on the Hom-branching law in another direction:

Corollary 7.7 *Let $\pi' \in \text{Irr}(G_n)$. Let $\pi \in \text{Irr}(G_{n+1})$.*

(1) *Suppose π is not 1-dimensional. Then*

$$\text{Hom}_{G_n}(\pi', \pi|_{G_n}) = 0.$$

(2) *Suppose π is 1-dimensional. Then*

$$\text{Hom}_{G_n}(\pi', \pi|_{G_n}) \neq 0$$

if and only if π' is also 1-dimensional and $\pi' = \pi|_{G_n}$.

Proof The second statement is trivial. We consider the first one. Since π is not one-dimensional, the level of π is not 1 by Zelevinsky classification. By Theorem 6.2 and Proposition 2.5, $\pi^{[1]}$ and ${}^{[1]}\pi$ have no common irreducible submodule if $\pi^{(1)} \neq 0$ and ${}^{(1)}\pi \neq 0$. Then at least one of

$$\text{Hom}_{G_n}(\pi', \pi^{[1]}) = 0 \quad \text{or} \quad \text{Hom}_{G_n}(\pi', {}^{[1]}\pi) = 0.$$

On the other hand, we have that for all $i \geq 2$, by Frobenius reciprocity,

$$\text{Hom}_{G_n}(\pi', \text{ind}_{R_{n-i+1}}^{G_n} \pi^{[i]} \boxtimes \psi_{i-1}) \cong \text{Hom}_{G_{n+1-i} \times G_{i-1}}(\pi'_{N_{i-1}}, \pi^{[i]} \boxtimes \Pi_{i-1}),$$

and

$$\text{Hom}_{G_n}(\pi', \text{ind}_{R_{n-i+1}}^{G_n} [i] \pi \boxtimes \psi_{i-1}) \cong \text{Hom}_{G_{i-1} \times G_{n+1-i}}(\pi'_{N_{n+1-i}}, \Pi_{i-1} \boxtimes [i] \pi)$$

However, in the last two equalities, the Hom in the right hand side is always zero (see Lemma 5.3). Now applying a Bernstein–Zelevinsky filtration to obtain the corollary. \square

8 Preserving indecomposability of Bernstein–Zelevinsky induction

We shall use the following criteria of indecomposable representations:

Lemma 8.1 *Let G be a connected reductive p -adic group. Let π be a smooth representation of G . The only idempotents in $\text{End}_G(\pi)$ are 0 and the identity (up to automorphism) if and only if π is indecomposable.*

Proof If π is not indecomposable, then any projection to a direct summand gives a non-identity idempotent. On the other hand, if $\sigma \in \text{End}_G(\pi)$ is a non-identity idempotent (i.e. $\sigma(\pi) \neq \pi$), then $\pi \cong \text{im}(\sigma) \oplus \text{im}(1 - \sigma)$. \square

Lemma 8.2 *Set $\Pi = \Pi_k$. Let $\mathfrak{s} \in \mathfrak{B}(G_k)$. Then ${}^{(k)}(\Pi_{\mathfrak{s}}) \cong \mathfrak{Z}_{\mathfrak{s}}$, where $\mathfrak{Z}_{\mathfrak{s}}$ is the Bernstein center of $\mathfrak{R}_{\mathfrak{s}}(G_k)$.*

Proof We have

$${}^{(k)}(\Pi_{\mathfrak{s}}) \cong \text{Hom}_{\mathbb{C}}(\mathbb{C}, {}^{(k)}(\Pi_{\mathfrak{s}})) \cong \text{Hom}_{G_k}(\Pi, \Pi_{\mathfrak{s}}) \cong \text{End}_{G_k}(\Pi_{\mathfrak{s}}),$$

where the second isomorphism follows from the second adjointness theorem in [15], and the third isomorphism follows from the Bernstein decomposition. Now the lemma follows from that the final endomorphism is isomorphic to $\mathfrak{Z}_{\mathfrak{s}}$ by [8]. \square

Theorem 8.3 *Let π be an admissible indecomposable smooth representation of G_{n-i} . For each $\mathfrak{s} \in \mathfrak{B}(G_n)$, the Bernstein component $(\text{ind}_{R_{n-i}}^{G_n} \pi \boxtimes \psi_i)_{\mathfrak{s}}$ is indecomposable whenever it is nonzero.*

Proof We first prove the following:

Lemma 8.4 *Let $\pi \in \text{Alg}(G_{n-i})$ be admissible. Then, as algebras,*

$$\text{End}_{G_n}(\text{ind}_{R_{n-i}}^{G_n} \pi \boxtimes \psi_i) \cong \text{End}_{G_{n-i}}(\pi) \otimes \mathfrak{Z}_i,$$

where $\mathfrak{Z}_i = \prod_{\mathfrak{s} \in \mathfrak{B}(G_i)} \mathfrak{Z}_{\mathfrak{s}}$ is the Bernstein center.

Proof Let $\mathfrak{s}_1, \mathfrak{s}_2 \in \mathfrak{B}(G_{n-i})$. Let π_1, π_2 be two admissible G_{n-i} representations in $\mathfrak{R}_{\mathfrak{s}_1}(G_{n-i})$ and $\mathfrak{R}_{\mathfrak{s}_2}(G_{n-i})$ respectively. Then

$$\begin{aligned} & \text{Hom}_{G_n}(\text{ind}_{R_{n-i}}^{G_n} \pi_1 \boxtimes \psi_i, \text{ind}_{R_{n-i}}^{G_n} \pi_2 \boxtimes \psi_i) \\ & \cong \prod_{\mathfrak{s} \in \mathfrak{R}(G_n)} \text{Hom}_{G_n}(\text{ind}_{R_{n-i}}^{G_n} \pi_1 \boxtimes \psi_i, (\text{ind}_{R_{n-i}}^{G_n} \pi_2 \boxtimes \psi_i)_{\mathfrak{s}}) \\ & \cong \prod_{\mathfrak{s} \in \mathfrak{R}(G_n)} \text{Hom}_{G_n}(\pi_1, {}^{(i)}(\text{ind}_{R_{n-i}}^{G_n} \pi_2 \boxtimes \psi_i)_{\mathfrak{s}}) \quad [16, \text{Lemma 2.4}] \\ & \cong \prod_{\mathfrak{u} \in \mathfrak{R}(G_i)} \text{Hom}_{G_{n-i}}(\pi_1, {}^{(i)}(\pi_2 \times (\Pi_i)_{\mathfrak{u}})) \end{aligned}$$

We remark that in order to apply the second adjointness theorem [15, Lemma 2.4], we need to get the first isomorphism. For the third isomorphisms, see discussions in Sect. 5.1.

Now we apply the geometric lemma on ${}^{(i)}(\pi_2 \times (\Pi_i)_{\mathfrak{u}})$ and we obtain a filtration on ${}^{(i)}(\pi_2 \times (\Pi_i)_{\mathfrak{u}})$ whose successive quotients are of the form

$${}^{(j)}(\pi_2) \times {}^{(k)}((\Pi_i)_{\mathfrak{u}}),$$

where $j + k = i$. Using Lemma 5.3,

$$\text{Hom}_{G_{n-i}}(\pi_1, {}^{(j)}(\pi_2) \times {}^{(k)}((\Pi_i)_{\mathfrak{u}})) = 0$$

unless $j = 0$ and $k = i$. Combining the above isomorphisms, we now have

$$\begin{aligned} & \text{Hom}_{G_n}(\text{ind}_{R_{n-i}}^{G_n} \pi_1 \boxtimes \psi_i, \text{ind}_{R_{n-i}}^{G_n} \pi_2 \boxtimes \psi_i) \\ & \cong \prod_{\mathfrak{u} \in \mathfrak{B}(G_i)} \text{Hom}_{G_{n-i}}(\pi_1, \pi_2 \otimes {}^{(i)}((\Pi_i)_{\mathfrak{u}})) \\ & \cong \prod_{\mathfrak{u} \in \mathfrak{B}(G_i)} \text{Hom}_{G_{n-i}}(\pi_1, \pi_2 \otimes \mathfrak{Z}_{\mathfrak{u}}) \\ & \cong \prod_{\mathfrak{u} \in \mathfrak{B}(G_i)} \text{Hom}_{G_{n-i}}(\pi_1, \pi_2) \otimes \mathfrak{Z}_{\mathfrak{u}} \\ & \cong \text{Hom}_{G_{n-i}}(\pi_1, \pi_2) \otimes \mathfrak{Z} \end{aligned}$$

We remark that the third and last isomorphisms require the admissibility of π_1 and π_2 . The second isomorphism follows from Lemma 8.2. One further traces the above isomorphisms to see that they give an isomorphism preserving the algebra structure. Now we specialize to $\pi_1 = \pi_2 = \pi$ and we obtain the lemma when each of π_1, π_2 lying in one Bernstein component.

The general case follows by writing $\pi = \bigoplus_{\mathfrak{s}} \pi_{\mathfrak{s}}$. □

Now the theorem follows from Lemmas 8.1 and 8.3 since the right hand side has only 0 and 1 as idempotents if and only if the left hand side does. □

Let M_n be the mirabolic subgroup in G_{n+1} , that is the subgroup containing all matrices with the last row of the form $(0, \dots, 0, 1)$. We have the following consequence restricted from M_n to G_n . Here G_n is viewed as the subgroup of M_n via the embedding $g \mapsto \text{diag}(g, 1)$.

Corollary 8.5 *Let π be an irreducible smooth representation of M_n . Then for any $s \in \mathfrak{B}(G_n)$, π_s is indecomposable.*

Proof This follows from [7, Corollary 3.5] and Theorem 8.3. □

9 Appendix: Speh representation approximation

9.1 Two lemmas

Some studies on parabolic induction can be found in [24,32,34]. We give a proof of the following specific cases, using the theory of derivatives. We use notations in Sect. 7.2.

Lemma 9.1 *Let $m \in \mathbb{Z}$. Let $\Delta = [v^a \rho, v^b \rho]$. If $\Delta' = [v^k \rho, v^l \rho]$ such that*

$$-\frac{m-1}{2} + a \leq k \leq l \leq \frac{m-1}{2} + b,$$

then

$$u(m, \Delta) \times \langle \Delta' \rangle \cong \langle \Delta' \rangle \times u(m, \Delta),$$

and is irreducible.

Proof When $-(m-1)/2 + a - k \notin \mathbb{Z}$, this case is easier by using (2.1) and we omit the details. We now consider $-(m-1)/2 + a - k \in \mathbb{Z}$.

The statement is not difficult when Δ is a singleton $[\rho]$ because

$$u(m, \Delta) \cong \text{St}([v^{-(m-1)/2} \rho, v^{(m-1)/2} \rho])$$

and $\text{St}(\Delta') \times \langle \Delta'' \rangle \cong \langle \Delta'' \rangle \times \text{St}(\Delta')$ whenever $\Delta'' \subset \Delta'$. (Indeed, one can also prove the latter fact by similar arguments as below by noting that the Zelevinsky multisegment of any simple composition factor in $\text{St}(\Delta') \times \langle \Delta'' \rangle$ contains a segment $\tilde{\Delta}$ with $b(\tilde{\Delta}) \cong b(\Delta')$ and at least one segment $\hat{\Delta}$ with $b(\hat{\Delta}) \cong b(\Delta'')$.)

We now assume Δ is not a singleton. We consider two cases:

- (1) Case 1: $l = b + \frac{m-1}{2}$. Suppose τ is a composition factor of $u(m, \Delta) \times \langle \Delta' \rangle$ with the associated Zelevinsky multisegment m . Then we know that at least two segments Δ_1, Δ_2 in m takes the form $b(\Delta_1) \cong b(\Delta_2) \cong v^{\frac{m-1}{2}+b} \rho$. If $\tau^{(i^*)}$ is the highest derivative of τ , then we know that the cuspidal support of $\tau^{(i^*)}$ does not contain $v^{\frac{m-1}{2}} \rho$. We also have that $\tau^{(i^*)}$ is a composition factor of $(u(m, \Delta) \times \langle \Delta' \rangle)^{(i^*)}$. The only possibility is that $i^* = m + 1$ i.e. $(u(m, \Delta) \times \langle \Delta' \rangle)^{(i^*)} = u(m, \Delta^-) \times \langle (\Delta')^- \rangle$, which the latter one is irreducible by induction. This proves

the lemma. Since taking derivative is an exact functor, we have that $u(m, \Delta) \times \langle \Delta' \rangle$ is irreducible. Using the Gelfand–Kazhdan involution [6, Section 7], we have that $\langle \Delta' \rangle \times u(m, \Delta) \cong u(m, \Delta) \times \langle \Delta' \rangle$.

- (2) Case 2: $l < b + \frac{m-1}{2}$. The argument is similar. Again suppose τ is a composition factor of $u(m, \Delta) \times \langle \Delta' \rangle$. Using an argument similar to above, we have that the level of τ is either $m + 1$ or m . However, if the level of τ is m , then $\tau^{(m)}$ would be $u(m, \Delta^-) \times \langle \Delta' \rangle$, which is irreducible by induction. Then it would imply that the number of segments in Zelevinsky multisegment of $\tau^{(m)}$ is m and contradicts that the number of segments for the Zelevinsky multisegment of the highest derivative of an irreducible representation π must be at most that for π . Hence, the level of τ must be $m + 1$. Now repeating a similar argument as in (1) and using the induction, we obtain the statements. □

Lemma 9.2 *Let Δ be a segment. Then*

$$u(m, \Delta^-) \times v^{(m-1)/2} \langle \Delta \rangle \cong v^{(m-1)/2} \langle \Delta \rangle \times u(m, \Delta^-)$$

is irreducible.

Proof The statement is clear if Δ is a singleton. For the general case, we note by simple counting that the Zelevinsky multisegment of any composition factor must contain a segment Δ_1 with $b(\Delta_1) \cong v^{(m-1)/2} b(\Delta)$ and at least one segment Δ_2 with $b(\Delta_2) \cong v^{(m-1)/2} b(\Delta^-)$. Then one proves the statement by a similar argument using the highest derivative as in the previous lemma. □

9.2 Spheh representation approximation

For a Spheh multisegment

$$m = \{ \Delta, v^{-1} \Delta, \dots, v^{-k} \Delta \},$$

define $b(m) = b(\Delta)$.

Proposition 9.3 *Let $m = \{ \Delta_1, \dots, \Delta_k \}$. Then there exists Spheh multisegments m_1, \dots, m_r satisfying the following properties:*

- (1) $m = m_1 + \dots + m_r$;
- (2) *For each Spheh multisegment m_i and any $j > i$, there is no segment Δ in m_i such that $m_j + \{ \Delta \}$ is a Spheh multisegment;*
- (3) $\langle m \rangle$ *is the unique submodule of $\langle m_1 \rangle \times \dots \times \langle m_r \rangle$;*
- (4) $b(m_i) \not\leq b(m_j)$ *if $i < j$;*
- (5) *if $m_i \cap m_j \neq \emptyset$ and $i \leq j$, then $m_j \subset m_i$.*

Proof We shall label the segments m in the way that for $i < j$, either (i) $b(\Delta_i) \not\leq b(\Delta_j)$ and (ii) if $b(\Delta_i) = b(\Delta_j)$, then $a(\Delta_i) \not\leq a(\Delta_j)$. Let $\Delta = \Delta_1$. Let k be the largest integer ($k \geq 0$) such that $\Delta, v^{-1} \Delta, \dots, v^{-k} \Delta$ are segments in m . We claim that

$$\langle m \rangle \hookrightarrow \langle m' \rangle \times \langle m \setminus m' \rangle \tag{9.29}$$

and moreover $\langle m \rangle$ is the unique submodule of $\langle m' \rangle \times \langle m \setminus m' \rangle$. By induction, the claim proves (1) and (3), and the remaining ones follow from the inductive construction.

We now prove the claim. We shall prove by induction that for $i = 0, \dots, k$,

$$\langle m \rangle \hookrightarrow \langle m_i \rangle \times \zeta(m \setminus m_i) \hookrightarrow \zeta(m),$$

where $m_i = \{\Delta, v^{-1}\Delta, \dots, v^{-i}\Delta\}$. When $i = 0$ the statement is clear from the definition. Now suppose that we have the inductive statement for i . To prove the statement for $i + 1$. We now set n to be the collection of all segments Δ' in m such that $b(\Delta') = b(\Delta), v^{-1}b(\Delta), \dots, v^{-i}b(\Delta)$ and $v^{-i}a(\Delta)$ precedes $a(\Delta')$. We also set \bar{n} to be the collection of all segments Δ' in m such that $b(\Delta') = b(\Delta), v^{-1}b(\Delta), \dots, v^{-i}b(\Delta)$ and $a(\Delta')$ precedes $v^{-i-1}a(\Delta)$. By using (2.1) several times, we have that $\zeta(n) \times \zeta(\bar{n}) \cong \zeta(\bar{n} + n)$ and

$$\zeta(m \setminus m_i) \cong \zeta(n) \times \zeta(\bar{n}) \times \zeta(m \setminus (n + \bar{n} + m_i)). \tag{9.30}$$

From our construction (and $i \neq k$), we have that $v^{-i-1}\Delta$ is a segment in $m \setminus (n + \bar{n} + m_i)$ and

$$\zeta(m \setminus (n + \bar{n} + m_i)) = \langle v^{-i-1}\Delta \rangle \times \zeta(m \setminus (n + \bar{n} + m_{i+1}))$$

On the other hand

$$\langle m_i \rangle \times \zeta(n) \times \zeta(\bar{n}) \times \langle v^{-i-1}\Delta \rangle \tag{9.31}$$

$$\cong \zeta(n) \times \langle m_i \rangle \times \zeta(\bar{n}) \times \langle v^{-i-1}\Delta \rangle \tag{9.32}$$

$$\cong \zeta(n) \times \langle m_i \rangle \times \langle v^{-i-1}\Delta \rangle \times \zeta(\bar{n}) \tag{9.33}$$

$$\hookrightarrow \zeta(n) \times \langle m_{i+1} \rangle \times \zeta(\bar{n}) \tag{9.34}$$

$$\cong \langle m_{i+1} \rangle \times \zeta(n) \times \zeta(\bar{n}) \tag{9.35}$$

The first and last isomorphisms follow from Lemma 9.3. The injectivity in the forth line comes from the uniqueness of submodule in $\zeta(m_{i+1})$. The second isomorphism follows from again by (2.1). Combining (9.30), the above series of isomorphisms and the inductive case, we have:

$$\zeta(m) \hookrightarrow \langle m_{i+1} \rangle \times \zeta(n) \times \zeta(\bar{n}) \times \zeta(m \setminus (n + \bar{n} + m_{i+1})).$$

This gives the desired injectivity by using the uniqueness of submodule of $\zeta(m)$ and proves the claim. □

We shall need a variation which is more flexible in our application.

Lemma 9.4 *Let $m = m(m, \Delta)$ be a Speh multisegment. Let n_1 be a Zelevinsky multisegment such that for any segment Δ' in n_1 satisfying $b(\Delta) < b(\Delta')$ or $b(\Delta) \cong b(\Delta')$. Let n_2 be a Zelevinsky multisegment such that for any segment Δ' in n_2 satisfying either one of the following properties:*

- (1) $b(\Delta) \neq b(\Delta')$; or
- (2) $b(\Delta) \neq b(\Delta')$, or if $b(\Delta) < b(\Delta')$, then $(\Delta')^- = \Delta$.

Then

$$\langle \mathfrak{n}_1 + \mathfrak{m} + \mathfrak{n}_2 \rangle \hookrightarrow \zeta(\mathfrak{n}_1) \times \langle \mathfrak{m} \rangle \times \zeta(\mathfrak{n}_2) \hookrightarrow \zeta(\mathfrak{n}_1 + \mathfrak{m} + \mathfrak{n}_2).$$

In particular, $\zeta(\mathfrak{n}_1) \times \langle \mathfrak{m} \rangle \times \zeta(\mathfrak{n}_2)$ has unique submodule isomorphic to $\langle \mathfrak{n}_1 + \mathfrak{m} + \mathfrak{n}_2 \rangle$.

Remark 9.5 Case (2) covers case (1). But for the purpose of clarity of an argument used in Sect. 7.4, we divide into two cases. We also recall that the case $b(\Delta) \neq b(\Delta')$ includes the possibility $b(\Delta) \cong b(\Delta')$.

Proof For all cases, we have that

$$\zeta(\mathfrak{n}_1) \times \zeta(\mathfrak{m} + \mathfrak{n}_2) \cong \zeta(\mathfrak{n}_1 + \mathfrak{m} + \mathfrak{n}_2).$$

Using (9.29) for (1) we obtain the lemma. For (2), let \mathfrak{n}' be all the segments in \mathfrak{n}_2 with the property that $(\Delta')^- \cong \Delta$. Then we have that

$$\zeta(\mathfrak{n}') \times \zeta(\mathfrak{m}) \times \zeta(\mathfrak{n}_2 \setminus \mathfrak{n}') \hookrightarrow \zeta(\mathfrak{n}') \times \zeta((\mathfrak{m} + \mathfrak{n}_2) \setminus \mathfrak{n}') \hookrightarrow \zeta(\mathfrak{m} + \mathfrak{n}_2).$$

By Lemma 9.2, we have that

$$\zeta(\mathfrak{n}') \times \zeta(\mathfrak{m}) \times \zeta(\mathfrak{n}_2 \setminus \mathfrak{n}') \cong \zeta(\mathfrak{m}) \times \zeta(\mathfrak{n}') \times \zeta(\mathfrak{n}_2 \setminus \mathfrak{n}') \cong \zeta(\mathfrak{m}) \times \zeta(\mathfrak{n}_2),$$

which proves the lemma. □

10 Appendix

Let G be a connected reductive group over a non-Archimedean local field.

Lemma 10.1 *Let π be a smooth representation of G . Let π admits a filtration*

$$0 = \pi'_0 \subset \pi'_1 \subset \dots \subset \pi'_r = \pi$$

Suppose π admits an irreducible subquotient τ . Let $\pi_k = \pi'_k / \pi'_{k-1}$ for $k = 1, \dots, r$. Then there exists s such that π_s contains an irreducible subquotient isomorphic to τ .

Proof Let λ_1, λ_2 be submodules of π such that $\lambda_2 / \lambda_1 = \tau$. Now we have a filtration on λ_p ($p = 1, 2$):

$$0 \subset (\pi'_1 \cap \lambda_p) \subset (\pi'_2 \cap \lambda_p) \subset \dots \subset (\pi'_r \cap \lambda_p) = \lambda_p.$$

Now λ_2 / λ_1 admits a filtration

$$0 \subset \gamma_1 \subset \gamma_2 \subset \dots \subset \gamma_r,$$

where $\gamma_k = (\pi'_k \cap \lambda_2)/(\pi'_k \cap \lambda_1)$ and so

$$\gamma_{k+1}/\gamma_k \cong \frac{(\pi'_{k+1} \cap \lambda_2)/(\pi'_{k+1} \cap \lambda_1)}{(\pi'_k \cap \lambda_2)/(\pi'_k \cap \lambda_1)}.$$

This implies $\pi'_{k+1} \cap \lambda_2$ and so π'_{k+1} has irreducible subquotient isomorphic to τ . \square

Lemma 10.2 *Let π be a non-zero smooth representation of G . Let τ be a non-zero G -submodule of π . Suppose π admits a filtration on*

$$0 = \pi_0 \subset \pi_1 \subset \dots \subset \pi_r = \pi.$$

Then there exists a non-zero G -submodule τ' of τ such that

$$\tau' \hookrightarrow \pi_{k+1}/\pi_k.$$

for some k .

Proof Let k be the smallest positive integer such that

$$\tau \cap \pi_k \neq 0.$$

Define $\tau' = \tau \cap \pi_k$. The non-zero G -submodule τ' embeds to π_k/π_{k-1} as desired. \square

Lemma 10.3 *Let $0 \neq \pi \in \text{Alg}(G)$. Suppose π admits two G -filtrations:*

$$0 = \pi_0 \subset \pi_1 \subset \pi_2 \subset \dots \subset \pi_r = \pi$$

and

$$0 = \pi'_0 \subset \pi'_1 \subset \pi'_2 \subset \dots \subset \pi'_s = \pi.$$

Then there exists a non-zero $\tau \in \text{Alg}(G)$ such that for some i, j ,

$$\tau \hookrightarrow \pi_{i+1}/\pi_i, \quad \tau \hookrightarrow \pi'_{j+1}/\pi'_j.$$

Proof When $\pi_1 \neq 0$, we consider the smallest integer k such that $\pi_1 \cap \pi'_k \neq 0$. The general case is similar. \square

Acknowledgements This article is a part of the project in studying Ext-branching laws, and the author would like to thank Gordan Savin for a number of helpful discussions. Part of the idea on indecomposability for restriction was developed during the participation in the program of 'On the Langlands Program: Endoscopy and Beyond' at IMS of National University of Singapore in January 2019. The author would like to thank the organizers for their warm hospitality. The author would like to thank the anonymous referee for useful comments and a remark related to Remark 6.4.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Adams, J.D., Prasad, D., Savin, G.: Euler–Poincaré characteristic for the oscillator representation. In: Cogdell, J., Kim, J.L., Zhu, C.B. (eds.) Representation Theory, Number Theory, and Invariant Theory. Progress in Mathematics, vol. 323. Birkhäuser, Basel (2017)
2. Aizenbud, A., Gourevitch, D., Rallis, S., Schiffmann, G.: Multiplicity one theorems. *Ann. Math. (2)* **172**(2), 1407–1434 (2010)
3. Bernstein, J.N.: Le centre de Bernstein. In: Deligne, P. (ed.) Travaux en Cours, Representations of reductive groups over a local field, pp. 1–32, Hermann, Paris (1984)
4. Bernstein, J.N.: P-invariant distributions on $GL(N)$ and the classification of unitary representations of $GL(N)$ (non-Archimedean case). In: Lie Group Representations, II (College Park, Md., 1982/1983), Lecture Notes in Mathematics, vol. 1041, pp. 50–102. Springer, New York (1984)
5. Bernstein, J.: Representations of p-Adic Groups, Lecture Notes. Harvard University, Cambridge (1992)
6. Bernstein, I.N., Zelevinsky, A.V.: Representations of the group $GL(n, F)$, where F is a non-archimedean local field. *Russ. Math. Surv.* **31**(3), 1–68 (1976)
7. Bernstein, I.N., Zelevinsky, A.V.: Induced representations of reductive p-adic groups, I. *Ann. Sci. Ecole Norm. Sup.* **10**, 441–472 (1977)
8. Bushnell, C., Henniart, G.: Generalized Whittaker models and the Bernstein center. *Am. J. Math.* **125**(3), 513–547 (2003)
9. Bushnell, C.J., Kutzko, P.C.: The Admissible Dual of $GL(N)$ via Compact Open Subgroups. *Annals Mathematics Studies*. Princeton University Press, Princeton (1993)
10. Bushnell, C.J., Kutzko, P.C.: Semisimple types in GL_n . *Compos. Math.* **119**(1), 53–97 (1999)
11. Chan, K.Y.: Duality for Ext-groups and extensions of discrete series for graded Hecke algebras. *Adv. Math.* **294**, 410–453 (2016)
12. Chan, K.Y.: Restriction for general linear groups: the local non-tempered Gan–Gross–Prasad conjecture (non-Archimedean case). [arXiv:2006.02623](https://arxiv.org/abs/2006.02623), submitted
13. Chan, K.Y., Savin, G.: Bernstein–Zelevinsky derivatives: a Hecke algebra approach. *IMRN* (2019). <https://doi.org/10.1093/imrn/rnx138>
14. Chan, K.Y., Savin, G.: the Iwahori-component of the Gelfand–Graev representations. *Math. Z.* **288**, 125–133 (2018). <https://doi.org/10.1007/s00209-017-1882-3>
15. Chan, K.Y., Savin, G.: A vanishing Ext-branching theorem for $(GL_{n+1}(F), GL_n(F))$. *Duke Math. J.* [arXiv:1803.09131](https://arxiv.org/abs/1803.09131), to appear
16. Chan, K.Y., Savin, G.: Iwahori component of Bessel models. *Proc. Am. Math. Soc.* **148**, 1487–1497 (2020)
17. Gan, W.T., Gross, B.H., Prasad, D.: Symplectic local root numbers, central critical L values, and restriction problems in the representation theory of classical groups. *Sur les conjectures de Gross et Prasad. I. Asterisque* **346**, 1–109 (2012)

18. Gan, W.T., Gross, B.H., Prasad, D.: Branching laws for classical groups: the non-tempered case. *Compos. Math.* [arXiv:1911.02783](https://arxiv.org/abs/1911.02783), to appear
19. Gurevich, M.: On restriction of unitarizable representations of general linear groups and the non-generic local Gan-Gross-Prasad conjecture. *JEMS* [arXiv: 1808.02640](https://arxiv.org/abs/1808.02640), to appear
20. Haines, T.: The stable Bernstein center and test functions for Shimura varieties. In: *Automorphic forms and Galois Representations*, vol. 2, London Mathematical Society Lecture Notes Series, vol. 415, pp. 118–186. London Mathematical Society, London (2014)
21. Jacquet, H., Shalika, J.: The Whittaker models of induced representations. *Pac. J. Math.* **109**(1), 107–120 (1983)
22. Lang, S.: Restriction of the oscillator representation to dual pairs: some projective cases. [arXiv:1711.10562](https://arxiv.org/abs/1711.10562)
23. Lapid, E., Mínguez, A.: On a determinantal formula of Tadić. *Am. J. Math.* **136**(1), 111–142 (2014)
24. Lapid, E., Mínguez, A.: On parabolic induction on inner forms of the general linear group over a non-Archimedean local field. *Sel. Math. New Ser.* **22**, 2347–2400 (2016)
25. Lusztig, G.: Affine Hecke algebras and their graded versions. *J. Am. Math. Soc.* **2**, 599–635 (1989)
26. Nori, M., Prasad, D.: On a duality theorem of Schneider–Stuhler. *Journal für die reine und angewandte Mathematik* **762**, 261–280 (2020)
27. Prasad, D.: On the decomposition of a representation of $GL(3)$ restricted to $GL(2)$ over a p -adic field. *Duke Math. J.* **69**, 167–177 (1993)
28. Prasad, D.: An Ext-analogue of branching laws. In: *Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018*. Vol. II. Invited lectures, pp. 1367–1392. World Scientific Publishing, Hackensack (2018)
29. Schneider, P.: U-Stuhler, representation theory and sheaves on the Bruhat–Tits building. *Inst. Hautes Etudes Sci. Publ. Math.* **85**, 97–191 (1997)
30. Sakellaridis, Y., Venkatesh, A.: Periods and harmonic analysis on spherical varieties. *Astérisque* **396** (2017)
31. Sun, B., Zhu, C.B.: Multiplicity one theorems: the Archimedean case. *Ann. Math.* (2) **175**(1), 23–44 (2012)
32. Tadić, M.: On the classification of irreducible unitary representations of $GL(n)$ and the conjectures of Bernstein and Zelevinsky. *Ann. Sci. École Norm. Sup.* **19**, 335–382 (1986)
33. Tadić, M.: Unitary representations of $GL(n)$, derivatives in the non-Archimedean case, V. *Mathematikertreffen Zagreb-Graz (Mariatrost/Graz, 1986)*, Ber. Math.-Statist. Sect. Forschungsgesellsch. Joanneum, vol. 274, Forschungszentrum Graz, Graz, pp. Ber. No. 281, 19 (1987)
34. Tadić, M.: On the reducibility points beyond the ends of complementary series of p -adic general linear groups. *J. Lie Theory* **25**(1), 147–183 (2015)
35. Zelevinsky, A.: Induced representations of reductive p -adic groups II. *Ann. Sci. Ecole Norm. Sup.* **13**, 154–210 (1980)