

POLYNOMIAL RETRACTS AND THE JACOBIAN CONJECTURE

VLADIMIR SHPILRAIN AND JIE-TAI YU

ABSTRACT. Let $K[x, y]$ be the polynomial algebra in two variables over a field K of characteristic 0. A subalgebra R of $K[x, y]$ is called a retract if there is an idempotent homomorphism (a *retraction*, or *projection*) $\varphi : K[x, y] \rightarrow K[x, y]$ such that $\varphi(K[x, y]) = R$. The presence of other, equivalent, definitions of retracts provides several different methods of studying and applying them, and brings together ideas from combinatorial algebra, homological algebra, and algebraic geometry. In this paper, we characterize all the retracts of $K[x, y]$ up to an automorphism, and give several applications of this characterization, in particular, to the well-known Jacobian conjecture.

1. INTRODUCTION

Let $K[x, y]$ be the polynomial algebra in two variables over a field K of characteristic 0. A subalgebra R of $K[x, y]$ is called a *retract* if it satisfies any of the following equivalent conditions:

- (R1) There is an idempotent homomorphism (a *retraction*, or *projection*) $\varphi : K[x, y] \rightarrow K[x, y]$ such that $\varphi(K[x, y]) = R$.
- (R2) There is a homomorphism $\varphi : K[x, y] \rightarrow R$ that fixes every element of R .
- (R3) $K[x, y] = R \oplus I$ for some ideal I of the algebra $K[x, y]$.
- (R4) $K[x, y]$ is a projective extension of R in the category of K -algebras. In other words, there is a split exact sequence $1 \rightarrow I \rightarrow K[x, y] \rightarrow R \rightarrow 1$, where I is the same ideal as in (R3) above.

Examples. K ; $K[x, y]$; any subalgebra of the form $K[p]$, where $p \in K[x, y]$ is a *coordinate* polynomial (i.e., $K[p, q] = K[x, y]$ for some polynomial $q \in K[x, y]$). There are other, less obvious, examples of retracts: if $p = x + x^2y$, then $K[p]$ is a retract of $K[x, y]$, but p is not coordinate since it has a fiber $\{p = 0\}$ which is reducible, and therefore is not isomorphic to a line. Even less obvious examples are retracts generated by $p = xy$ or $p = x^2 - y^2$.

The very presence of several equivalent definitions of retracts shows how natural these objects are. Later on, we shall also comment on a very natural geometric meaning of retracts.

In [8], Costa has proved that every proper retract of $K[x, y]$ (i.e., one different from K and $K[x, y]$) has the form $K[p]$ for some polynomial $p \in K[x, y]$, i.e., is isomorphic to a polynomial K -algebra in one variable. A natural problem now is

Received by the editors March 11, 1997 and, in revised form, August 20, 1997.

1991 *Mathematics Subject Classification*. Primary 13B25, 13P10; Secondary 14E09, 16S10.

The second author's research was partially supported by RGC Fundable Grant 344/024/0002.

to somehow characterize those polynomials $p \in K[x, y]$ that generate a retract of $K[x, y]$. Since the image of a retract under any automorphism of $K[x, y]$ is again a retract, it would be reasonable to characterize retracts up to an automorphism of $K[x, y]$, i.e., up to a “change of coordinates”. We give an answer to this problem by proving the following:

Theorem 1.1. *Let $K[p]$ be a retract of $K[x, y]$. Then there is an automorphism ψ of $K[x, y]$ that takes the polynomial p to $x + y \cdot q$ for some polynomial $q = q(x, y)$. A retraction for $K[\psi(p)]$ is given then by $x \rightarrow x + y \cdot q; y \rightarrow 0$.*

Geometrically, our Theorem 1.1 says that (in case $K = \mathbf{C}$) every polynomial retraction of a plane is a “parallel” projection (sliding) on a fiber of a coordinate polynomial (which is isomorphic to a line) along the fibers of another polynomial (which generates a retract of $K[x, y]$).

Our proof of this result is based on the well-known Abhyankar-Moh theorem ([1], Main Theorem). We note in passing that Theorem 1.1 also yields a characterization of retracts of a free associative algebra $K\langle x, y \rangle$ (see Theorem 2.1 in the next section) if one uses a natural lifting. In Section 2, we also make an observation on retracts of a polynomial algebra in arbitrarily many variables (Proposition 2.2).

Theorem 1.1 yields another useful characterization of retracts of $K[x, y]$:

Corollary 1.2. *A polynomial $p \in K[x, y]$ generates a retract of $K[x, y]$ if and only if there is a polynomial mapping of $K[x, y]$ that takes p to x . The “if” part is actually valid for a polynomial algebra in arbitrarily many variables.*

Theorem 1.1 has several interesting applications, in particular, to the notorious

Jacobian conjecture ([13]). *If for a pair of polynomials $p, q \in K[x, y]$ the corresponding Jacobian matrix is invertible, then $K[p, q] = K[x, y]$.*

For a survey and background on this problem, the reader is referred to [5].

Now we establish a link between retracts of $K[x, y]$ and the Jacobian conjecture by means of the following:

Conjecture “R”. *If for a pair of polynomials $p, q \in K[x, y]$ the corresponding Jacobian matrix is invertible, then $K[p]$ is a retract of $K[x, y]$.*

This statement is formally much weaker than the Jacobian conjecture, since, instead of asking for p to be a coordinate polynomial, we only ask for p to generate a retract, and this property is much less restrictive as can be seen from our Theorem 1.1. However, the point is that these conjectures are actually equivalent:

Theorem 1.3. *Conjecture “R” implies the Jacobian conjecture.*

There are several different ways of proving Theorem 1.3 using the characterization of retracts given in Theorem 1.1; we believe that the proof we give here (using Newton polygons) is particularly simple.

As a corollary, we show that the Jacobian conjecture is equivalent to other (formally) weaker statements:

Corollary 1.4. *Each of the following claims is equivalent to the Jacobian conjecture. Suppose (p, q) is a Jacobian pair. Then:*

- (i) For some coordinate polynomial g , $K[x, y] = K[p] + \langle g \rangle$, where $\langle g \rangle$ is the ideal of $K[x, y]$ generated by the polynomial g .
- (ii) For some coordinate polynomial g , $K[x, y] = K[g] + \langle p \rangle$.

Corollary 3.1 in Section 3 also seems interesting, since it improves several known partial results on the two-variable Jacobian conjecture.

Conjecture “R” has the following geometric interpretation. Suppose we have a polynomial $p \in \mathbf{C}[x, y]$ which has a Jacobian mate. Then every fiber $\{p = c, c \in \mathbf{C}\}$ is a non-singular curve in the complex plane. Moreover, by a result of Kaliman [12], we can restrict our attention to the situation where every fiber of p is irreducible, i.e., is a connected curve.

If there is another polynomial fiber $\{h = c_0\}$ which is isomorphic to a line and intersects every fiber of p at exactly one point, then we can arrange a geometric projection of the plane onto the curve $\{h = c_0\}$ by sliding a point on a fiber $\{p = c\}$ toward the intersection point of $\{p = c\}$ with $\{h = c_0\}$. This geometric projection will also be algebraic, i.e., $\mathbf{C}[p]$ will be a retract of $\mathbf{C}[x, y]$ in this case.

The only problem is to show that there is a fiber (isomorphic to a line) which intersects every fiber of p at *exactly* one point. In particular, we have:

Corollary 1.5 (cf. [10]). *Suppose φ is a polynomial mapping of $\mathbf{C}[x, y]$ with invertible Jacobian matrix. If φ is injective on some line, then φ is an automorphism.*

We note here that the aforementioned result of Kaliman [12] calls for an example of a non-coordinate polynomial $p \in \mathbf{C}[x, y]$ that has a non-vanishing gradient and all of whose fibers are irreducible. A series of polynomials like that has been constructed in [4].

We also note that if we replaced the condition that the polynomial $p(x, y)$ have a Jacobian mate by the weaker condition that it have a non-vanishing gradient, then Conjecture “R” would not be true. The following counterexample has been communicated to us by A. van den Essen: $p(x, y) = x + (x + x^2y)^2$.

Another application of retracts to the Jacobian conjecture (somewhat indirect, though) is based on the “ φ^∞ -trick” familiar in combinatorial group theory (see [16]). For a polynomial mapping $\varphi : K[x, y] \rightarrow K[x, y]$, denote by $\varphi^\infty(K[x, y]) = \bigcap_{k=1}^\infty \varphi^k(K[x, y])$ the *stable image* of φ . Then we have:

Theorem 1.6. *Let φ be a polynomial mapping of $K[x, y]$. If the Jacobian matrix of φ is invertible, then either φ is an automorphism, or $\varphi^\infty(K[x, y]) = K$.*

Our proof of Theorem 1.6 is based on recent results of Formanek [11] and Connell-Zweibel [7]. Lemma 3.2, which is crucial for our proof of the theorem, seems to be of independent interest.

Obviously, if φ fixes a polynomial $p \in K[x, y]$, then $p \in \varphi^\infty(K[x, y])$. Therefore, we have:

Corollary 1.7. *Suppose φ is a polynomial mapping of $K[x, y]$ with invertible Jacobian matrix. If $\varphi(p) = p$ for some non-constant polynomial $p \in K[x, y]$, then φ is an automorphism.*

This yields the following interesting re-formulation of the Jacobian conjecture: if φ is a polynomial mapping of $K[x, y]$ with invertible Jacobian matrix, then for some automorphism α , the mapping $\alpha \cdot \varphi$ fixes a non-constant polynomial.

2. RETRACTS OF $\mathbf{K}\langle\mathbf{x}, \mathbf{y}\rangle$

We start with

Proof of Theorem 1.1. Let $K[p]$, $p \in K[x, y]$, be a retract of $K[x, y]$ (note that by a result of Costa [8], every retract of $K[x, y]$ has this form).

Let the corresponding retraction be given by $\phi : x \rightarrow q_1(p); y \rightarrow q_2(p)$ for some one-variable polynomials q_1, q_2 .

Since ϕ is a retraction, polynomials $q_1(p), q_2(p)$ should generate $K[p]$. Suppose q_1 and q_2 have degree $n \geq 1$ and $m \geq 1$, respectively. Then, by the Abhyankar-Moh theorem [1], either n divides m , or m divides n . Suppose $\deg(q_1) = k \cdot \deg(q_2)$ for some integer $k \geq 1$.

Then we make the following change of coordinates: $x \rightarrow \tilde{x} = x - c \cdot y^k; y \rightarrow \tilde{y} = y$, where the coefficient $c \in K^*$ is chosen so that in the polynomial $q_1 - c \cdot q_2^k$, the leading terms cancel out.

In these new coordinates, our retraction acts as follows: $\phi : \tilde{x} \rightarrow q_1 - c \cdot q_2^k = \tilde{q}_1; \tilde{y} \rightarrow q_2 = \tilde{q}_2$. The polynomials \tilde{q}_1 and \tilde{q}_2 are easily seen to be another generating set of $K[p]$, but the sum of degrees of \tilde{q}_1 and \tilde{q}_2 is less than that of q_1 and q_2 .

Continuing this process, we shall eventually arrive at a pair of polynomials one of which is a constant $c \in K$. Denote the other one by h ; then we must have $K[h] = K[p]$, i.e., $h = c_1 \cdot p + c_2$ for some $c_1 \in K^*, c_2 \in K$.

Thus, we have shown that for some automorphism $\psi \in \text{Aut}(K[x, y])$ (“change of coordinates”), the composition $\phi\psi$ takes x to $c_1 \cdot p + c_2$, and y to c . It follows that $c_1 \cdot \psi(p(x, y)) + c_2 = x + (y - c) \cdot q(x, y)$ for some polynomial $q(x, y)$. Applying a linear transformation completes the proof. \square

Proof of Corollary 1.2. (1) Suppose $p \in K[x, y]$ generates a retract of $K[x, y]$. Then, by Theorem 1.1, for some automorphism $\psi \in \text{Aut}(K[x, y])$, the polynomial $\psi(p)$ has the form $x + y \cdot q(x, y)$. Let ϕ be a mapping of $K[x, y]$ that takes x to x ; y to 0. Then $\phi(\psi(p)) = x$.

(2) We are going to prove the “if” part for a polynomial algebra $K[x_1, \dots, x_n]$ in arbitrarily many variables.

Let $\varphi(p) = x_1$. Consider the following mapping of $K[x_1, \dots, x_n]$: $\psi : x_1 \rightarrow p; x_i \rightarrow 0, i = 2, \dots, n$. Then

$$(1) \quad \psi(\varphi(p)) = p.$$

Denote $\varrho = \psi\varphi$. Then, by (1), $\varrho(p) = p$, which means ϱ fixes every element of $K[p]$. Also, it is clear that $\varrho(K[x_1, \dots, x_n]) = K[p]$. Therefore, ϱ is a retraction of $K[x_1, \dots, x_n]$, and $K[p]$ a retract. \square

Now we give a characterization of retracts of a free associative algebra $K\langle x, y \rangle$:

Theorem 2.1. *Let R be a proper retract of $K\langle x, y \rangle$. There is an automorphism ψ of $K\langle x, y \rangle$ that takes R to $K\langle v \rangle = K[v]$ for some element v of the form $x + w(x, y)$, where $w(x, y)$ belongs to the ideal of $K\langle x, y \rangle$ generated by y .*

Proof. First of all, every element of the given form generates a retract of $K\langle x, y \rangle$. Indeed, the corresponding retraction is given by $x \rightarrow v; y \rightarrow 0$.

Now we are going to show that every retract of $K\langle x, y \rangle$ has the form $K\langle v \rangle = K[v]$ for some element $v \in K\langle x, y \rangle$. From the definition (R1) of a retract, we see that every retract of $K\langle x, y \rangle$ can be generated by two elements $\varphi(x)$ and $\varphi(y)$.

Another easy observation is that if R is a retract of $K\langle x, y \rangle$ with the corresponding retraction ϕ , then R^α is a retract of $K[x, y]$, and the corresponding retraction is ϕ^α . Here R^α denotes the image of R under the natural abelianization mapping $\alpha : K\langle x, y \rangle \rightarrow K[x, y]$ (note that the kernel of this mapping is the commutator ideal of $K\langle x, y \rangle$), and ϕ^α is the natural abelianization of ϕ . The best way to see it is to apply the definition (R2) of a retract.

Upon combining these two observations with what we know about retracts of $K[x, y]$, we see that generators of our retract R must be of the form $v_1(x, y) = q_1(x, y) + w_1(x, y)$ and $v_2(x, y) = q_2(x, y) + w_2(x, y)$, where w_1, w_2 belong to the commutator ideal of $K\langle x, y \rangle$, and $q_1^\alpha = h_1(p), q_2^\alpha = h_2(p)$, where $p = p(x, y) \in K[x, y]$ is a polynomial that generates a retract of $K[x, y]$.

Moreover, since R^α should be equal to $K[p]$, we should have $K[h_1(p), h_2(p)] = K[p]$, so that we are in a position to apply the Abhyankar-Moh theorem. Repeating the argument from the proof of Theorem 1.1, we see that after applying an appropriate automorphism $\psi \in \text{Aut}(K[x, y])$ (“change of coordinates”), our pair $(h_1(p), h_2(p))$ becomes $(\psi(h_1(p)), 0)$.

Since every automorphism of $K[x, y]$ can be lifted to an automorphism of $K\langle x, y \rangle$ (see [6], Theorem 8.5), we have shown that, after applying an automorphism if necessary, our retract R can be generated by two elements, of which one (call it u) belongs to the commutator ideal of $K\langle x, y \rangle$, and the other one (call it v) is of the form $x + y \cdot q(x, y) + w(x, y)$, where $w(x, y)$ belongs to the commutator ideal of $K\langle x, y \rangle$ (which is contained in the ideal of $K\langle x, y \rangle$ generated by y), and the corresponding retraction is $\phi : x \rightarrow v; y \rightarrow u$.

Then, since ϕ^α annihilates y^α (see above), $u = \phi(y)$ should belong to the commutator ideal of $K\langle x, y \rangle$.

On the other hand, ϕ should fix every element of R , in particular, the element u . Suppose $u \neq 0$. Write u in the form $u = m_1 + \tilde{u}$, where m_1 is the sum of the lowest degree terms, i.e., any monomial of \tilde{u} has degree greater than that of monomials in m_1 . Then, since u belongs to the commutator ideal of $K\langle x, y \rangle$, every monomial in m_1 depends on y . Therefore, the image of any monomial in m_1 under the endomorphism ϕ is a sum of monomials whose degree is greater than that of monomials in m_1 (this sum might be equal to zero, but it does not affect the argument). The same applies to monomials of \tilde{u} .

Thus, there is no way we can have $\phi(u) = u$; this contradiction shows that $u = 0$, which completes the proof of the theorem. \square

We conclude this section with an observation on retracts of a polynomial algebra in arbitrarily many variables.

Proposition 2.2. *Let R be a proper retract of $K[x_1, \dots, x_n]$ generated by polynomials p_1, \dots, p_n , $n \geq 2$. Then p_1, \dots, p_n are algebraically dependent.*

Proof. Let $\phi : K[x_1, \dots, x_n] \rightarrow R$ be a retraction, so that $\phi(R) = R$. In particular, ϕ restricted to R is an automorphism of R . If the polynomials p_1, \dots, p_n were algebraically independent, then a result of [3] and [7] would imply that ϕ is an automorphism of $K[x_1, \dots, x_n]$ as well. In that case, we would have $R = K[x_1, \dots, x_n]$, which means R is not a proper retract, hence a contradiction. \square

We note that it is an open problem whether or not any retract of $K[x_1, \dots, x_n]$, $n \geq 3$, can be generated by algebraically independent polynomials. This problem

is related to (some forms of) the well-known *cancellation problem* – see [8] for discussion.

3. THE JACOBIAN CONJECTURE

First we prove that our Conjecture “R” implies the Jacobian conjecture.

Proof of Theorem 1.3. Let $\varphi : x \rightarrow p(x, y); y \rightarrow q(x, y)$ be a polynomial mapping of $K[x, y]$ with invertible Jacobian matrix.

If we assume that our Conjecture “R” is true, then $p(x, y)$ generates a retract of $K[x, y]$; hence by Theorem 1.1, for some automorphism $\psi \in \text{Aut}(K[x, y])$, the polynomial $\psi(p)$ has the form $x + y \cdot h(x, y)$. Therefore, upon combining φ with ψ if necessary, we may assume that $p(x, y)$ itself has this form.

Now we appeal to a result of [3] and [15] concerning Newton polygons of a Jacobian pair $(p(x, y), q(x, y))$. The Newton polygon of a polynomial $f = f(x, y) = \sum a_{ij}x^i y^j$ is the convex hull of $\{(i, j) \mid a_{ij} \neq 0\} \cup \{(0, 0)\}$.

The result of [15] we need is that if $(p(x, y), q(x, y))$ is a Jacobian pair, but $\varphi : x \rightarrow p(x, y); y \rightarrow q(x, y)$ is not an automorphism, then the Newton polygons of $p(x, y)$ and $q(x, y)$ are radially similar.

Now, by way of contradiction, suppose φ is not an automorphism. Look at the Newton polygon of $p(x, y)$. We see that it has an edge of length 1, namely, the one between the vertices $(0, 0)$ and $(1, 0)$. It follows that the similarity ratio for the Newton polygons of $p(x, y)$ and $q(x, y)$ is an integer; in particular, $q(x, y)$ has the form $\sum_{i=1}^k c_i x^i + y \cdot f(x, y)$ for some $c_i \in K$, $c_k \neq 0$.

Then replace the pair (p, q) with $(p, q - c_k p^k)$. This new pair clearly has the same properties as (p, q) does: it is a Jacobian pair, but the corresponding mapping is not an automorphism. However, the highest degree of monomials of the form x^m in the second polynomial has been decreased.

Therefore, we can repeatedly apply our argument (note that $p(x, y)$ does not change), until we get a pair (p, g) , where g has no monomials of the form x^m . But in that case, the Newton polygon of g has no edges along the x -axis; hence the Newton polygons of p and g cannot be radially similar. This contradiction completes the proof of Theorem 1.3. \square

Proof of Corollary 1.4. (i) Without loss of generality, we may assume that both p and g have zero constant term. Moreover, upon applying an automorphism to all polynomials under consideration if necessary, we may assume $g = x$.

First we show that if $K[x, y] = K[p] + \langle x \rangle$, then the sum is actually direct.

By way of contradiction, suppose we have

$$(2) \quad x \cdot u = \sum_{i=1}^m c_i \cdot p^i$$

for some non-zero polynomial $u = u(x, y)$ and constants $c_i \in K$. Since the left-hand side of (2) is divisible by x , the right-hand side should be divisible by x , too. This is only possible if p itself is divisible by x , but in that case, the Newton polygon of p would not have an edge along the y -axis, which contradicts p having a Jacobian mate (see e.g. [15]).

Thus, we have shown that $K[x, y] = K[p] \oplus \langle x \rangle$. By the definition (R3) of a retract, this implies $K[p]$ is a retract of $K[x, y]$. Therefore, by our Theorem 1.3, $K[p, q] = K[x, y]$.

(ii) Arguing as in (i), we get $p \cdot u = \sum_{i=1}^m c_i \cdot x^i$. In this case, p cannot depend on y , so $p = p(x)$; but then the equality $K[x, y] = K[x] + \langle p \rangle$ is not possible.

Thus, $K[x, y] = K[x] \oplus \langle p \rangle$. This clearly implies $p = c \cdot y + f(x)$ for some $c \in K^*$ and $f(x) \in K[x]$. The result follows. \square

Corollary 3.1. *Let K be an algebraically closed field, and let $(p, q), p, q \in K[x, y]$, be a Jacobian pair. Suppose p has the form $p = x + g$ for some $g \in K[x, y]$, which is divisible by a homogeneous polynomial. Then $K[p, q] = K[x, y]$.*

Proof. If p is linear, then we are done. Suppose p is non-linear, and suppose g is divisible by a non-constant homogeneous polynomial h .

If h is divisible by x , then p itself is divisible by x . In that case, the Newton polygon of p does not have an edge along the y -axis, which contradicts p having a Jacobian mate (see [15]) unless $p = x$.

If h is divisible by y , then $K[p]$ is a retract of $K[x, y]$. If not, then there is $c \in K^*$ such that the homomorphism $x \rightarrow p; y \rightarrow cp$ takes h to 0. This is obviously a retraction, so again, $K[p]$ is a retract of $K[x, y]$.

Applying our Theorem 1.3 yields the result. \square

Van den Essen and Tutaj [9] have shown that if a Jacobian pair (p, q) is of the form $(x + h_1, y + h_2)$, where both h_1 and h_2 are homogeneous polynomials, then $K[p, q] = K[x, y]$. It is notable that our Corollary 3.1 not only relaxes the condition on the form of polynomials, but, most importantly, our condition is imposed on *one* polynomial only.

To prove Theorem 1.6, we need the following result, which is also of independent interest:

Lemma 3.2. *Let $\varphi : x \rightarrow p(x, y); y \rightarrow q(x, y)$ be a polynomial mapping of $K[x, y]$ with invertible Jacobian matrix. Suppose $\varphi(K[x, y])$ contains a coordinate polynomial. Then φ is an automorphism.*

Proof. Upon composing φ with an automorphism if necessary, we may assume that $x \in \varphi(K[x, y])$. This implies $K[p, q, x] = K[p, q]$, and, therefore, $K(p, q, x) = K(p, q)$, where $K(p, q)$ is the quotient field of $K[p, q]$.

On the other hand, by a result of [11], (p, q) being a Jacobian pair implies $K(p, q, x) = K(x, y)$.

Therefore, we have $K(p, q) = K(x, y)$, which by Keller’s theorem [13] implies $K[p, q] = K[x, y]$. \square

Proof of Theorem 1.6. Suppose φ is *not* an automorphism. Then, by Lemma 3.2, $\varphi(K[x, y])$ contains no coordinate polynomials. In particular, the degree of any non-constant polynomial in $\varphi(K[x, y])$ is at least 2. By an inductive argument, we show now that the degree of any non-constant polynomial in $\varphi^k(K[x, y])$ is at least $(k + 1)$; this will imply $\varphi^\infty(K[x, y]) = K$.

Consider an algebra $\varphi^k(K[x, y])$ for some $k \geq 1$. Since φ is injective (this is ensured by the Jacobian condition), the polynomials $\varphi^k(x)$ and $\varphi^k(y)$ are algebraically independent.

If φ restricted to $\varphi^k(K[x, y])$ were an automorphism of $\varphi^k(K[x, y])$, then, by a result of [7], φ would be an automorphism of $K[x, y]$, contrary to our assumption. (Note that $\varphi^k(K[x, y])$ is invariant under φ since, by induction, $\varphi^k(K[x, y]) \subseteq \varphi^{k-1}(K[x, y])$ implies $\varphi^{k+1}(K[x, y]) \subseteq \varphi^k(K[x, y])$.) Hence, $\varphi|_{\varphi^k(K[x, y])}$ is not an automorphism of $\varphi^k(K[x, y])$, and our previous argument yields that $\varphi^{k+1}(K[x, y])$ contains no coordinate polynomials of $\varphi^k(K[x, y])$. In particular, the degree of any

polynomial in $\varphi^{k+1}(K[x, y])$ is greater than that of any coordinate polynomial in $\varphi^k(K[x, y])$. This completes the induction. \square

A. van den Essen has pointed out to us that in the case $K = \mathbf{C}$, a more complicated (geometric) proof of Theorem 1.6 was given by Kraft [14], who also proved that if a polynomial mapping φ of $\mathbf{C}[x, y]$ is *not* birational (i.e., φ does not induce an automorphism of the quotient field $\mathbf{C}(x, y)$), then the stable image of φ is a retract of $\mathbf{C}[x, y]$. This yields a natural question – is the same true for *any* polynomial mapping? (cf. [16]). \square

Acknowledgements. We are indebted to A. Campbell and D. Markushevich for numerous insightful discussions. The first author is grateful to the Department of Mathematics of the University of Hong Kong for its warm hospitality during his visit when this work was initiated. A crucial part of this work was done when the first author was visiting the Mathematical Sciences Research Institute in Berkeley, and the second author was visiting the Max Planck Institut für Mathematik in Bonn. The warm hospitality and stimulating atmosphere of these institutions are greatly appreciated.

REFERENCES

- [1] S.S. Abhyankar, T.-T. Moh, *Embeddings of the line in the plane*, J. Reine Angew. Math. **276** (1975), 148–166. MR **52**:407
- [2] S.S. Abhyankar, *Lectures on expansion techniques in algebraic geometry*, Notes by Balwant Singh. Tata Institute of Fundamental Research Lectures on Mathematics and Physics, **57**. Tata Institute of Fundamental Research, Bombay, 1977. MR **80m**:14016
- [3] H. Appelgate, H. Onishi, *The Jacobian conjecture in two variables*, J. Pure Appl. Algebra **37** (1985), 215–227. MR **87b**:14005
- [4] E. Artal-Bartolo, P. Cassou-Nogues, I. Luengo Velasco, *On polynomials whose fibers are irreducible with no critical points*, Math. Ann. **299** (1994), 477–490. MR **95g**:14016
- [5] H. Bass, E. Connell and D. Wright, *The Jacobian conjecture: reduction of degree and formal expansion of the inverse*, Bull. Amer. Math. Soc. **7** (1982), 287–330. MR **83k**:14028
- [6] P.M. Cohn, *Free rings and their relations*, Second edition, Academic Press, London, 1985. MR **87e**:16006
- [7] E. Connell, J. Zweibel, *Subrings invariant under polynomial maps*, Houston J. Math. **20** (1994), 175–185. MR **95g**:13006
- [8] D. Costa, *Retracts of polynomial rings*, J. Algebra **44** (1977), 492–502.
- [9] A. van den Essen, H. Tutaj, *A remark on the two-dimensional Jacobian conjecture*, J. Pure Appl. Algebra **96** (1994), 19–22. MR **55**:2876
- [10] J. Gwozdziwicz, *Injectivity on one line*, Bull. Soc. Sci. Lett. Lodz Ser. Rech. Deform. **15** (1993), 59–60. MR **95f**:14024
- [11] E. Formanek, *Observations about the Jacobian conjecture*, Houston J. Math. **20** (1994), 369–380. MR **95f**:14027
- [12] S. Kaliman, *On the Jacobian conjecture*, Proc. Amer. Math. Soc. **117** (1993), 45–51. MR **93e**:14017
- [13] O. Keller, *Ganze Cremona-Transformationen*, Monatsh. Math. Phys. **47** (1939), 299–306.
- [14] H. Kraft, *On a question of Yosef Stein*, Automorphisms of affine spaces (Curacao, 1994), 225–229, Kluwer Acad. Publ., Dordrecht, 1995. MR **96i**:14013
- [15] J. Lang, *Newton polygons of Jacobian pairs*, J. Pure Appl. Algebra **72** (1991), 39–51. MR **92i**:14012
- [16] E.C. Turner, *Test words for automorphisms of free groups*, Bull. London Math. Soc. **28** (1996), no. 3, 255–263. MR **96m**:20039

DEPARTMENT OF MATHEMATICS, THE CITY COLLEGE OF NEW YORK, NEW YORK, NEW YORK 10031

E-mail address: shpil@groups.sci.ccny.cuny.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HONG KONG, POKFULAM ROAD, HONG KONG

E-mail address: yujt@hkusua.hku.hk