

26 **Abstract:**

27 BACKGROUND: Heterotrimeric G proteins take part in membrane-mediated cell-signalling
28 and have a role in e.g. hormonal regulation. This study clarifies the expression and
29 localization of the G protein subunit $G\alpha_{i2}$ in the human endometrium and fallopian tube and
30 changes in $G\alpha_{i2}$ expression in human endometrium during the menstrual cycle. METHODS:
31 The expression of $G\alpha_{i2}$ was identified by PCR, and localization confirmed by immunostaining.
32 Cyclic changes in $G\alpha_{i2}$ expression during the menstrual cycle were evaluated by quantitative
33 real time PCR. RESULTS: We found $G\alpha_{i2}$ to be expressed in human endometrium, fallopian
34 tube tissue and fallopian tube primary epithelial cells. Our studies revealed enriched
35 localization of $G\alpha_{i2}$ in human fallopian tube cilia and in endometrial glands. We showed that
36 $G\alpha_{i2}$ expression in human endometrium changes significantly during the menstrual cycle.
37 CONCLUSIONS: $G\alpha_{i2}$ is specifically localized in oviductal cilia of rat and human and is
38 likely to have a cilia-specific role in reproduction. Significantly variable expression of $G\alpha_{i2}$
39 during the menstrual cycle suggests it might be under hormonal regulation in the female
40 reproductive tract *in vivo*.

41

42

43

44

45

46

47

48

49

50 Introduction

51

52 Among the cell-surface receptors, G protein-coupled receptors are the most widespread and
53 diverse, playing an essential regulatory role in cell growth, hormonal regulation, sensory
54 perception and neuronal activity (Hepler and Gilman, 1992). In reproduction, G protein-
55 coupled receptors have a neuroendocrine regulatory role in gonadotropin-releasing hormone
56 (GnRH) -induced secretion of luteinising hormone (LH) and follicle-stimulating hormone
57 (FSH) from the anterior pituitary gland (Chi *et al.*, 1993; Tsutsumi *et al.*, 1992). In gonads, G
58 protein-coupled receptors mediate gonadotropin signalling (Loosfelt *et al.*, 1989; McFarland
59 *et al.*, 1989; Minegishi *et al.*, 1991; Minegishi *et al.*, 1990; Sprengel *et al.*, 1990), thus
60 regulating the synthesis and secretion of sex hormones.

61

62 G protein-coupled receptors communicate via heterotrimeric G proteins, which are recognized
63 as crucial elements in various types of membrane-mediated cell-signalling. Heterotrimeric G
64 proteins consist of α -, β - and γ -subunits. According to the α -subunits, G proteins are divided
65 into four classes (G_s , G_i , G_q and G_{12}) (Hepler and Gilman, 1992). Proteins of the G_i family are
66 the most diverse and interact with a wide variety of G protein-coupled receptors. For example,
67 they take part in hormonal regulation via interaction with GnRH (Hawes *et al.*, 1993;
68 Krsmanovic *et al.*, 2003; Krsmanovic *et al.*, 2001; Stanislaus *et al.*, 1998), FSH (Arey *et al.*,
69 1997) and LH receptors (Herrlich *et al.*, 1996). Moreover, G_i family proteins play a role in the
70 signal transduction of rapid, nongenomic actions of estrogen (Benten *et al.*, 2001) and
71 progesterone (Karteris *et al.*, 2006; Zhu *et al.*, 2003).

72

73 The dual balance between G_i and G_s signalling in the regulation of adenylyl cyclase has been
74 well established. Proteins of G_i -family can inhibit adenylyl cyclase (AC) and thus decrease

75 intracellular cAMP concentration (Bokoch *et al.*, 1984; Katada *et al.*, 1984). Via this pathway,
76 G_i-family protein G α_{i2} has been shown to take part in adrenergic signalling, controlling
77 myometrium relaxation in the rat during pregnancy (Mhaouty *et al.*, 1995). In the human
78 myometrium, the levels of G α_{i2} have been shown to decrease during pregnancy, suggesting
79 that the consequent, altered balance between G α_{i2} and G_s could be responsible for maintaining
80 the relaxation of uterus during pregnancy (Europe-Finner *et al.*, 1993). Although the role of
81 G α_{i2} in myometrium has been thoroughly studied, the presence or the role of G α_{i2} elsewhere
82 in the human reproductive tract remains unclear.

83

84 Immunohistochemical studies in the rat have shown that G α_{i2} is specifically localized in
85 tissues having motile cilia with a characteristic 9+2 ultrastructure. Such a specific localization
86 in rat oviductal, tracheal and brain ependymal cilia (Shinohara *et al.*, 1998) implies that G α_{i2}
87 may well serve a physiological function distinct from those of the other G α subunits. It is
88 probable that G α_{i2} might play a cilia-specific physiological role. Interestingly, proteomic
89 analysis has revealed G α_{i2} as a resident axonemal protein of the human bronchial cilia
90 (Ostrowski *et al.*, 2002). To date, however, there are no reports providing evidence of the
91 localization of G α_{i2} in any other human ciliated tissues, such as fallopian tubes. In this study,
92 we identify the presence and localization of G α_{i2} in tissues which are primarily in contact
93 with gametes, and provide environment for fertilization, early development of the embryo as
94 well as implantation, i.e., the human fallopian tube and endometrium. We have also evaluated
95 the potential changes in G α_{i2} expression in human endometrium during the menstrual cycle to
96 reveal any potential hormonal regulation of this G protein subunit in humans.

97

98

99

100 **Materials and methods**

101

102 **Endometrial tissue collection and preparation for immunohistochemistry**

103 The current study was approved by the Local Ethics Committee and informed written consent
104 was obtained prior to the collection of tissue samples. For immunohistochemical

105 investigations, tissue samples were obtained from 6 fertile women, and for genomic studies,

106 endometrial biopsies were obtained from 21 fertile women. All the women taking part in the

107 investigation had regular cycles, showed no evidence of any pathological uterine disorder, and

108 had not used oral contraception or an intrauterine device during the previous three months.

109 Biopsies were obtained in the operating theatre between 2 and 29 days after the last menstrual

110 period (LMP). The mean age of the women taking part in the study was 35 (range 24-40)

111 years, and each had had at least one previous successful pregnancy.

112

113 Endometrial biopsies for immunohistochemistry were immediately snap-frozen and stored in
114 liquid nitrogen until processed. Cryosections were cut at 5 μm and stored at -70°C until use.

115 For genomic studies, endometrial biopsies were immediately placed in RNAlater (Ambion,

116 Huntingdon, U.K.), followed by immersion in liquid nitrogen until processed.

117

118 **Fallopian tube tissue collection and preparation for immunohistochemistry**

119 Human fallopian tube tissues were collected from 9 patients undergoing total abdominal

120 hysterectomy for benign gynaecological conditions. The mean age of the women taking part

121 in the study was 42 (range 33-56) years.

122

123 Fallopian tube tissue samples for immunohistochemistry were immediately fixed in 10%

124 formalin overnight and embedded in paraffin. Paraffin sections were cut at 5 μm . For genomic

125 studies, fallopian tube tissue samples were immediately placed in RNAlater (Ambion), and
126 stored for 24 hours at 4°C followed by immersion and storage in liquid nitrogen until
127 processed.

128

129 **Cell culture**

130 Fallopian tube tissue samples for primary epithelial cell cultures were obtained as follows:

131 fallopian tubes were placed in Hank's solution immediately after collection, cut open

132 longitudinally and incubated 1 h with 0.25 % collagenase (at 37°C, 95% O₂, 5% CO₂). The

133 cells were scraped gently using a sterile blade, washed with red blood cell lysing buffer

134 (Sigma-Aldrich) and then 2-3 times with culture media (DMEM-F12). The cells were plated

135 into 75 ml flasks. Fallopian tube primary epithelial cells were cultured at +37°C in DMEM

136 (F12) culture media (Invitrogen, Paisley, UK) supplemented with 1% penicillin and

137 streptomycin (Sigma-Aldrich), 10% fetal calf serum (Invitrogen) and L-glutamine (Invitrogen)

138 in 5% CO₂ atmosphere.

139

140 **RNA isolation and cDNA synthesis**

141 Tissues were removed from RNAlater and homogenised in 3 ml of TRIreagent (Sigma-

142 Aldrich) using an Ultra-Turrax homogenizer for 2 min. Total RNA from the tissues and

143 pelleted cells stored in TRIreagent was extracted following standard protocol supplied by the

144 manufacturer. Total RNA was treated with Dnase I (DNA-freeTM, Ambion) to remove

145 genomic DNA contamination from the samples. First strand cDNA synthesis was performed

146 using oligo dT primers (Metabion, Martinsried, Germany) and reverse transcription by

147 SuperScript II (200 U/μl, Invitrogen, Paisley, UK). Negative controls were prepared without

148 the enzyme (non-reverse transcribed controls, RT controls).

149

150 PCR

151 PCR was performed with the constructed cDNAs, Platinum Blue PCR Super Mix (Invitrogen)
152 and primers from Metabion. We used the following primer pairs: β -actin forward 5'-TGA
153 CCC AGA TCA TGT TTG AGA CC-3' and β -actin reverse 5'-GGA GGA GCA ATG ATC
154 TTG ATC TTC-3', $G\alpha_{i2}$ forward 5'-CTT GTC TGA GAT GCT GGT AAT GG-3' and $G\alpha_{i2}$
155 reverse 5'-CTC CCT GTA AAC ATT TGG ACT TG-3'. The amplification was run for 35
156 cycles under the following conditions: 95° 30 sec, 58° or 65° 30 sec, 72° 30 sec. Amplified
157 sequences were 643 and 212 base pairs for $G\alpha_{i2}$ and β -actin respectively. Annealing
158 temperatures of 58° (β -actin) and 65° ($G\alpha_{i2}$) were used. All experiments included RT controls
159 as well as negative controls (no cDNA). PCR products were separated on 1.2 % agarose gel.

160

161 Quantitative real time PCR

162 Quantitative real time PCR was performed with the constructed cDNAs and the same primers
163 that were used in PCR reactions. SYBR Green Jump Start (Sigma-Aldrich) master mix
164 (containing 10 μ l SYBR Green, 7 μ l Water, 1 μ l of each primer and 1 μ l cDNA) was added to
165 each well of PCR plate and amplification was performed under the following conditions: 50
166 cycles (95° 30 sec, 58° or 65° 30 sec, 72° 30 sec). All experiments included RT controls and
167 negative controls (no cDNA).

168

169 Results were analyzed using iCycler (Biorad laboratories Ltd, Hemel Hempstead, UK). To
170 compare relative quantities of $G\alpha_{i2}$ expression during the menstrual cycle, endometrial
171 biopsies were divided into three groups; menstrual (LMP + 1-4 ; n = 3; LMP +1, +4 and +4),
172 proliferative (LMP + 5-14 ; n = 9; early proliferative LMP +5, +5 and +7, mid-proliferative
173 LMP +8, +9 and +10, late proliferative LMP +11, +12 and +13) and secretory (LMP + 15-29 ;
174 n = 9; early secretory LMP +16, +16 and +17, mid-secretory LMP +20, +21 and +22, late

175 secretory LMP +26, +28 and +29). Relative $G\alpha_{i2}$ expression quantities were compared
176 between these groups. The threshold cycle values were normalised against threshold value of
177 human β -actin. The results were expressed as mean \pm S.E.M. Statistical analysis was
178 performed by using one-way ANOVA with Tukey's multiple comparison test.
179 $p < 0.05$ was considered significant.

180

181 **Immunohistochemistry**

182 Cryosections of endometrium were thawed by immersion (15 min at 20 °C) into fixative
183 containing 4 % paraformaldehyde (Sigma-Aldrich, Poole, UK) in 0.1 M PBS, pH 7.4. The
184 slides were then washed with PBS (2x5 min), and further fixed by immersion in -20°C
185 methanol (4 min) followed immediately by treatment with -20°C acetone (2 min). After 2x5
186 min washes with PBS, endogenous peroxidase activity was blocked by 5% H_2O_2 (in distilled
187 water) treatment (5 min). The slides were then washed with deionized water (2x5 min) and
188 PBS (2x5 min). After this, the protocol follows the same blocking and staining protocol as
189 described for paraffin sections.

190

191 Fallopian tube paraffin sections were firstly dewaxed in xylene, rehydrated through a series of
192 ethanols and finally washed with PBS. Endogenous peroxidase activity was quenched by a 20
193 min incubation with 3% H_2O_2 (v/v) in methanol. Antigen retrieval was performed by
194 microwave irradiation in 10mM citrate buffer, pH 6.0 (12 min). The slides were allowed to
195 cool in the buffer and then washed with PBS (2x3 min).

196

197 Vectastain Elite ABC Kit (Vector Laboratories, Peterborough, UK) was used according to the
198 manufacturers instructions for both cryosections and paraffin sections, with the following
199 modifications. Slides were blocked in blocking buffer containing 250 μ l avidin D / ml (1 h

200 RT). Mouse anti-G_iα-2 monoclonal antibody, MAB3077 (Chemicon International, Temecula,
201 CA) was diluted into Dako antibody diluent (Dako UK Ltd, Cambridgeshire, UK) containing
202 250 µl biotin / ml, and incubated overnight at 4 °C (cryosections 1:1000, paraffin sections
203 1:500). Primary antibody was omitted in negative controls. The slides were washed with PBS
204 (5 min), and incubated with secondary antibody (1:200 Biotinylated anti-mouse (Vector
205 Laboratories)) for 30 min at 20 °C. The slides were washed as before and incubated for 30
206 min with Vectastain ABC reagent (Vector Laboratories). After washing, binding was
207 visualized by incubation with substrate DAB or DAB-Ni for 8 min (Vector Laboratories). The
208 slides were rinsed with tap water (5 min) and PBS (3 min) and counterstained by using 10%
209 haematoxylin (10 min). Following thorough rinse in tap water, slides were dehydrated
210 through a series of ethanols, cleared in xylene and coverslipped with DePex mounting
211 medium (VWR International, Lutterworth, UK).

212

213 The endometrial biopsy specimens were timed according to LMP and morphology and
214 divided into three groups, menstrual, proliferative or secretory. The slides were imaged using
215 a x40 objective on an Olympus CKX41 microscope. Digital images were captured with a
216 Nikon Coolpix 5400 camera and identically edited in Adobe Photoshop (Adobe Systems,
217 Mountain View, CA).

218

219 **Results**

220

221 *PCR reveals the expression of $G\alpha_{i2}$ gene in human reproductive tissues.*

222 We used human fallopian tube tissue and human endometrial biopsies to study the expression
223 of $G\alpha_{i2}$ by PCR. Our data revealed that $G\alpha_{i2}$ is expressed in human fallopian tube and human
224 endometrium (Figure 1 A, B). Our studies also confirmed that $G\alpha_{i2}$ is expressed in primary
225 cultures of fallopian tube epithelial cells (Figure 1 C). Control experiments with non-reverse
226 transcribed RNA of each sample confirmed that there was no contamination of human DNA
227 in the samples.

228

229 *Immunohistochemistry shows specific localization of $G\alpha_{i2}$ protein in fallopian tube cilia and*
230 *enrichment in endometrial glands.*

231 Immunostaining on human fallopian tube paraffin sections showed specific localization of
232 $G\alpha_{i2}$ protein in fallopian tube epithelial cells and the cilia (Figure 2 C). Positive staining was
233 also seen in the cytoplasm of epithelial cells, surrounding the nuclei. In endometrial tissue,
234 $G\alpha_{i2}$ staining was enriched in endometrial glands, but was present also in stroma (Figure 2 A,
235 B).

236

237 *Quantitative real time PCR shows alterations in $G\alpha_{i2}$ gene expression during the menstrual*
238 *cycle.*

239 We carried out quantitative real time PCR experiment on endometrial biopsies spanning the
240 menstrual cycle (Figure 3). Based on the phase of the menstrual cycle of each patient, the
241 biopsies were designated in three groups, namely menstrual (LMP + 1-4), proliferative (LMP
242 + 5-14) and secretory (LMP + 15-29).

243

244 Our results demonstrated that endometrial expression of $G\alpha_{i2}$ gene changed during the cycle.
245 The expression reached its peak in secretory phase. The expression of $G\alpha_{i2}$ gene in secretory
246 phase was significantly higher ($p < 0.05$) compared to that of the other phases.

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268 **Discussion**

269

270 The present study demonstrates the existence and localization of $G\alpha_{i2}$ in human endometrium
271 and fallopian tube. Our data establishes the specific localization of $G\alpha_{i2}$ in the fallopian tube
272 epithelial cells, particularly in the cilia of fallopian tube epithelial cells. In human
273 endometrium, we have demonstrated that localization of $G\alpha_{i2}$ is enriched in endometrial
274 glands. We have also shown that $G\alpha_{i2}$ expression in human endometrium changes
275 significantly during the menstrual cycle with maximum expression in the secretory phase,
276 providing evidence that expression of this G_i subunit might be under hormonal regulation in
277 the female reproductive tract *in vivo*.

278

279 The presence of G protein subunit $G\alpha_{i2}$ in rat myometrial membranes was first reported by
280 Milligan *et al.* (1989) and the finding was later supported by a study suggesting differential
281 regulation of $G\alpha_{i2}$ and $G\alpha_{i3}$ in rat myometrium during gestation (Tanfin *et al.*, 1991). In
282 human myometrium, the levels of G protein subunits $G\alpha_{i1}$, $G\alpha_{i3}$, $G\alpha_q$ and $G\alpha_{11}$ have been
283 shown to remain constant in pregnant and non-pregnant women, while levels of $G\alpha_{i2}$ decrease
284 during pregnancy. The simultaneous, substantial increase in myometrial G_s suggested that the
285 balance between $G\alpha_{i2}$ and G_s might be essential in regulating relaxation of the uterus during
286 pregnancy (Europe-Finner *et al.*, 1993). Besides this, G_i family proteins have been suggested
287 to be functionally linked to α_2 adrenergic signalling in human myometrium during pregnancy
288 (Breuiller *et al.*, 1990). Later studies in the rat have confirmed the involvement of $G\alpha_{i2}$ and
289 $G\alpha_{i3}$ in α_2/β_2 adrenergic signalling in the maintenance of uterus relaxation during rat
290 pregnancy (Mhaouty *et al.*, 1995).

291

292 Unlike the thoroughly studied myometrium, the presence and role of $G\alpha_{i2}$ in other regions of
293 the reproductive tract has remained largely obscure. Although the presence of G_i family
294 proteins have been described in human endometrium during artificial cycles of hormone
295 replacement therapy, those studies rely solely on data from immunoblotting, using an
296 antibody unable to discriminate between the closely related $G\alpha_{i1}$ and $G\alpha_{i2}$ (Bernardini *et al.*,
297 1995, 1999). Therefore, prior to our study, cyclical changes in $G\alpha_{i2}$ expression have not been
298 reported in humans. Quantitative PCR showed that $G\alpha_{i2}$ expression in human endometrium *in*
299 *vivo* significantly increased towards secretory phase of the menstrual cycle. This suggested
300 that sex hormones, like oestrogen or progesterone, might regulate the expression of this G_i
301 subunit in human endometrium. Furthermore, immunostaining clearly demonstrated the main
302 localization of $G\alpha_{i2}$ in endometrial glands and partially in endometrial stroma.

303

304 It is likely that $G\alpha_{i2}$ is hormonally regulated in the human endometrium. Earlier studies on rat
305 myometrium have shown that estradiol administration during rat pregnancy increases the
306 levels of both $G\alpha_{i2}$ and $G\alpha_{i2}$ mRNA, while progesterone has no effect on $G\alpha_{i2}$ expression.
307 Instead, progesterone was reported to cause a decrease in $G\alpha_q$ subunit expression (Cohen-
308 Tannoudji *et al.*, 1995). Other studies in pregnant rat myometrium have suggested a
309 regulatory role for progesterone in control of β_2 receptors (Maltier *et al.*, 1989) and G_s
310 proteins (Elwardy-Merezak *et al.*, 1994), as well as in upregulation β_2 receptor expression
311 (Vivat *et al.*, 1992). Apart from the studies by Bernardini *et al.* (1995; 1999) the potential role
312 for sex hormones in regulation of G proteins in the human has remained largely unexplored.

313

314 In the present study, we have reported for the first time the localization of $G\alpha_{i2}$ in fallopian
315 tube epithelial cilia. In fallopian tubes, ciliary beat is essential for gamete transport in
316 association with the tubal secretory flow and muscle contractility. Furthermore, fallopian

317 tubes have been proposed to act as sperm reservoirs, where the ciliated epithelial cells interact
318 with sperm (Baillie *et al.*, 1997; Pacey *et al.*, 1995a; Pacey *et al.*, 1995b; Reeve *et al.*, 2003).
319 Fallopian tube epithelial cells have also been demonstrated to preserve the viability of sperm
320 (Kervancioglu *et al.*, 1994; Kervancioglu *et al.*, 2000; Murray and Smith, 1997). Given the
321 fact that $G\alpha_{i2}$ is specifically localized in rat tissue motile cilia with a characteristic 9+2
322 ultrastructure, namely in rat oviductal, tracheal and brain ependymal cilia (Shinohara *et al.*,
323 1998), it seems evident that this G_i subunit might have a cilia-specific physiological role.
324 Apart from proteomic analysis providing evidence of $G\alpha_{i2}$ as a resident axonemal protein of
325 the human bronchial cilia (Ostrowski *et al.*, 2002), there are no reports describing $G\alpha_{i2}$ in any
326 other human ciliated tissue. In addition to positive immunostaining of fallopian tube cilia, we
327 reported here positive immunostaining surrounding the nuclei. This presumably represents
328 pre-stage $G\alpha_{i2}$ which is still in synthesis, or alternatively, $G\alpha_{i2}$ which is ready for transport
329 into cilia by intraflagellar transport mechanisms. This intracellular machinery is vital for
330 assembly and maintenance of the cilia, as it transports essential particles, such as proteins
331 synthesised in the cytoplasm of cell, into the cilia, and returns the turnover products to the
332 cytoplasm of cell (Rosenbaum and Witman, 2002).

333

334 Studies with $G\alpha_{i2}$ -knockout mice have established a crucial regulatory role for the $G\alpha_{i2}$
335 subunit in immunological processes (Dalwadi *et al.*, 2003; Fan *et al.*, 2005; Han *et al.*, 2005;
336 Jiang *et al.*, 1997; Rudolph *et al.*, 1995; Rudolph *et al.*, 1995; Zhang *et al.*, 2005). $G\alpha_{i2}$ has
337 been revealed to control regulation of T-cell proliferation (Zhang *et al.*, 2005) and B cell
338 development (Dalwadi *et al.*, 2003). Furthermore, $G\alpha_{i2}$ has been suggested to mediate
339 chemokine signalling (Han *et al.*, 2005). However, reports of $G\alpha_{i2}$ -knockout studies have not
340 provided any information on potential involvement of this G_i subunit in modulation of mice
341 fertility. Interestingly, a recent study on $G\alpha_{i2}$ -knockout mice showed $G\alpha_{i2}$ to differentially

342 regulate inflammatory mediator production in response to microbial stimuli and proposed a
343 TLR-signalling regulating, anti-inflammatory role for $G\alpha_{i2}$ by an yet unknown mechanism
344 (Fan *et al.*, 2005). Regarding the potential link between TLR-signalling and $G\alpha_{i2}$ in female
345 reproductive tract, it is noteworthy that our previous studies showing the localization pattern
346 of several TLRs (Fazeli *et al.*, 2005) showed a similar pattern of localisation compared to that
347 we now report for $G\alpha_{i2}$. Future studies should be directed towards understanding whether
348 $G\alpha_{i2}$ might share signalling pathways with TLRs, and potentially have a TLR-signalling
349 regulating role in human reproductive tract.

350

351 In conclusion, our studies reveal the presence of $G\alpha_{i2}$ in human endometrium and fallopian
352 tube epithelium, especially the cilia of fallopian tube epithelial cells. To the best of our
353 knowledge, this is the first report of the localization of $G\alpha_{i2}$ in ciliated reproductive tissue in
354 the human. We also report here, for the first time, the alterations in $G\alpha_{i2}$ expression during
355 human menstrual cycle. Our data implies this G_i family subunit might be under hormonal
356 regulation in the female reproductive tract *in vivo*. Further studies are required to clarify the
357 physiological role of $G\alpha_{i2}$ in the female reproductive tract.

358

359

360

361 **Acknowledgments**

362 We wish to thank Chris Bruce and Sarah Elliott for skilful technical assistance. This study
363 was financially supported by The Association of Finnish Pharmacies, Finnish Pharmaceutical
364 Society, Alfred Kordelin Foundation, Farnos Science and Research Foundation, Helsingin
365 Sanomat Centennial Foundation and The Saastamoinen Foundation.

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386 **References**

387

- 388 Arey, B. J., Stevis, P. E., Deecher, D. C., Shen, E. S., Frail, D. E., Negro-Vilar, A. and Lopez,
389 F. J. (1997) Induction of promiscuous G protein coupling of the follicle-stimulating
390 hormone (FSH) receptor: a novel mechanism for transducing pleiotropic actions of
391 FSH isoforms. *Mol Endocrinol* **11**, 517-26.
- 392 Baillie, H. S., Pacey, A. A., Warren, M. A., Scudamore, I. W. and Barratt, C. L. (1997)
393 Greater numbers of human spermatozoa associate with endosalpingeal cells derived
394 from the isthmus compared with those from the ampulla. *Hum Reprod* **12**, 1985-92.
- 395 Benten, W. P., Stephan, C., Lieberherr, M. and Wunderlich, F. (2001) Estradiol signaling via
396 sequestrable surface receptors. *Endocrinology* **142**, 1669-77.
- 397 Bernardini, L., Moretti-Rojas, I., Brush, M., Rojas, F. J. and Balmaceda, J. P. (1995) Status of
398 hCG/LH receptor and G proteins in human endometrium during artificial cycles of
399 hormone replacement therapy. *J Soc Gynecol Investig* **2**, 630-5.
- 400 Bernardini, L., Moretti-Rojas, I., Brush, M., Rojas, F. J. and Balmaceda, J. P. (1999) Changes
401 in expression of adenylyl cyclase activity in human endometrium during hormone
402 replacement therapy and ovarian stimulation. *Mol Hum Reprod* **5**, 955-60.
- 403 Bokoch, G. M., Katada, T., Northup, J. K., Ui, M. and Gilman, A. G. (1984) Purification and
404 properties of the inhibitory guanine nucleotide-binding regulatory component of
405 adenylyl cyclase. *J Biol Chem* **259**, 3560-7.
- 406 Breuiller, M., Rouot, B., Litime, M. H., Leroy, M. J. and Ferre, F. (1990) Functional coupling
407 of the alpha 2-adrenergic receptor-adenylyl cyclase complex in the pregnant human
408 myometrium. *J Clin Endocrinol Metab* **70**, 1299-304.
- 409 Chi, L., Zhou, W., Prikhozhan, A., Flanagan, C., Davidson, J. S., Golembo, M., Illing, N.,
410 Millar, R. P. and Sealfon, S. C. (1993) Cloning and characterization of the human
411 GnRH receptor. *Mol Cell Endocrinol* **91**, R1-6.
- 412 Cohen-Tannoudji, J., Mhaouty, S., Elwardy-Merezak, J., Lecrivain, J. L., Robin, M. T.,
413 Legrand, C. and Maltier, J. P. (1995) Regulation of myometrial Gi2, Gi3, and Gq
414 expression during pregnancy. Effects of progesterone and estradiol. *Biol Reprod* **53**,
415 55-64.
- 416 Dalwadi, H., Wei, B., Schrage, M., Spicher, K., Su, T. T., Birnbaumer, L., Rawlings, D. J.
417 and Braun, J. (2003) B cell developmental requirement for the G alpha i2 gene. *J*
418 *Immunol* **170**, 1707-15.
- 419 Elwardy-Merezak, J., Maltier, J. P., Cohen-Tannoudji, J., Lecrivain, J. L., Vivat, V. and
420 Legrand, C. (1994) Pregnancy-related modifications of rat myometrial Gs proteins:
421 ADP ribosylation, immunoreactivity and gene expression studies. *J Mol Endocrinol*
422 **13**, 23-37.
- 423 Europe-Finner, G. N., Phaneuf, S., Watson, S. P. and Lopez Bernal, A. (1993) Identification
424 and expression of G-proteins in human myometrium: up-regulation of G alpha s in
425 pregnancy. *Endocrinology* **132**, 2484-90.
- 426 Fan, H., Zingarelli, B., Peck, O. M., Teti, G., Tempel, G. E., Halushka, P. V., Spicher, K.,
427 Boulay, G., Birnbaumer, L. and Cook, J. A. (2005) Lipopolysaccharide- and gram-
428 positive bacteria-induced cellular inflammatory responses: role of heterotrimeric
429 Galpha(i) proteins. *Am J Physiol Cell Physiol* **289**, C293-301.
- 430 Fazeli, A., Bruce, C. and Anumba, D. O. (2005) Characterization of Toll-like receptors in the
431 female reproductive tract in humans. *Hum Reprod* **20**, 1372-8.

- 432 Han, S. B., Moratz, C., Huang, N. N., Kelsall, B., Cho, H., Shi, C. S., Schwartz, O. and Kehrl,
433 J. H. (2005) Rgs1 and Gnai2 regulate the entrance of B lymphocytes into lymph nodes
434 and B cell motility within lymph node follicles. *Immunity* **22**, 343-54.
- 435 Hawes, B. E., Barnes, S. and Conn, P. M. (1993) Cholera toxin and pertussis toxin provoke
436 differential effects on luteinizing hormone release, inositol phosphate production, and
437 gonadotropin-releasing hormone (GnRH) receptor binding in the gonadotrope:
438 evidence for multiple guanyl nucleotide binding proteins in GnRH action.
439 *Endocrinology* **132**, 2124-30.
- 440 Hepler, J. R. and Gilman, A. G. (1992) G proteins. *Trends Biochem Sci* **17**, 383-7.
- 441 Herrlich, A., Kuhn, B., Grosse, R., Schmid, A., Schultz, G. and Gudermann, T. (1996)
442 Involvement of Gs and Gi proteins in dual coupling of the luteinizing hormone
443 receptor to adenylyl cyclase and phospholipase C. *J Biol Chem* **271**, 16764-72.
- 444 Jiang, M., Boulay, G., Spicher, K., Peyton, M. J., Brabet, P., Birnbaumer, L. and Rudolph, U.
445 (1997) Inactivation of the G alpha i2 and G alpha o genes by homologous
446 recombination. *Receptors Channels* **5**, 187-92.
- 447 Karteris, E., Zervou, S., Pang, Y., Dong, J., Hillhouse, E. W., Randevara, H. S. and Thomas, P.
448 (2006) Progesterone signaling in human myometrium through two novel membrane G
449 protein-coupled receptors: potential role in functional progesterone withdrawal at term.
450 *Mol Endocrinol* **20**, 1519-34.
- 451 Katada, T., Northup, J. K., Bokoch, G. M., Ui, M. and Gilman, A. G. (1984) The inhibitory
452 guanine nucleotide-binding regulatory component of adenylate cyclase. Subunit
453 dissociation and guanine nucleotide-dependent hormonal inhibition. *J Biol Chem* **259**,
454 3578-85.
- 455 Kervancioglu, M. E., Djahanbakhch, O. and Aitken, R. J. (1994) Epithelial cell coculture and
456 the induction of sperm capacitation. *Fertil Steril* **61**, 1103-8.
- 457 Kervancioglu, M. E., Saridogan, E., Aitken, R. J. and Djahanbakhch, O. (2000) Importance of
458 sperm-to-epithelial cell contact for the capacitation of human spermatozoa in fallopian
459 tube epithelial cell cocultures. *Fertil Steril* **74**, 780-4.
- 460 Krsmanovic, L. Z., Mores, N., Navarro, C. E., Arora, K. K. and Catt, K. J. (2003) An agonist-
461 induced switch in G protein coupling of the gonadotropin-releasing hormone receptor
462 regulates pulsatile neuropeptide secretion. *Proc Natl Acad Sci U S A* **100**, 2969-74.
- 463 Krsmanovic, L. Z., Mores, N., Navarro, C. E., Tomic, M. and Catt, K. J. (2001) Regulation of
464 Ca²⁺-sensitive adenylyl cyclase in gonadotropin-releasing hormone neurons. *Mol*
465 *Endocrinol* **15**, 429-40.
- 466 Loosfelt, H., Misrahi, M., Atger, M., Salesse, R., Vu Hai-Luu Thi, M. T., Jolivet, A.,
467 Guiochon-Mantel, A., Sar, S., Jallal, B., Garnier, J. *et al.* (1989) Cloning and
468 sequencing of porcine LH-hCG receptor cDNA: variants lacking transmembrane
469 domain. *Science* **245**, 525-8.
- 470 Maltier, J. P., Benghan-Eyene, Y. and Legrand, C. (1989) Regulation of myometrial beta 2-
471 adrenergic receptors by progesterone and estradiol-17 beta in late pregnant rats. *Biol*
472 *Reprod* **40**, 531-40.
- 473 McFarland, K. C., Sprengel, R., Phillips, H. S., Kohler, M., Rosemblyt, N., Nikolics, K.,
474 Segaloff, D. L. and Seeburg, P. H. (1989) Lutropin-choriogonadotropin receptor: an
475 unusual member of the G protein-coupled receptor family. *Science* **245**, 494-9.
- 476 Mhaouty, S., Cohen-Tannoudji, J., Bouet-Alard, R., Limon-Boulez, I., Maltier, J. P. and
477 Legrand, C. (1995) Characteristics of the alpha 2/beta 2-adrenergic receptor-coupled
478 adenylyl cyclase system in rat myometrium during pregnancy. *J Biol Chem* **270**,
479 11012-6.

- 480 Milligan, G., Tanfin, Z., Goureau, O., Unson, C. and Harbon, S. (1989) Identification of both
481 Gi2 and a novel, immunologically distinct, form of Go in rat myometrial membranes.
482 *FEBS Lett* **244**, 411-6.
- 483 Minegishi, T., Nakamura, K., Takakura, Y., Ibuki, Y., Igarashi, M. and Minegishi, T. (1991)
484 Cloning and sequencing of human FSH receptor cDNA. *Biochem Biophys Res*
485 *Commun* **175**, 1125-30.
- 486 Minegishi, T., Nakamura, K., Takakura, Y., Miyamoto, K., Hasegawa, Y., Ibuki, Y., Igarashi,
487 M. and Minegishi, T. (1990) Cloning and sequencing of human LH/hCG receptor
488 cDNA. *Biochem Biophys Res Commun* **172**, 1049-54.
- 489 Murray, S. C. and Smith, T. T. (1997) Sperm interaction with fallopian tube apical membrane
490 enhances sperm motility and delays capacitation. *Fertil Steril* **68**, 351-7.
- 491 Ostrowski, L. E., Blackburn, K., Radde, K. M., Moyer, M. B., Schlutzer, D. M., Moseley, A.
492 and Boucher, R. C. (2002) A proteomic analysis of human cilia: identification of novel
493 components. *Mol Cell Proteomics* **1**, 451-65.
- 494 Pacey, A. A., Davies, N., Warren, M. A., Barratt, C. L. and Cooke, I. D. (1995a)
495 Hyperactivation may assist human spermatozoa to detach from intimate association
496 with the endosalpinx. *Hum Reprod* **10**, 2603-9.
- 497 Pacey, A. A., Hill, C. J., Scudamore, I. W., Warren, M. A., Barratt, C. L. and Cooke, I. D.
498 (1995b) The interaction in vitro of human spermatozoa with epithelial cells from the
499 human uterine (fallopian) tube. *Hum Reprod* **10**, 360-6.
- 500 Reeve, L., Ledger, W. L. and Pacey, A. A. (2003) Does the Arg-Gly-Asp (RGD) adhesion
501 sequence play a role in mediating sperm interaction with the human endosalpinx?
502 *Hum Reprod* **18**, 1461-8.
- 503 Rosenbaum, J. L. and Witman, G. B. (2002) Intraflagellar transport. *Nat Rev Mol Cell Biol* **3**,
504 813-25.
- 505 Rudolph, U., Finegold, M. J., Rich, S. S., Harriman, G. R., Srinivasan, Y., Brabet, P., Boulay,
506 G., Bradley, A. and Birnbaumer, L. (1995) Ulcerative colitis and adenocarcinoma of
507 the colon in G alpha i2-deficient mice. *Nat Genet* **10**, 143-50.
- 508 Rudolph, U., Finegold, M. J., Rich, S. S., Harriman, G. R., Srinivasan, Y., Brabet, P., Bradley,
509 A. and Birnbaumer, L. (1995) Gi2 alpha protein deficiency: a model of inflammatory
510 bowel disease. *J Clin Immunol* **15**, 101S-105S.
- 511 Shinohara, H., Asano, T., Kato, K., Kameshima, T. and Semba, R. (1998) Localization of a G
512 protein Gi2 in the cilia of rat ependyma, oviduct and trachea. *Eur J Neurosci* **10**, 699-
513 707.
- 514 Sprengel, R., Braun, T., Nikolics, K., Segaloff, D. L. and Seeburg, P. H. (1990) The testicular
515 receptor for follicle stimulating hormone: structure and functional expression of
516 cloned cDNA. *Mol Endocrinol* **4**, 525-30.
- 517 Stanislaus, D., Ponder, S., Ji, T. H. and Conn, P. M. (1998) Gonadotropin-releasing hormone
518 receptor couples to multiple G proteins in rat gonadotrophs and in GGH3 cells:
519 evidence from palmitoylation and overexpression of G proteins. *Biol Reprod* **59**, 579-
520 86.
- 521 Tanfin, Z., Goureau, O., Milligan, G. and Harbon, S. (1991) Characterization of G proteins in
522 rat myometrium. A differential modulation of Gi2 alpha and Gi3 alpha during
523 gestation. *FEBS Lett* **278**, 4-8.
- 524 Tsutsumi, M., Zhou, W., Millar, R. P., Mellon, P. L., Roberts, J. L., Flanagan, C. A., Dong,
525 K., Gillo, B. and Sealfon, S. C. (1992) Cloning and functional expression of a mouse
526 gonadotropin-releasing hormone receptor. *Mol Endocrinol* **6**, 1163-9.
- 527 Vivat, V., Cohen-Tannoudji, J., Revelli, J. P., Muzzin, P., Giacobino, J. P., Maltier, J. P. and
528 Legrand, C. (1992) Progesterone transcriptionally regulates the beta 2-adrenergic
529 receptor gene in pregnant rat myometrium. *J Biol Chem* **267**, 7975-8.

530 Zhang, Y., Finegold, M. J., Jin, Y. and Wu, M. X. (2005) Accelerated transition from the
531 double-positive to single-positive thymocytes in G alpha i2-deficient mice. *Int*
532 *Immunol* **17**, 233-43.

533 Zhu, Y., Rice, C. D., Pang, Y., Pace, M. and Thomas, P. (2003) Cloning, expression, and
534 characterization of a membrane progestin receptor and evidence it is an intermediary
535 in meiotic maturation of fish oocytes. *Proc Natl Acad Sci U S A* **100**, 2231-6.

536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573

574 **Figure legends**

575

576 **Figure 1.** PCR showed $G\alpha_{i2}$ expression in fallopian tube tissue (A), human endometrium
577 tissue (B) and fallopian tube primary epithelial cells (C). PCR products were separated on 1.2
578 % agarose gel. 1: β -actin (643 base pairs), 2: β actin RT control, 3: $G\alpha_{i2}$ (212 base pairs), 4:
579 $G\alpha_{i2}$ RT control, M_W : molecular weight (base pairs).

580

581 **Figure 2.** Immunostaining showing localization of $G\alpha_{i2}$ in human endometrial cryosections
582 and fallopian tube paraffin embedded sections. $G\alpha_{i2}$ is enriched in endometrial glands,
583 proliferative phase (A), secretory phase (B). Immunostaining of human fallopian tube paraffin
584 embedded sections (C) indicated specific localization of $G\alpha_{i2}$ in fallopian tube epithelial cells
585 and the cilia. $G\alpha_{i2}$ (brown). Negative control slides were incubated with diluent only. All the
586 slides were counterstained with haematoxylin (blue). Scale bar: 100 μ m (A, B), 40 μ m (C).

587

588 **Figure 3.** Quantitative real time PCR uncovered variable expression of $G\alpha_{i2}$ gene in
589 endometrium during the menstrual cycle. Endometrial biopsies were designated in three
590 groups according to menstrual history of the patient (menstrual n=3, proliferative and
591 secretory n=9). The figure illustrates mean \pm SEM of normalised $G\alpha_{i2}$ gene expression. *
592 Secretory phase was significantly different from the other phases, $p < 0.05$; One-way
593 ANOVA with Tukey's multiple comparison test.

594

595

596

597

598 **Figure legends**

599

600 **Figure 1.** PCR showed $G\alpha_{i2}$ expression in fallopian tube tissue (A), human endometrium
601 tissue (B), immortalized fallopian tube epithelial cell line (OE-E6/E7) (C) and fallopian tube
602 primary epithelial cells (D). PCR products were separated on 1.2 % agarose gel. 1: β -actin
603 (643 base pairs), 2: β actin RT control, 3: $G\alpha_{i2}$ (212 base pairs), 4: $G\alpha_{i2}$ RT control, M_W :
604 molecular weight (base pairs).

605

606 **Figure 2.** Immunostaining shows localization of $G\alpha_{i2}$ in human endometrial cryosections and
607 fallopian tube paraffin embedded sections. $G\alpha_{i2}$ is enriched in endometrial glands,
608 proliferative phase (A), secretory phase (B). Immunostaining of human fallopian tube paraffin
609 embedded sections (C) indicated specific localization of $G\alpha_{i2}$ in epithelial cells and the cilia.
610 $G\alpha_{i2}$ (brown): Chemicon MAB3077 primary antibody was used with dilutions of 1:1000 for
611 endometrial cryosections and 1:500 for paraffin embedded fallopian tube sections. DAB or
612 DAB-Ni was used as a chromogen (endometrial cryosections and paraffin embedded fallopian
613 tube sections, respectively). Negative control slides were incubated with diluent only. All the
614 slides were counterstained with haematoxylin (blue). Scale bar: 100 μ m.

615

616 **Figure 3.** Western blot analysis confirmed the presence of $G\alpha_{i2}$ in immortalized fallopian
617 tube epithelial cell line (OE-E6/E7). A: G protein standard, (2 μ l / lane) Bovine brain
618 immunoblot standard, Calbiochem. B: Homogenate of fallopian tube epithelial cells, (60 μ g /
619 lane).


620

621 **Figure 4.** Quantitative real time PCR uncovered variable expression of $G\alpha_{i2}$ in endometrium
622 during the menstrual cycle. Endometrial biopsies were designated in three groups according
623 to menstrual history of the patient (menstrual n=3, proliferative and secretory n=9). The figure
624 illustrates mean \pm SEM of normalised $G\alpha_{i2}$ gene expression. * Secretory phase was

625 significantly different from the other phases, $p < 0.05$; One-way ANOVA with Tukey's
626 multiple comparison test.

627

628 **Figures**



643
644
645
646

Figure 1. KS Mönkkönen et al.



647

648



G α_{i2}

Negative control



A



B



C



691
692
693
694
695
696

Figure 2. KS Mönkkönen et al.

697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727




Figure 3. KS Mönkkönen et al.




Figure 4. KS Mönkkönen et al.