

COMPUTER SCIENCE PUBLICATION

A LOWER BOUND FOR INTERVAL ROUTING IN
GENERAL NETWORKS

Savio S.H. Tse and Francis C.M. Lau


Technical Report TR-94-09

July 1994



DEPARTMENT OF COMPUTER SCIENCE
FACULTY OF ENGINEERING
UNIVERSITY OF HONG KONG
POKFULAM ROAD
HONG KONG

UNIVERSITY OF HONG KONG
LIBRARY



*This book was a gift
from*

Dept. of Computer Science
The University of Hong Kong

A Lower Bound for Interval Routing in General Networks

Savio S.H. Tse and Francis C.M. Lau*

Department of Computer Science
The University of Hong Kong

July 1994

Abstract

Interval routing is a space-efficient routing method for point-to-point communication networks. The method has drawn considerable attention in recent years because of its being incorporated into the design of a commercially available routing chip. The method is based on proper labeling of edges of the graph with intervals. An optimal labeling would result in routing of messages through the shortest paths. Optimal labelings have existed for regular as well as some of the common topologies, but not for arbitrary graphs. In fact, it has already been shown that it is impossible to find optimal labelings for arbitrary graphs. In this paper, we prove a $7D/4 - 1$ lower bound for interval routing in arbitrary graphs, where D is the diameter—*i.e.*, the best any interval labeling scheme could do is to produce a longest path having a length of at least $7D/4 - 1$.

Keywords: communication networks, interval routing, lower bounds, routing algorithms.

*Correspondence: Dr F.C.M. Lau, Department of Computer Science, The University of Hong Kong, Hong Kong / Email: fcmlau@csd.hku.hk / Fax: (+852) 559 8447

1 Introduction

Routing is an important operation in communication networks. Since it is frequently invoked, it is worth the while to try to optimize the number of steps taken to route a message from one node to another. Obviously, we can achieve optimal routing by keeping a table of size $O(n)$ in each node, n being the number of nodes in the network. For large networks, this may not be practical. Various methods that use much less space have been proposed, including *interval routing* which has been adopted in the design of a commercially available routing chip. The idea of interval routing is to label every node with a number from a linearly and cyclicly ordered set, for example $\{0, \dots, n-1\}$, and every (directed) edge with an interval (of range of node numbers). To understand how the method works, refer to Figure 1 which shows an example of a simple network, complete with node numbers and interval labels, and a path traversed by a message (from Node 2 to Node 0) by following the interval labels. In the figure, an interval label of the form $\langle i, j \rangle$ corresponds to the range of node numbers from i to j ; intervals of the form $\langle k \rangle$ contain the single node number k . The message, being destined for Node 0, first takes the edge to Node 3 because 0 is contained in the interval $\langle 3, 0 \rangle$, and then it takes the edge to Node 4 because 0 is contained in $\langle 4, 0 \rangle$, and so on. It can be seen that at most $O(d)$ space is needed at a node, where d is the node's degree. The idea of interval




Figure 1: Example of interval routing

routing was first proposed by Santoro and Khatib [6] who used a spanning tree at every

node to carry out the assignment of interval labels. As a result, not all the edges are used for routing in their scheme. Later on, van Leeuwen and Tan extended the method to make use of all the edges [8]. Their labeling scheme can produce optimal labelings for common topologies such as trees, rings, complete graphs, and some grids. An optimal labeling is such that the routing of a message from any node to any other node would take the shortest path in the graph. Their labeling scheme as well as Santoro and Khatib's, however, are not able to generate optimal labelings for arbitrary graphs. Ružička proved that it is impossible to find optimal interval labelings for arbitrary graphs. Specifically, Ružička found a graph for which he proved that no interval labeling can result in a longest path which is shorter than $3D/2 + 1/2$ (or $1.5D + 1/2$). In this paper, we give an improved lower bound of $7D/4 - 1$ (or $1.75D - 1$). We use a graph which bears certain resemblance to Ružička's graph, but is slightly more complicated.

Our lower bound result suggests that Santoro and Khatib's labeling algorithm [6], which produces paths that are no longer than $2D$ for arbitrary networks, is very close to the best possible. Their labeling algorithm, however, might suffer from bottleneck problems due to the use of a spanning tree for the routes. Interval labeling has been incorporated into the latest routing chip, the C104, by Inmos [3, 4], which undoubtedly would add to the need of finding even better interval labeling algorithms. The graph as presented in this paper can be used as a test case for measuring the goodness of such algorithms.

In the following, we assume that interval labels are cyclic.¹ In addition to interval labels, there could also be *null labels* and *complement labels* [5]. An edge labeled with a null label is never taken in routing messages. An edge with a complement label is taken when the interval label of all other edges fail to contain the destination node number. It can be easily seen that if null labels are allowed in our graph which will be presented in Section 3, a lower bound of $2D - 1$ would result. Therefore, in the following, we consider only interval and complement labels. Obviously, a node can have at most one complement label. We first present a lower bound for the case of using only interval labels (Section 3). Then in Section 4 we modify the graph to allow for the use of complement labels, and the modification is such that the complement labels, no matter

¹The scheme is called *linear interval routing* when non-cyclic labels are used [1, 2].

where they are placed, cannot help to shorten the longest path; as a result, we have the same lower bound for the latter case based on the modified graph.

2 Definitions, Notations, and Properties

The network in question is an undirected graph, $G = (E, V)$, where E is the set of edges, and V the set the nodes. Every edge in E is actually made up of two directed edges, one for each direction (as in Figure 1). There are n nodes in V . To implement interval routing, each node is labeled with a unique integer, called a *node number*, from the set $L = \{0, \dots, n - 1\}$. For simplicity, we assume a node's number is the same as the node's name.

Every edge in each direction is labeled with an *interval label* (or *interval*) which is of the form $\langle p, q \rangle$, where $p, q \in L$. For $u, v \in V$ that are directly connected, $L(u, v)$ denotes the interval label for the edge that goes from u to v . A node m (or its number) is said to be contained in $\langle p, q \rangle$ if (1) $p \leq m \leq q$ for $p \leq q$, or (2) $p \leq m \leq n - 1$ or $0 \leq m \leq q$ otherwise. We use the notation $u \prec v \prec w$, $u, v, w \in L$, to denote the cyclic ordering of node numbers. Naturally, $0 \prec 1 \prec \dots \prec n - 1 \prec 0$.

In the following, subsets of node numbers that are contained in some interval often occur inside expressions—we use the set notation to denote them. For example, $\{u, v, w\}$ refers to three node numbers, u, v, w , that are contained in some interval and whose order is not specified. The expression $u \prec \{v, w\} \prec x \dots$ means that v and w are contained in some interval and that they are ordered after u and before x , but the order of v and w is not known.

Property 2.1 (*Completeness*) *The set of interval labels for edges directed from a node u is complete. That is, every node in $V \neq u$ must be contained in some u 's interval.*

Property 2.2 (*No ambiguity*) *The interval labels for edges directed from a node u are disjoint. That is, for node $v \neq u$, v is contained in exactly one of these intervals.*

Property 2.3 (*No bouncing*) *For any edge $(u, v) \in E$, there exists no node $w \neq u, v$ such that w is contained in both $L(u, v)$ and $L(v, u)$.*

Given node u , Property 2.2 implies that $L(u, v) \cap L(u, w) = \emptyset$, where (u, v) and (u, w) are any two edges directed from u . And Property 2.3 implies that $L(u, v) \cap L(v, u) = \emptyset$.

3 Lower Bound

We are going to be more specific about the graph G based on which we will derive our lower bound. Figure 2 shows the details of G which consists of three identical “flaps”, each of length $2k$ (edges), $k \geq 3$, extending from a middle axis. The set of nodes V is




Figure 2: The graph G

made up of $\{u_{i,j}, v_{i,j}, w_{i,j} | 1 \leq i \leq 8, 1 \leq j \leq 2k - 1\} \cup \{m_i | 1 \leq i \leq 8\} \cup \{u_0, v_0, w_0\}$. The number of nodes, n , is therefore equal to $3 \times 8 \times (2k - 1) + 8 + 3 = 48k - 13$. The diameter of G , D , is $4k$. The lower bound we are going to prove is $7D/4 - 1$ —i.e., there exists no labeling scheme such that the longest path in G is shorter than $7D/4 - 1$. In Figure 2, we included an example of a path which is of length $7D/4 - 1$ to give an idea of the magnitude of this value. We will prove the bound by contradiction.

If there is a labeling scheme such that the longest path is shorter than $7D/4 - 1$, then

the following three lemmas hold.

Lemma 3.1 *For every $i \in \{1, \dots, 8\}$, there exists an interval label that contains*

$$\{u_{i,k+1}, u_{i,2k-1}, v_{i,2k-1}, w_{i,2k-1}\}$$

but does not contain $\{u_{1,1}, u_{2,1}, \dots, u_{8,1}\}$.

Proof: Consider $u_{i,k}$. $L(u_{i,k}, u_{i,k+1})$ must contain $\{u_{i,k+1}, u_{i,2k-1}, v_{i,2k-1}, w_{i,2k-1}\}$ and $L(u_{i,k}, u_{i,k-1})$ must contain $\{u_{1,1}, u_{2,1}, \dots, u_{8,1}\}$, but by Property 2.2, these two interval labels are disjoint. \square

Lemma 3.2 *For every $i \in \{1, \dots, 8\}$, there exist three disjoint intervals containing $\{u_{i,2k-1}, u_{i,k-1}\}$, $\{v_{i,2k-1}, v_{i,k-1}\}$, and $\{w_{i,2k-1}, w_{i,k-1}\}$, respectively.*

Proof: By considering the three edges directed from $m_i, i = 1, \dots, 8$. \square

Lemma 3.3 *There exist four or more disjoint intervals each of which contains*

$$\{u_{i,2k-1}, v_{i,2k-1}, w_{i,2k-1}\},$$

where $i \in \{1, \dots, 8\}$.

Proof: Without loss of generality, suppose $u_{1,1} \prec u_{2,1} \prec \dots \prec u_{8,1} \prec u_{1,1}$. Consider u_0 . If there is a labeling scheme such that the longest path is shorter than $7D/4 - 1$, then $L(u_0, u_{i,1})$ contains $\{u_{i,1}, u_{i,k+1}\}$, for $i = 1, \dots, 8$. Since all intervals of the same node are disjoint (Property 2.2), we have

$$\{u_{1,1}, u_{1,k+1}\} \prec \{u_{2,1}, u_{2,k+1}\} \prec \dots \prec \{u_{8,1}, u_{8,k+1}\} \prec \{u_{1,1}, u_{1,k+1}\}.$$

By ignoring some of the node numbers, we have

$$u_{1,1} \prec u_{2,k+1} \prec u_{3,1} \prec u_{4,k+1} \prec u_{5,1} \prec u_{6,k+1} \prec u_{7,1} \prec u_{8,k+1} \prec u_{1,1}.$$

And then by Lemma 3.1, we have

$$\begin{aligned} u_{1,1} \prec \{u_{2,k+1}, u_{2,2k-1}, v_{2,2k-1}, w_{2,2k-1}\} \prec u_{3,1} \prec \{u_{4,k+1}, u_{4,2k-1}, v_{4,2k-1}, w_{4,2k-1}\} \prec u_{5,1} \\ \prec \{u_{6,k+1}, u_{6,2k-1}, v_{6,2k-1}, w_{6,2k-1}\} \prec u_{7,1} \prec \{u_{8,k+1}, u_{8,2k-1}, v_{8,2k-1}, w_{8,2k-1}\} \prec u_{1,1} \end{aligned}$$

or

$$\begin{aligned} & \{u_{2,2k-1}, v_{2,2k-1}, w_{2,2k-1}\} \prec \{u_{4,2k-1}, v_{4,2k-1}, w_{4,2k-1}\} \\ & \prec \{u_{6,2k-1}, v_{6,2k-1}, w_{6,2k-1}\} \prec \{u_{8,2k-1}, v_{8,2k-1}, w_{8,2k-1}\}. \end{aligned}$$

□

We denote these four subsets of intervals by C_2, C_4, C_6, C_8 , respectively. Figure 3 shows the axis portion of G and the locations of these four subsets.




Figure 3: Four interval subsets at the centre

Theorem 3.1 *There exists no labeling scheme such that the longest path is shorter than $7D/4 - 1$.*

Proof: Assume that a longest path of length shorter than $7D/4 - 1$ exists. Consider C_4 in Lemma 3.3 and all the possible orderings of the three node numbers that were shown.

$$\begin{aligned} & C_2 \prec u_{4,2k-1} \prec \underline{u_{4,2k-1}} \prec w_{4,2k-1} \prec C_6 \prec C_8 \\ & C_2 \prec u_{4,2k-1} \prec \underline{w_{4,2k-1}} \prec v_{4,2k-1} \prec C_6 \prec C_8 \\ & C_2 \prec v_{4,2k-1} \prec \underline{u_{4,2k-1}} \prec w_{4,2k-1} \prec C_6 \prec C_8 \\ & C_2 \prec v_{4,2k-1} \prec \underline{w_{4,2k-1}} \prec u_{4,2k-1} \prec C_6 \prec C_8 \\ & C_2 \prec w_{4,2k-1} \prec \underline{u_{4,2k-1}} \prec v_{4,2k-1} \prec C_6 \prec C_8 \\ & C_2 \prec w_{4,2k-1} \prec \underline{v_{4,2k-1}} \prec u_{4,2k-1} \prec C_6 \prec C_8 \end{aligned}$$

Note that there are three possible choices for the middle place (underlined above) among the three places of C_4 . All together there are four middle places for C_2, C_4, C_6, C_8 , respectively, which are to be occupied by four elements from the following three sets.

$$\begin{aligned} & \{u_{2,2k-1}, u_{4,2k-1}, u_{6,2k-1}, u_{8,2k-1}\} \\ & \{v_{2,2k-1}, v_{4,2k-1}, v_{6,2k-1}, v_{8,2k-1}\} \\ & \{w_{2,2k-1}, w_{4,2k-1}, w_{6,2k-1}, w_{8,2k-1}\} \end{aligned}$$

Hence one set will contribute at least two elements to the middle places. Without loss of generality, suppose that the first set above contributes two elements to the middle places of C_4 and C_6 ; that is

$$\underbrace{v_{4,2k-1} \prec \underline{u_{4,2k-1}} \prec w_{4,2k-1}}_{C_4} \prec \underbrace{v_{6,2k-1} \prec \underline{u_{6,2k-1}} \prec w_{6,2k-1}}_{C_6}.$$

By Lemma 3.2, we have

$$v_{4,2k-1} \prec \{u_{4,2k-1}, u_{4,k-1}\} \prec w_{4,2k-1} \prec v_{6,2k-1} \prec \{u_{6,2k-1}, u_{6,k-1}\} \prec w_{6,2k-1}$$

or

$$v_{4,2k-1} \prec u_{4,k-1} \prec w_{4,2k-1} \prec v_{6,2k-1} \prec u_{6,k-1} \prec w_{6,2k-1}.$$

Now consider $u_{4,k}$, $L(u_{4,k}, u_{4,k-1})$ must contain $\{u_{4,k-1}, u_{1,1}, u_{2,1}, \dots, u_{8,1}\}$, and $L(u_{4,k}, u_{4,k+1})$ must contain $\{v_{4,2k-1}, w_{4,2k-1}\}$. Therefore, we have

$$v_{4,2k-1} \prec \{u_{4,k-1}, u_{1,1}, u_{2,1}, \dots, u_{8,1}\} \prec w_{4,2k-1} \prec v_{6,2k-1} \prec u_{6,k-1} \prec w_{6,2k-1}.$$

Similarly, $L(u_{6,k}, u_{6,k-1})$ must contain $\{u_{6,k-1}, u_{1,1}, u_{2,1}, \dots, u_{8,1}\}$, and $L(u_{6,k}, u_{6,k+1})$ must contain $\{v_{6,2k-1}, w_{6,2k-1}\}$. But in order for $\{v_{6,2k-1}, w_{6,2k-1}\}$ to be in the same interval, the interval must include $\{u_{4,k-1}, u_{1,1}, u_{2,1}, \dots, u_{8,1}\}$ (in order to not include $u_{6,k-1}$) according to the above cyclic order. Hence, $L(u_{6,k}, u_{6,k+1}) \cap L(u_{6,k}, u_{6,k-1}) = \{u_{1,1}, u_{2,1}, \dots, u_{8,1}\} \neq \emptyset$, which violates Property 2.2. \square

4 Graphs with Complement Labels

In the above, we allowed only interval labels for the labeling scheme. Here, we extend the lower bound result to cover also labeling schemes that use complement labels in

addition to interval labels. For any node, one can assign at most one complement label to the node. We transform the graph, G , we used in the previous section to another graph, G' . Based on G' , we show that the lower bound for labeling schemes that can use complement labels is the same as before.

The new graph, G' , as shown in Figure 4, has 36 layers and four flaps. The newly




Figure 4: The graph G'

added parts are represented by dashed lines in the figure. The previous graph, G , is a proper subgraph of G' .

Theorem 4.1 *There exists no labeling scheme (complement labels allowed) such that the longest path is shorter than $7D/4 - 1$.*

Proof: The nodes of G' that can have a complement label are u_0, v_0, w_0, x_0 and $m_i, i = 1, \dots, 12$. All the other nodes are of degree two and therefore have either two interval labels or one interval label and a null label. Since each of u_0, v_0, w_0, x_0 can have at most one complement label, the distribution of the complement labels of u_0, v_0, w_0, x_0 covers at most four layers (see the example in the figure—the bottom four layers), leaving at

most 32 layers that are without complement labels at the tips of the four flaps. Then, consider the m_i 's: since each m_i can be assigned at most one complement label, and each m_i has four edges, we can find at least $32/4 = 8$ m_i 's for which three of their four flaps have no complement labels. Without loss of generality, suppose these three flaps are the ones tipped at u_0, v_0, w_0 , respectively. As a result, we are left with a subgraph which is free of complement labels; this subgraph is either G or one that properly contains G . Let's concentrate on G and ignore for the moment those nodes that are not in G . Since every node of subgraph G must be able to reach every other node of G , we have an interval routing problem for subgraph G which is exactly equal to the interval routing problem we had in the previous section. Therefore, by Theorem 3.1, there exists at least one path in G whose length is longer than or equal to $7D/4 - 1$; let this path be P . Now we put the nodes we just ignored back into the picture. It is then easy to see that P cannot be shortened by noting the following.

1. The added nodes would not affect the proof of Theorem 3.1 which is based on subsets of intervals relevant to the nodes in G .
2. Any alternate path for the same source and destination as P that goes through one or more nodes in $G' - G$ cannot be shorter than P .

Hence, there exists a path in G' that is longer than or equal to $7D/4 - 1$. \square

5 Conclusion

Ružička used a graph that has two flaps [5], and we use here one that has three flaps; looking at the way we proved the bound, however, using four or more flaps might not give rise to a better bound. The graph we used (for the one with interval labels only) has eight layers. In fact, we could have made it six layers, but then the proof (of Theorem 3.1) would become more complicated (and less interesting). On the other hand, if we made $k \geq 2$ instead of 3, the proof would become simpler. We did not consider the case of linear interval routing (*i.e.*, using non-cyclic labels) which should be just a simple extension of what we have done: its bound is expected to be worst (bigger) than $7D/4 - 1$ because of the reduced flexibility of the interval labels. One obvious

future direction is to consider multiple labels per edge. Intuitively, the more labels are assigned to an edge, the better the chance of finding optimal or near-optimal labelings for arbitrary graphs. Using a strategy similar to the one used here, we have proved a lower bound of $5D/4 - 1$ (or $1.25D - 1$) for 2-label interval routing in arbitrary graphs [7].

References

- [1] E.M. Bakker, J. van Leeuwen, and R.B. Tan, Linear Interval Routing Schemes. Technical Report RUU-CS-91-7, Department of Computer Science, Utrecht University (1991).
- [2] P. Fraigniaud and C. Gavoille, Interval Routing Schemes. Research Report No. 94-04, Laboratoire de L'Informatique du Parallélisme, Ecole Normale Supérieure de Lyon (1994).
- [3] J. Hinton and A. Pinder, *Transputer Hardware and System Design*. Prentice-Hall, New York (1993).
- [4] M.D. May, P.W. Thompson, and P.H. Welch (eds.), *Networks, Routers and Transputers*. IOS Press, Amsterdam (1993).
- [5] P. Ružička, A note on the efficiency of an interval routing algorithm. *The Computer Journal* **34** (1991) 475–476.
- [6] N. Santoro and R. Khatib, Labelling and implicit routing in networks. *The Computer Journal* **28** (1985) 5–8.
- [7] S.S.H. Tse and F.C.M. Lau, On 2-label Interval Routing. Technical Report, Department of Computer Science, The University of Hong Kong (1994).
- [8] J. van Leeuwen and R.B. Tan, Interval routing. *The Computer Journal* **30** (1987) 198–307.

X09000732



P 004.65 T88
Tse, Savio S. H.
A lower bound for interval
routing in general networks
Hong Kong : Department of
Computer Science, Faculty of
Engineering, University of

