

CLINICAL REPORT
OF THE
TSAN YUK HOSPITAL
FOR THE YEAR
1960



ISSUED BY THE
DEPARTMENT OF OBSTETRICS AND GYNAECOLOGY
OF THE
UNIVERSITY OF HONG KONG

PRINTED BY
THE SOUTH CHINA MORNING POST, LIMITED
HONG KONG

CONTENTS

	<i>Page</i>
I. MEDICAL STAFF	1
II. ABBREVIATIONS	3
III. TABLE OF COMPARATIVE STATISTICS, 1951—1960	4
IV. STATISTICAL SUMMARY	5
V. DETAILED OBSTETRICAL REPORT	

ARRANGEMENT OF TABLES:—

Table: 1. Associated Diseases	6
2. Abortion	8
3. Cardiac Disease	9
4. Toxaemia of Pregnancy	11
5. Pre-eclampsia	12
6. Eclampsia	14
7. Unclassified Toxaemia	16
8. Recurrent Toxaemia	18
9. Essential Hypertension	20
10. Chronic Nephritis	21
11. Persistent Posterior Position of the Occiput and Transverse Arrest	22
12. Breech Deliveries	23
13. Face and Brow Presentation	26
14. Transverse and Oblique Lie	28
15. Compound Presentation	30
16. Prolapse and Presentation of Cord	31
17. Hydramnios	32
18. Prolonged Labour	33
19. Labour Following Previous Caesarean Section	38
20. Induction of Labour	39
21. Forceps Deliveries	41
22. Version in Labour	42
23. Caesarean Sections	43
24. Rupture of Uterus	45
25. Contracted Pelvis	47

CONTENTS *(Contd.)*

	<i>Page</i>
26. Antepartum Haemorrhage	64
27. Accidental Antepartum Haemorrhage ..	65
28. Placenta Praevia	69
29. A.P.H. of Unknown Origin	72
30. A.P.H.—Extra-Placental Causes	73
31. Post-partum Haemorrhage	74
32. Manual Removal of Placenta	75
33. Tumours Associated with Pregnancy, Labour or Puerperium	76
34. Multiple Pregnancy	77
35. Puerperal Pyrexia	82
VI. DETAILS OF MATERNAL DEATHS	84
VII. DETAILED PAEDIATRIC REPORT	
ARRANGEMENT OF TABLES:—	
36. Summary of Foetal Loss	85
37. Premature Live-born Infants	87
38. Congenital Malformations	88
39. Birth Trauma	89
40. Neo-natal Infections	90
41. Haemolytic Diseases	90
VIII. X-RAY DEPARTMENT STATISTICS	91

MEDICAL STAFF OF THE TSAN YUK HOSPITAL

Daphne Chun, O.B.E., F.R.C.S. (Ed.), F.R.C.O.G.

Professor of Obstetrics & Gynaecology, University of Hong Kong.
Consulting Obstetrician & Gynaecologist to the Government.

H. Y. Cheng, M.B., B.S.

Medical Superintendent.

C. A. Braga, M.B., B.S., M.R.C.O.G.

D.P.C. Chan, M.B., B.S., F.R.C.S. (Ed.), F.R.F.P.S. (Glas.), M.R.C.O.G.

Chew Wei, M.B., B.S., M.R.C.O.G.

P.S. Kan, M.B., B.S., M.R.C.O.G.

*Dorothy Kwan, M.B., B.S., M.R.C.O.G.

*Y. K. Tsao, M.B., B.S., M.R.C.O.G.

Lecturers, University of Hong Kong.

Chung Ho Kei, M.B., B.S. (July—Dec.)

Therese Lu, M.B., B.S. (Jan.—Aug.)

Soo Hung Nin, M.B., B.S. (July—Dec.)

Assistant Lecturers, University of Hong Kong.

Chung Ho Kei, M.B., B.S.

Hui Chiu Ling, Jane, M.B., B.S.

Lee Kin Hung, M.B., B.S.

A. C. de B. Lopes, M.B., B.S.

Soo Hung Nin, M.B., B.S.

(Jan.—June)

Clinical Assistants, University of Hong Kong.

Joseph Cheung, M.B., B.S. (Jan.—Dec.)

Lee Kin Hung, M.B., B.S., (July—Dec.)

A. C. de B. Lopes, M.B., B.S. (July—Dec.)

Medical Officers.

* Part-time.

MEDICAL STAFF OF THE TSAN YUK HOSPITAL (Continued)

Chan Hip Sing, M.B., B.S.

Chan Ming Yee, M.B., B.S.

Chow Tsin-Si, Therese, M.B., B.S.

Kwa Will Pong, Mathew, M.B., B.S.

Lau Chung Hi, Denis, M.B., B.S.

Leung Ding Bong, Ronald, M.B., B.S.

Tam Cherk Shing, M.B., B.S.

Wong Chik Sun, M.B., B.S.

(Jan.—June)

Lee Tin-Sang, M.B., B.S.

Ma Kin Fun, M.B., B.S.

Mok Chan Hung, M.B., B.S.

Su Chung Yee, M.B., B.S.

Thom Fong Yan, M.B., B.S.

Wong Chun Lit, M.B., B.S.

Wong Kam Fun, M.B., B.S.

(July—Dec.)

House-Officers.

ABBREVIATIONS

A.C.H.	After coming head.
A.P.H.	Antepartum haemorrhage.
A.P.O.	Antero-posterior outlet.
A.R.M.	Artificial rupture of membranes.
B.B.A.	Born before arrival.
B.W.	Birth weight.
B.P.	Blood pressure.
C.	Child.
C-P disproportion	Cephalo-pelvic disproportion.
C.S.	Caesarean Section.
C.C.S.	Classical Caesarean Section.
D.	Died.
D.A.A.	Discharged against advice.
E.U.A.	Examination under anaesthesia.
F.D.	Fully dilated.
G.A.	General anaesthesia.
I.D.I.	Induction delivery interval.
I.P. Eclampsia	Intra-partum eclampsia.
I.U.D.	Intra-uterine death.
I.V.	Intravenous.
I.V.P.	Intravenous pyelogram.
L.	Live.
L.B.	Live birth.
L.S.C.S.	Lower segment Caesarean Section.
M.	Mother.
Mac.	Macerated.
N.N.D.	Neo-natal death.
O.A.	Occipito-anterior.
O.B.E.	Oil, bath and enema (Medical induction of labour).
O.C.	Obstetrical conjugate.
O.P.	Occipito-posterior.
P.E.	Pre-eclampsia.
P.N.D.	Peri-natal death.
P.P. Eclampsia	Post-partum eclampsia.
P.P.H.	Post-partum haemorrhage.
P.P.S.	Post-partum sterilization.
Q.M.H.	Queen Mary Hospital.
S.B.	Stillbirth.
Spont.	Spontaneous delivery.
T.B.	Tuberculosis.
T.D.	Transverse diameter.
V.D.R.L.	Venereal disease research laboratory.
Wks.	Weeks.

STATISTICAL SUMMARY

Adult patients admitted	7,251
Patients delivered in hospital	6,266
Primigravidae	2,525
Multiparae	3,741
Patients admitted after delivery	32
Primigravidae	2
Multiparae	30
Total patients delivered	6,298
Abortions	17
Patients discharged undelivered	119
Maternal deaths (Died delivered ... 1 } (Died undelivered . 1 }	2
Maternal death rate (per 1,000 live & stillbirths)	0.32
Infants born in hospital	6,333
Infants born before admission (B.B.A.)	32
Total infants born	6,365
Stillbirths	61
Stillbirth rate (per 1,000 live & stillbirths)	9.58
Total number of live born infants	6,304
Neo-natal deaths	78
Neo-natal death rate (per 1,000 live births)	12.37
Perinatal deaths	135
Perinatal death rate (per 1,000 live & stillbirths)	21.21
Ante-natal clinic:—	
New patients	6,665
Re-attendances	28,275
Total	34,940
Post-natal clinic:—	
Total attendances	3,653

TABLE I

**ASSOCIATED DISEASES & CONDITIONS NOT DETAILED
ELSEWHERE**

Diseases of Blood:—	
Iron deficiency anaemia	1,552
Megaloblastic anaemia	1
Thalassaemia minor	2
Bleeding tendency	1
Haemolytic jaundice	1
Thrombocytopaenic purpura	1
Diseases of Cardiovascular System:—	
Dextrocardia	1
Diseases of Respiratory System:—	
Pulmonary T.B.	236
Bronchial asthma	11
Chronic bronchitis with emphysema	1
Pulmonary emphysema	1
Bronchiectasis	6
Upper respiratory tract infection	2
Previous thoracoplasty	1
Benign neoplasm of lung	1
Pneumonia	1
Diseases of Nervous System:—	
Schizophrenia	9
Mental defect	1
Mental illness	1
Bell's palsy	1
Idiopathic epilepsy	2
Median carpo-tunnel syndrome	1
Postphrenic crush diaphragmatic paralysis	1
Parasitic Diseases:—	
Ankylostomiasis	7
Ascariasis	3
Trichomoniasis	58
Moniliasis	50
Diseases of Reticulo-endothelial System:—	
Hepatosplenomegaly	1
Previous splenectomy	1
Cervical lymphadenitis	1
Diseases of Gastro-intestinal Tract:—	
Granulomatous epulis	1
Cardiospasm	1
Carcinoma of liver	1
Cholangiohepatitis	6
Acute appendicitis	1
Ischio-rectal abscess	1
Gastro-enteritis	3
Diseases of Genito-urinary System:—	
Urinary tract infection	6
Chronic pyelonephritis	1
Left hydronephrosis with ureteric stone	1
Post renal uraemia	1
Chyluria	2
Haematuria of unknown cause	2
Cystitis	1
Tuberculosis of urinary tract	1
Paraurethral cyst	1
Condyloma accuminata	3

Previous ectopic pregnancy	2
Thick intact hymen	1
Previous Manchester operation	2
Bartholin cyst	1
Cervical polyp	5
Incompetent os (habitual abortion)	1
Prolapse of uterus	8
Previous ruptured uterus	1
Previous stab wound of uterus	1
Lipoma of labium majus	1
Infectious Diseases:—	
Syphilis	114
Bacillary dysentery	11
Virus hepatitis	4
Leprosy	2
Scarlet fever	1
Poliomyelitis	1
Diseases of Endocrine Glands:—	
Thyrotoxicosis	17
Nontoxic diffuse goitre	1
Nontoxic nodular goitre	1
Diabetes mellitus	4
Prediabetic state	1
Cushing's syndrome	1
Diseases of Bones & Joints:—	
T.B. Hip	3
T.B. Spine	1
Old T.B. sacro-iliac joint & ilium	1
Ankylosed left hip	1
Right cervical rib	1
Osteomyelitis of jaw	1
Diseases of Ears, Nose & Throat:—	
Chronic sinusitis	1
Haemolytic streptococcal infection of throat	1
Avitaminosis:—	3
Diseases of Eye:—	
Conjunctivitis	2
Diseases of Skin:—	
Tinea infection	2
Chronic eczema	1
Lichen simplex	1
Contact dermatitis	4
Neurodermatitis	1
Miscellaneous:—	
Jaundice of unknown cause	1
Infected insect bite	1
Streptococcal infection of foot	1
Abscess of finger	1
Carcinoma of breast	2
Breast abscess	2
Fibroadenoma of breast	1
Hyperemesis gravidarum	1

TABLE 2
ABORTION

Total No. of Cases	17
Booked	14
Non-booked	3
Primiparae	5
Multiparae	12
Incidence	0.27%
Maternal Death	—
On admission:—	
Threatened	—
Inevitable	9
Missed	8
(Habitual)	1
Outcome—	
Complete	14
Incomplete	3

TABLE 3

CARDIAC DISEASE

Total No. of Cases	60
Booked	58
Non-booked	2
Primiparae	26
Multiparae	34
Incidence	0.95%
Maternal death	1 (undelivered)
S. B.	—
N. N. D.	—
P. N. D.	—
Abortion	1

ANATOMICAL LESIONS:—

A. Rheumatic heart disease	52	
(a) Mitral lesion with regular rhythm	43	} 45
Mitral lesion with auricular fibrillation	2	
(b) Mitral and aortic lesions with regular rhythm	6	} 7
Mitral and aortic lesions with auricular fibrillation	1	
B. Congenital heart disease	8	
(a) Atrial septal defect	2	
(b) Ventricular septal defect	1	
(c) Patent ductus arteriosus	2	
(d) Pulmonary stenosis	2	
(e) Dextrocardia	1	

ASSOCIATED COMPLICATIONS:—

Anaemia	7
Toxaemia	1
Upper respiratory infections	2
Premature labour	7
Contracted pelvis	5

CARDIAC DISEASE (Continued)

FUNCTIONAL GRADING AT THE TIME OF DELIVERY:—

Functional Class	Rheumatic		Congenital				
	Mitral	Mitral + Aortic	Atrial septal defect	Ventricular septal defect	Patent ductus arteriosus	Pulmonary stenosis	Dextrocardia
I .. .	21	1	1	1	2	2	1
II	19	—	—	—	—	—	—
III	1	2	—	—	—	—	—
IV	4	4	1	—	—	—	—

METHOD OF DELIVERY AND FOETAL LOSS:—

Method of delivery	No of cases	S.B.	N.N.D.	P.N.D.
Spontaneous	27	—	—	—
Low forceps	22	—	—	—
Mid forceps	1	—	—	—
Breech delivery	1	—	—	—
C. S.	6	—	—	—
Caesarean hysterectomy	1	—	—	—
Total	58*			

MATERNAL DEATH:—	1
Undelivered	1
Before labour	1
In labour	—
Delivered	—

* 1 mother died undelivered before labour
 1 mother had an abortion at the 28th week } Both are non-booked cases.

TABLE 4

TOXAEMIA OF PREGNANCY

	<i>No. of cases</i>
Pre-eclampsia	358 (70.33%)
Eclampsia	13 (2.55%)
Unclassified Toxaemia	115 (22.59%)
Recurrent Toxaemia	12 (2.36%)
Essential Hypertension	9 (1.77%)
Chronic Nephritis	2 (0.39%)
TOTAL	509
INCIDENCE	8.08%
Total No. of babies	534
Multiple Pregnancies	24 (1 set of triplets)
Maternal death	—
S.B.	9 (1.69%)
N.N.D.	12 (2.29% of L.B.)
P.N.D.	21 (3.93%)

PRE-ECLAMPSIA (Continued)

BIRTH WEIGHT & FOETAL LOSS:—

<i>Birth Weight</i>	<i>No. of cases</i>	<i>S.B.</i>	<i>N.N.D.</i>	<i>P.N.D.</i>
(A) Spontaneous labour				
Under 1,500	5	1	3	4
1,501 — 2,000	12	1	3	4
2,001 — 2,500	42	1	2	3
2,501 — 3,000	111	3	1	4
3,001 — 3,500	109	—	1	1
3,501 — 4,000	44	—	—	—
4,001 & over	5	—	—	—
(B) Induced labour				
Under 1,500	—	—	—	—
1,501 — 2,000	—	—	—	—
2,001 — 2,500	7	1	—	1
2,501 — 3,000	16	—	1	1
3,001 — 3,500	18	1	—	1
3,501 — 4,000	11	—	—	—
4,001 & over	—	—	—	—

INDICATIONS FOR C.S.:—

	<i>No. of cases</i>
Foetal Distress	8
Placenta Praevia	5
Failed Induction	3
Failed Trial of Labour (Contracted pelvis)	2
Deformed Pelvis	2
Contracted Pelvis	11
Accidental Haemorrhage	3
Prolapse of Cord	1
Elderly Primigravida	2
Prolonged Labour	2
Uterine Inertia	5
Cervical Dystocia	1
Breech & Uterine Abnormality	1
Unripe Cervix	2
Previous Caesarean Section	2
Total	50*

* There were 3 sets of twins.

TABLE 6
ECLAMPSIA

Total No. of cases	13
Booked	11
Non-booked	2
Primiparae	9
Multiparae	4
Incidence:—Over all	0.21%
Of toxæmia	2.55%
Total No. of babies	13
Multiple pregnancies	—
Maternal death	—
S.B.	—
N.N.D.	—
P.N.D.	—

Reg. No.	Previous Pregnancies		Maturity	Condition on admission	If in labour	Bejore admission	Fits			Total	Proteinuria		Quantity in first 24 hrs.	Oedema
	Age	28 wks					After 28 wks	Ante-partum	Intra-partum		Post-partum	Maximum		
PRIMIPARA														
BOOKED														
317	26	—	—	40	Good	Yes	—	—	—	1	1	+	—	+
411	24	—	—	39	Unconscious	No	2	2	—	—	4	+++	—	+++
3264	21	—	—	38	Good	Yes	—	—	—	1	1	+	—	++
3877	23	—	—	39	„	„	—	—	1	1	2	++	—	++
4047	23	—	—	38	Fair	„	—	—	—	1	1	+++	—	++
4785	16	—	—	39	Unconscious	No	—	1	—	—	1	+	—	++
4826	26	—	—	40	Good	Yes	—	—	1	1	2	+++	—	+
5000	27	—	—	38	„	„	—	—	—	1	1	—	—	—
NON-BOOKED														
3015	22	—	—	37	Good	No	—	1	—	—	1	+++	—	+++
MULTIPARA														
BOOKED														
3076/59	39	—	5	?	Unconscious	B.B.A.	8	—	—	14	22	+++	—	+++
2741/60	24	1	1	38	Good	Yes	—	—	—	2	2	+	—	—
3046	43	—	3	37	„	No	—	—	1	—	1	+++	—	+++
NON-BOOKED														
1456	25	—	1	38	With a convulsion	No	2	2	—	—	4	—	—	3/4 gms. +-

Blood pressure Before 20th wk	Highest	No. days in hospital	If labour induced state method	Method of delivery	Treatment	Result		Weight of child (grams)	Remarks
						M.	C.		
115/75	180/100	—	—	Spont.	Pentothal 1.7 gms.	L.	L.	2225	Ante-natal visits — E.P. normal. In labour—B.P. 180/ 100. Discharge — B.P. 110/70.
—	180/125	32	—	Low forceps	" 7.1 gms.	L.	L.	2540	
—	170/100	1	—	Spont.	" 1 gm.	L.	L.	3090	
—	170/120	—	—	Low forceps	" 4.1 gms.	L.	L.	2863	Anaemia.
—	190/120	4	—	"	" 0.8 gms.	L.	L.	2977	Transferred to Q.M.H. because of subarachnoid hae- morrhage, 4th day after delivery.
—	165/110	—	—	L.S.C.S.	" 2.45 gms.	L.	L.	3629	Contracted pelvis.
110/75	170/120	14	—	"	" 5.3 gms.	L.	L.	2551	
—	165/100	14	—	Spont.	" 2 gms.	L.	L.	2381	
—	190/100	16	—	Breech extraction	Pentothal 2.5 gms.	L.	L.	2665	
—	220/105	30	—	Spont.	Pentothal 4.35 gms.	L.	L.	Unknown	B.B.A. P.P.S.
—	190/120	14	—	"	" 4.5 gms.	L.	L.	2495	
—	240/130	2	Stripping of membranes & A.R.M.	"	" 3.5 gms.	L.	L.	1616	On discharge—B.P. 170/100. To at- tend medical clinic.
—	160/100	7	—	Low forceps	Pentothal 5.15 gms.	L.	L.	2296	

TABLE 7

UNCLASSIFIED TOXAEMIA

Total No. of cases	115
Booked	94
Non-booked	21
Primiparae	39
Multiparae	76
Incidence	Overall 1.83%
	Of toxæmia 22.59%
Total No. of babies	118
Multiple pregnancies	3
Maternal death	—
S.B.	1 (0.85%)
N.N.D.	—
P.N.D.	1 (0.85%)

CLASSIFICATIONS:—

Mild—B.P. 140/90 — 160/100.

Severe—B.P. more than 160/100.

MANAGEMENT & FOETAL LOSS:—

Management	No. of cases	Booked			Non-booked			
		S.B.	N.N.D.	P.N.D.	No. of cases	S.B.	N.N.D.	P.N.D.
(A) Spontaneous labour								
(1) Spont. delivery								
Vertex	60	1	—	1	9	—	—	—
Breech	1	—	—	—	2	—	—	—
(2) Forceps	11	—	—	—	3	—	—	—
(3) C.S.	10	—	—	—	—	—	—	—
(B) Induced labour								
(1) Spont. delivery								
Vertex	10	—	—	—	6	—	—	—
Breech	1	—	—	—	—	—	—	—
(2) Forceps	—	—	—	—	4	—	—	—
(3) C.S.	1	—	—	—	—	—	—	—
TOTAL	94	1	—	1	24	—	—	—

UNCLASSIFIED TOXAEMIA (Continued)

BIRTH WEIGHT & FOETAL LOSS:—

<i>Birth Weight</i>	<i>No. of cases</i>	<i>S.B.</i>	<i>N.N.D.</i>	<i>P.N.D.</i>
(A) Spontaneous labour				
Under 1,500	1	1	—	1
1,501—2,000	1	—	—	—
2,001—2,500	8	—	—	—
2,501—3,000	36	—	—	—
3,001—3,500	38	—	—	—
3,501—4,000	11	—	—	—
4,000 & over	1	—	—	—
(B) Induced labour				
Under 1,500	—	—	—	—
1,501—2,000	—	—	—	—
2,001—2,500	5	—	—	—
2,501—3,000	11	—	—	—
3,001—3,500	5	—	—	—
3,501—4,000	—	—	—	—
4,000 & over	1	—	—	—

INDICATIONS FOR C.S.:—

	<i>No. of cases</i>
Elderly Primigravida	3
Foetal Distress	2
Contracted Pelvis & Failed Forceps	1
Prolonged Labour	2
Placenta Praevia & Contracted Pelvis	1
Previous Caesarean Section	2
TOTAL	<u>11</u>

TABLE 8

RECURRENT TOXAEMIA

Total No. of cases	12
Booked	12
Non-booked	—
Primiparae	—
Multiparae	12
Incidence	Overall 0.19%
		Of toxæmia 2.36%
Total No. of babies	12
Multiple pregnancies	—
Maternal death	—
S.B.	—
N.N.D.	—
P.N.D.	—

CLASSIFICATIONS:—

Mild—B.P. 140/90 — 160/100.

Severe—B.P. more than 160/100.

MANAGEMENT & FOETAL LOSS:—

Management	No. of cases	Mild			Booked			Severe		
		S.B.	N.N.D.	P.N.D.	No. of cases	S.B.	N.N.D.	P.N.D.		
(A) Spontaneous labour										
(1) Spont. delivery										
Vertex	4	—	—	—	2	—	—	—	—	—
Breech	1	—	—	—	—	—	—	—	—	—
(2) Forceps	1	—	—	—	1	—	—	—	—	—
(3) C.S.	—	—	—	—	—	—	—	—	—	—
(B) Induced labour										
(1) Spont. delivery										
Vertex	2	—	—	—	—	—	—	—	—	—
Breech	—	—	—	—	—	—	—	—	—	—
(2) Forceps	—	—	—	—	—	—	—	—	—	—
(3) C.S.	1	—	—	—	—	—	—	—	—	—
TOTAL	9	—	—	—	3	—	—	—	—	—

RECURRENT TOXAEMIA (Continued)

BIRTH WEIGHT & FOETAL LOSS:—

<i>Birth Weight</i>	<i>No. of cases</i>	<i>S.B.</i>	<i>N.N.D.</i>	<i>P.N.D.</i>
(A) Spontaneous labour				
Under 1,500	—	—	—	—
1,501 — 2,000	—	—	—	—
2,001 — 2,500	1	—	—	—
2,501 — 3,000	5	—	—	—
3,001 — 3,500	2	—	—	—
3,501 — 4,000	1	—	—	—
4,000 & over	—	—	—	—
(B) Induced labour				
Under 1,500	—	—	—	—
1,501 — 2,000	—	—	—	—
2,001 — 2,500	—	—	—	—
2,501 — 3,000	1	—	—	—
3,001 — 3,500	—	—	—	—
3,501 — 4,000	2	—	—	—
4,000 & over	—	—	—	—

INDICATIONS FOR C.S.:—

	<i>No. of cases</i>
C.P. disproportion & previous Caesarean Section	1
Total	<u>1</u>

TABLE 9
ESSENTIAL HYPERTENSION

Reg. No.	Age	Previous Pregnancies Before 28 wks	Asc	Maturity On adm.	At del.	Maximum	On discharge	Oedema	Blood Pressure Before 20th week	Highest	No. of days in hospital	Labour induced	Method of delivery	Result	Weight of child (grams)	Remarks
PRIMIPARA																
BOOKED																
65	34	—	—	31	42	—	—	—	140/100	170/110	77	—	L.S.C.S.	L.	2693	L.S.C.S. was done for borderline pelvis, post-maturity and cervix, not favourable for induction.
MULTIPARA																
BOOKED																
7944/59	42	—	7	32	42	—	—	—	—	170/100	25	Stripping of membranes	Spont.	L.	3060	Refused P.P.S. D.A.A.
691/60	23	—	2	32	36	—	+	—	—	170/110	13	—	L.S.C.S.	L.	2551	1st pregnancy—eclampsia. 2nd pregnancy—severe toxæmia. Treated with serpasil & apresoline. 2 previous C. S. done. P.P.S.
710	41	1	6	38	38	—	—	—	—	190/110	18	—	Spont.	L.	2750	P.P.S.
1097	23	—	1	32	32	—	—	210/100	—	210/100	32	—	"	L.	1673	N.N.D.
1450	41	—	8	28	37	—	+	—	—	150/110	46	—	"	L.	2948	Anæmia. Fibroids of uterus. Referred to Q.M.H. Gyn. unit.
2194	32	2	6	40	40	—	+	130/90	—	165/110	3	O.B.E. & stripping of membranes	"	L.	3515	
2830	42	1	7	33	38	—	—	—	—	180/100	23	—	"	L.	2825	B.P. 140/95 on discharge. Mild accidental hæmorrhage.
3249	40	2	1	38	38	—	—	140/90	—	140/90	—	—	L.S.C.S.	L.	3005	L.S.C.S. was done for cervical dystocia.
<p>Total No. of cases 9</p> <p>Booked 9</p> <p>Non-booked —</p> <p>Primiparæ 1</p> <p>Multiparæ 8</p> <p>Incidence—Over all 0.14%</p> <p>Of toxæmia 1.77%</p> <p>Of babies 9</p> <p>Multiple pregnancies —</p> <p>Maternal death —</p> <p>S.B. —</p> <p>N.N.D. 1 (11.11%)</p> <p>P.N.D. 1 (11.11%)</p> <p>Proteinuria —</p>																

TABLE 10
CHRONIC NEPHRITIS

Reg. No.	Age	Previous Pregnancies Before 28 wks. On Adm.	Maturity On Adm.	At del.	Maximum discharge	Oedema	Blood Pressure Before 20th wk.	Highest	No. of days in hospital	If labour induced	Method of delivery	Treatment	Result M. C.	Weight of child (grams)	Remarks
PRIMIPARA															
BOOKED															
144	23	1	34	36	2 gms.	—	150/110	33	O.B.E.	Spont.	—	—	L. L.	2500	
MULTIPARA															
BOOKED															
2944	38	1	6	33	35	—	160/110	22	—	Spont.	—	—	L. L.	1928	Blood urea (180 mgm.%) 4 years history. P.P.S.
Total No. of cases															
Booked															
Non-booked															
Primiparae															
Multiparae															
Incidence:—															
Over all															
Of toxæmia															
Total No. of babies															
Multiple pregnancies															
Maternal death															
S.B.															
N.N.D.															
P.N.D.															

TABLE 11

**PERSISTENT POSTERIOR POSITION OF THE OCCIPUT
AND TRANSVERSE ARREST**

Total No. of cases	171
Booked	165
Non-booked	6
Primiparae	113
Multiparae	58
Incidence	2.69%
Total No. of babies	171
Multiple pregnancies	5*
Maternal death	—
S. B.	—
N. N. D.	2 (1.17%)
P. N. D.	2 (1.17%)

SUMMARY OF LABOURS:—

Labour more than 24 hours	17
---------------------------	----

METHOD OF DELIVERY:—

Spontaneous delivery (face to pubes)	18
Manual rotation and forceps	72
Face to pubes and forceps	32
Failed manual rotation then C. S.	—
Failed forceps then C. S.	1
C. S.	48
Total	<u>171</u>

TYPES OF PELVIS:—

Gynaecoid	74
Anthropoid	4
Platypelloid	4
Android	1
X-ray not taken	88
Total	<u>171</u>

* Other twin in each case presented normally.

TABLE 12

BREECH DELIVERIES

Total No. of cases	240
Booked	218
Non-booked	22
Primiparae	110
Multiparae	130
Single pregnancies	216
Twin pregnancies	24
Triplet pregnancies	—
Total No. of babies	242
Incidence	3.8%
Caesarean section	56
Incidence	23.33%

UNCOMPLICATED BREECH DELIVERIES

Total No. of cases	100
Booked	95
Non-booked	5
Primiparae	40
Multiparae	60
Maternal death	—
S. B.	—
N. N. D.	1 (1%)
P. N. D.	1 (1%)

TYPE OF BREECH PRESENTATIONS AND FOETAL LOSS:—

<i>Position of legs</i>	<i>No. of cases</i>	<i>Primiparae</i>			<i>No. of cases</i>	<i>Multiparae</i>		
		<i>S.B.</i>	<i>N.N.D.</i>	<i>P.N.D.</i>		<i>S.B.</i>	<i>N.N.D.</i>	<i>P.N.D.</i>
Extended	22	—	—	—	28	—	1	1
Flexed	14	—	—	—	26	—	—	—
Footling	3	—	—	—	3	—	—	—
Not recorded ..	1	—	—	—	3	—	—	—
Total	40	—	—	—	60	—	1	1

BREECH DELIVERIES (Continued)

METHOD OF DELIVERIES, BIRTH WEIGHT AND FOETAL LOSS:—

Method of Deliveries	No. of cases	2501—3500 gms.			No. of cases	3501 gms. & over		
		S.B.	N.N.D.	P.N.D.		S.B.	N.N.D.	P.N.D.
Spontaneous ..	8	—	—	—	6	—	—	—
Assisted	30	—	—	—	8	—	—	—
Assisted with Forceps for A.C.H.	41	—	1	1	—	—	—	—
C. S.	5	—	—	—	2	—	—	—
Total	84	—	1	1	16	—	—	—

INDICATIONS FOR C.S.:—

	No. of cases
Elderly primigravida	2
Large foetus	2
Previous Caesarean Section .. .	3
Total	7

COMPLICATED BREECH DELIVERIES

Total No. of cases	140
Booked	123
Non-booked	17
Primiparae	68
Multiparae	72
Twin pregnancies	24
Total No. of babies	142
Maternal death	1
S. B.	14 (9.86%)
N. N. D.	9 (7.03%)
P. N. D.	22 (15.49%)

ASSOCIATED COMPLICATIONS*

	Primiparae	Multiparae
Contracted pelvis	32	21
Prematurity	24	28
Pre-eclampsia	12	16
Twin pregnancy	6	20
Prolapse of cord	6	6
Placenta praevia	1	4
Accidental haemorrhage .. .	2	3
A. P. H. of unknown origin ..	—	3
Hydramnios	—	3
Macerated foetus	—	3
Uterine inertia	3	—
Eclampsia	2	—
Mitral stenosis	2	—
Prolonged labour	—	1
Postmaturity	1	1
Uterus didelphys	1	1
Uterus bicornis unicollis .. .	1	—
Septate uterus	1	—
Uterine fibroid	—	1

* Some cases have more than one complication.

BRECH DELIVERIES (Continued)

METHOD OF DELIVERIES, BIRTH WEIGHT AND FOETAL LOSS:—

Birth Weight Method of Deliveries	2500 gms. and under			2501—3500 gms.			3501 gms. & over			
	No. of cases	S.B.	N.N.D.	No. of cases	S.B.	N.N.D.	No. of cases	S.B.	N.N.D.	P.N.D.
Spont.	11	6	2	3	—	—	—	—	—	—
Assisted	14	4	2	13	1	—	5	—	—	—
Assisted with forceps for A.C.H.	15	—	3	23	—	—	1	—	—	—
Breech extraction	3	1	1	4	—	—	—	—	—	—
Caesarean Section	9*	2	1	32	—	—	9	—	—	—
Total	52	13	9	75	1	—	15	—	—	—

* Both twins presented by breech delivered by same C. S.

INDICATIONS FOR C.S.:—

	No. of cases
Contracted pelvis	34
Placenta praevia	3
Pre-eclampsia	2
Uterine inertia	2
Eclampsia	1
Prolapse of cord	1
Prolonged labour	1
Premature rupture of membranes	1
Postmaturity	1
Hydramnios, hydrops foetalis	1
Uterus didelphys	1
Uterine fibroid	1
Total	49
Incidence of C. S.	35%

FACE & BROW PRESENTATION (Continued)

Reg. No.	Age	Pregnancies Before After 28 wks. 28 wks.	Maturity	Position	Cause	Treatment	Method of delivery	Result	Weight of child (grams)	Remarks
PRIMIPARA										
BOOKED										
2682	36	—	39	L.M.P.	—	—	Spont.	L.	3062	
3278	25	—	42	L.M.P.	—	—	L.S.C.S.	L.	3472	Diagnosed at 9 cm. dilation. Attempted rotation under G.A. Rotation under G.A. failed.
4044	28	—	40	R.M.P.	—	—	"	L.	3062	
MULTIPARA										
BOOKED										
1324	33	—	37	L.M.A.	—	—	Spont.	L.	2948	
2382	27	—	40	L.M.P.	Anencephaly	—	"	L.	2013	
2650	34	—	39	R.M.A.	—	—	"	L.	2722	
3380	29	—	40	L.M.P.	Anencephaly	—	"	L.	2722	
3488	30	—	41	R.M.P.	"	—	"	L.	3175	
4498	36	—	38	L.M.A.	—	—	Low forceps	L.	2948	
4539	25	—	41	R.M.A.	Anencephaly	—	Spont.	L.	2835	
4694	30	—	45	R.M.P. → M.A.	—	—	Manual rotation & mid forceps	L.	2693	
4698	34	—	32	R.M.A.	Anencephaly	—	Spont.	L.	2381	
4906	36	—	37	L.F.A.	—	—	L.S.C.S.	L.	5600	Contracted pelvis.
NON-BOOKED										
1238	29	—	45	R.M.P.	Borderline pelvis. Big baby	—	L.S.C.S.	L.	4196	

TABLE 14

TRANSVERSE AND OBLIQUE LIE
(in labour)

Total No. of cases	17
Booked	12
Non-booked	5
Primiparae	1
Multiparae	16
Incidence	0.27%
Total No. of babies	17
Multiple pregnancies	3
Maternal death	—
S.B.	2 (11.76%)
N.N.D.	—
P.N.D.	2 (11.76%)

MANAGEMENT & FOETAL LOSS:—

<i>Method of delivery</i>		<i>No. of cases</i>	<i>S.B.</i>	<i>N.N.D.</i>	<i>P.N.D.</i>
C.S.	C.C.S.	2	—	—	—
	L.S.C.S.	13	2	—	2
External	version	1	—	—	—
Internal	version & breech extraction	1	—	—	—
Total		17	2	—	2

ASSOCIATED CONDITIONS & FOETAL LOSS:—

<i>Associated conditions</i>	<i>No. of cases</i>	<i>S.B.</i>	<i>N.N.D.</i>	<i>P.N.D.</i>
Contracted pelvis	2	—	—	—
Prematurity	5	—	—	—
Multiple pregnancies	3	1	—	1
Prolapse of cord	2	1	—	1

TRANSVERSE AND OBLIQUE LIE (Continued)

Reg. No.	Age	Previous Pregnancies Before 28 wks.	After 28 wks.	Maturity	Obstetric complication	Size of os when diagnosed	Treatment	Result M. C.	Weight of child (grams)	Remarks
PRIMIPARA										
BOOKED										
5926	36	—	—	40	Mild P.E.	8 cm.	L.S.C.S.	L.	3544	
MULTIPARA										
BOOKED										
5926	36	1	5	38	—	7 cm.	L.S.C.S.	L.	3289	Prolapse of cord.
1331	37	1	6	42	Post-maturity	4 cm.	"	L.	3402	
1931	38	—	4	34	Anaemia	1 cm.	"	L.	2778	
2168	36	1	4	37	Rupture of uterus	6-7 cm.	L.S.C.S. and total hysterectomy	L.	1701	Prolapse of cord.
3952	47	—	3	39	P.E., Fibroid of uterus	Just admit one finger	L.S.C.S.	L.	3450	
4786	41	1	1	39	Deformed pelvis	Closed	C.C.S.	L.	3402	Previous C.S. done for contracted pelvis.
5136	35	—	6	42	—	7 cm.	L.S.C.S.	L.	3544	Anaemia, P.P.S.
5292	34	—	6	40	—	Closed	"	L.	2948	
5627	27	—	1	41	Septate uterus	—	Effective L.S.C.S.	L.	2863	Patent ductus. Previous C.S. done.
5764	30	—	4	38	Lax abdomen	F.D.	C.C.S.	L.	2722	P.P.S.
5845	30	1	4	32	Contracted pelvis	5 cm.	L.S.C.S.	L.	2438	Anaemia.
NON-BOOKED										
2239	32	—	6	36	—	4 cm.	L.S.C.S.	L.	2410	P.P.S.
5228	37	—	7	35	Prolapse of arm	"	"	L.	1843	A.P.H. of unknown origin.
MULTIPLE PREGNANCY										
NON-BOOKED										
260	26	—	1	38	Twin pregnancy—2nd twin, P.E.	5 cm.	L.S.C.S.	L.	1750	1st twin delivered in maternity home 3 hrs. ago.
1329	38	2	6	32	Twin pregnancy—2nd twin, Accidental haemorrhage.	F.D.	External version & Brech extraction	L.	2381	
2882	31	—	3	42	Twin pregnancy—2nd twin, P.E.	"	Internal version	L.	3033	

TABLE 16

PROLAPSE AND PRESENTATION OF CORD

Total No. of cases	25
Booked	20
Non-booked	5
Primiparae	7
Multiparae	18
Incidence	0.39%
Total No. of babies	25
Multiple pregnancies	2
Maternal death	—
S.B.	2 (8.00%)
N.N.D.	2 (8.70%)
P.N.D.	4 (16.00%)

PRESENTATION AND FOETAL LOSS:—

<i>Presentation</i>	<i>No. of cases</i>	<i>S.B.</i>	<i>N.N.D.</i>	<i>P.N.D.</i>
Vertex	9	1	1	2
Breech	12	—	1	1
Face & brow	—	—	—	—
Transverse Lie	2	1	—	1
Compound	2	—	—	—
Total	25	2	2	4

MANAGEMENT AND FOETAL LOSS:—

<i>Method of Delivery</i>	<i>No. of cases</i>	<i>S.B.</i>	<i>N.N.D.</i>	<i>P.N.D.</i>
Spont.	5	1	1	2
Forceps	7	—	1	1
Breech extraction	2	—	—	—
Int. version and breech extraction	1	—	—	—
C.S.	10	1	—	1
Total	25	2	2	4

TABLE 17

HYDRAMNIOS

Total No. of cases	13
Booked	9
Non-booked	4
Primiparae	1
Multiparae	12
Incidence	0.21%
Total No. of babies	13*
Multiple pregnancy	1
Maternal death	—
S.B.	5 (38.46%)
N.N.D.	4 (50.00%)
P.N.D.	9 (69.23%)
Cases associated with foetal abnormalities	8 (61.54%)

* Only one twin was associated with hydramnios.

TABLE 18

PROLONGED LABOUR

(Labour lasting more than 24 hours in primiparae, and more than 18 hours in multiparae.)

Total No. of cases	65
Booked	63
Non-booked	2
Primiparae	57
Multiparae	8
Incidence	1.03%
Total No. of babies	66
Multiple pregnancies	1
Maternal death	—
S.B.	2 (3.03%)
N.N.D.	1 (1.56%)
P.N.D.	3 (4.55%)

AGE GROUP & FOETAL LOSS:—

Age group & foetal loss	No. of cases	Primiparae			Multiparae		
		S.B.	N.N.D.	P.N.D.	S.B.	N.N.D.	P.N.D.
20 & under	3	—	—	—	—	—	—
21 — 25	20	1	—	1	—	—	—
26 — 30	26	—	1	1	—	—	—
31 — 35	8	1	—	1	—	—	—
36 & over	—	—	—	—	1	—	—

METHOD OF DELIVERY AND FOETAL LOSS:—

Method of delivery	No. of cases	Primiparae			Multiparae		
		S.B.	N.N.D.	P.N.D.	S.B.	N.N.D.	P.N.D.
Spontaneous	3	—	—	—	2	—	—
Low forceps	19	—	—	—	—	—	—
Mid forceps	7	—	—	—	1	—	—
C. S.	29*	2	1	3	5	—	—
Total	58	2	1	3	8	—	—

* A pair of twins delivered by Caesarean Section.

CAUSES OF DELAY AND FOETAL LOSS:—

Cause	No. of cases	Primiparae			Multiparae		
		S.B.	N.N.D.	P.N.D.	S.B.	N.N.D.	P.N.D.
Contracted pelvis	2	—	—	—	1	—	—
Uterine inertia	56*	2	1	3	6	—	—
Malpresentation	—	—	—	—	1	—	—
Total	58	2	1	3	8	—	—

* Including a pair of twins.

PROLONGED LABOUR (Continued)

Reg. No.	Age	Maturity	Height (inches)	Previous Pregnancies		Position of foetus		Other obstetric abnormalities	Cause of delay as diagnosed	Time of rupture of membranes i.e. hours before delivery	Duration of labour (hrs.)		
				Before 28 wks.	After 28 wks.	At onset of labour	At delivery				1st stage	2nd stage	3rd stage
PRIMIPARA													
BOOKED													
8097/59	29	39	63½	--	--	L.O.P.	L.O.A.	—	Uterine inertia	26:45	27:15	1:00	0:15
8166/59	27	40	57¾	--	--	R.O.P.	R.O.P.	—	—	31:40	31:00	—	—
14/60	23	39	59¼	—	—	L.O.P.	L.O.A.	Contracted pelvis	Contracted pelvis	13:55	30:00	1:55	0:05
25	30	39	56½	—	—	R.O.P.	R.O.P.	—	—	25:26	24:00	—	—
506	24	40	60	—	—	L.O.A.	L.O.A.	—	Uterine inertia	1:35	33:30	0:40	0:10
563	24	36	59	—	—	R.O.A.	R.O.A.	—	—	13:55	28:30	0:25	0:15
613	29	40	60¼	—	—	R.O.P.	R.O.P.	—	—	33:51	33:00	—	—
1075	29	41	56½	—	—	L.O.A.	L.O.A.	P.E.	—	12:00	29:41	0:03	0:03
1340	17	38	60	—	—	R.O.A.	R.O.A.	P.E.	—	5:55	29:05	0:15	0:10
1348	22	40	58¾	—	—	R.O.P.	R.O.P.	—	—	50:10	24:35	1:20	0:05
1370	31	41	60	—	—	L.O.A.	L.O.A.	—	—	80:15	31:10	0:05	0:05
1388	25	38	60	—	—	R.O.A.	R.O.A.	—	—	18:20	25:30	0:15	0:05
1409	29	41	59	—	—	L.O.T.	L.O.A.	—	—	6:45	25:30	—	—
1413	30	40	62	—	—	L.O.P.	L.O.P.	—	—	—	26:00	—	—
1440	29	41	61	—	—	R.S.A.	R.S.A.	—	—	11:35	25:00	—	—
1467	29	40	62½	—	—	R.O.P.	R.O.A.	—	—	2:50	25:55	0:10	0:05
1486	23	40	62½	—	—	R.O.T.	R.O.A.	—	—	7:00	37:00	—	—
1499	32	42	57½	—	—	R.O.T.	R.O.T.	Contracted pelvis	—	20:00	25:00	—	—
1817	26	40	61½	—	—	R.O.P.	R.O.P.	—	—	17:00	37:00	—	—
1895	26	39	58½	—	—	R.O.A.	R.O.A.	—	—	27:50	24:40	0:10	0:05
1965	27	40	60¼	—	—	L.O.A.	L.O.A.	Hydramnios	—	32:00	39:00	—	—
2083	35	40	59¾	—	—	L.O.A.	R.O.A.	Contracted pelvis	—	29:30	28:05	0:15	0:05
2273	27	37	61¼	—	—	R.O.A.	R.O.A.	—	—	4:10	24:45	0:25	0:05
2288	30	29	59¾	—	—	L.O.P.	L.O.T.	—	—	5:00	27:00	—	—
2393	28	40	63	—	—	L.O.P.	L.O.P.	—	—	5 days	41:50	0:25	0:05
2607	20	37	NS	—	—	R.O.A.	R.O.A.	—	—	4:00	27:40	0:20	0:05
2749	27	41	60½	—	—	L.O.P.	L.O.P.	—	—	3:00	26:00	—	—
2984	22	41	NS	—	—	L.O.P.	L.O.P.	P.E.	—	16:00	31:00	—	—
3024	35	41	59	—	—	R.O.P.	R.O.T.	—	—	7:30	32:00	—	—
3068	32	38	64	—	—	R.O.P.	R.O.P.	P.E.	—	31:30	24:12	—	—
3226	23	42	61¼	—	—	L.O.A.	L.O.A.	Uterus bicornis with complete septum	—	42:00	30:00	—	—
3526	28	37	57¾	—	—	R.O.P.	R.O.P.	Contracted pelvis.	—	12:00	> 24:00	—	—
3556	24	40	62¼	—	—	L.O.A.	L.O.A.	P.E.	—	7:00	25:55	0:25	0:05
3918	28	38	NS	—	—	L.O.A.	L.O.A.	—	—	17:00	> 24:00	—	—
3923	29	42	60½	—	—	L.O.T.	L.O.T.	P.E.	—	29:00	> 24:00	—	—
4081	29	43	59¼	—	—	R.O.T.	R.O.T.	P.E.	—	13:00	37:00	—	—
4198	24	40	56½	—	—	R.O.P.	R.O.P.	—	—	25:00	25:00	—	—
4262	25	39	60½	—	—	L.O.T.	L.O.T.	Contracted pelvis	—	2 days	29:00	—	—
4450	19	39	60	—	—	L.O.T.	L.O.A.	—	—	20:00	28:00	0:50	0:15
4802	29	38	57	—	—	R.O.P.	R.O.T.	—	—	12:30	25:00	0:05	0:10
4859	23	40	59½	—	—	L.O.A.	L.O.A.	—	—	4:00	26:00	0:15	0:10
4869	25	40	62	—	—	R.O.T.	R.O.T.	—	—	21:00	28:30	0:35	0:05
4950	24	36	60	—	—	L.O.A.	L.O.A.	—	—	11:00	24:55	0:15	0:10
4952	28	40	59	—	—	R.O.P.	R.O.P.	—	—	17:15	27:15	0:12	0:05
5005	24	39	54¾	—	—	L.O.T.	L.O.A.	Contracted pelvis	—	2:00	25:45	0:40	0:05
5027	24	40	59	—	—	L.O.A.	L.O.P.	—	—	?	36:00	—	—
5060	21	42	59	—	—	L.O.P.	L.O.P.	—	—	14:00	25:00	—	—
5074	23	37	61½	—	—	R.O.P.	P.O.P.	P.E.	—	13:00	28:30	0:20	0:01
5087	28	35	56¾	—	—	L.O.A.	L.O.A.	—	—	1:50	31:40	0:15	0:05

Treatment	Method of delivery	P.P.H.	Morbidity	Weight of child (grams)	Result		Child		Remarks
					M.	C.	Trauma	Resuscitation Yes/No	
Sedation	Spont.	—	—	2750	I.	L.	—	—	
—	L.S.C.S.	—	—	2835	L.	I.	—	—	
—	Manual rotation & mid forceps	—	—	2760	I.	I.	—	—	
—	L.S.C.S.	—	—	2665	L.	L.	—	—	
Sedation	Spont.	—	—	2780	I.	I.	—	—	Anaemia.
I.V. drip	Low forceps	—	—	2900	L.	I.	—	—	
—	L.S.C.S.	—	—	3000	L.	L.	—	—	
—	Low forceps	—	—	3202	L.	L.	—	—	
—	Spont.	—	—	3005	L.	I.	—	—	
—	Mid forceps	—	—	2920	L.	L.	—	—	Anaemia.
—	Low forceps	—	—	3549	L.	L.	—	—	
I.V. drip	"	—	—	3062	I.	L.	—	—	
I.V. drip & sedation	L.S.C.S.	—	—	3232	L.	L.	—	—	
"	"	—	—	2920	L.	I.	—	—	
"	"	—	—	3572	L.	I.	—	—	
"	Low forceps	—	—	3175	I.	I.	—	—	
"	L.S.C.S.	—	—	3402	I.	I.	—	—	
"	"	—	—	3033	I.	S.b.	—	—	
Pethidine, I.V. drip	"	—	—	3175	I.	I.	—	—	
—	Low forceps	—	—	2807	I.	L.	—	—	
Pitocin drip & sedation	L.S.C.S.	—	—	3799	I.	N.D.	—	—	
—	Mid forceps	—	—	2948	I.	L.	—	—	
—	Low forceps	—	—	2402	L.	L.	—	—	3° tear repaired
—	L.S.C.S.	—	—	3062	I.	L.	—	—	
—	Low forceps	—	—	2750	I.	I.	—	—	Anaemia.
Pethidine, I.V. drip	Mid forceps	—	—	3572	I.	L.	—	—	
"	L.S.C.S.	—	—	2977	L.	L.	—	—	
"	"	—	—	2920	L.	L.	—	—	
"	"	—	—	3629	L.	L.	—	—	Syphilis.
I.V. drip & sedation	"	—	—	2835	L.	L.	—	—	Anaemia. Fibroid.
Pethidine, I.V. drip	"	—	1	3005	L.	Mac.	—	—	Absence of left kidney.
I.V. drip & sedation	"	—	5	2863	L.	L.	—	Yes	
Sedation	Low forceps	—	—	3374	L.	L.	—	—	Trichomonas.
I.V. drip & sedation	L.S.C.S.	—	—	3118	L.	L.	—	Yes	
"	"	—	—	3175	L.	L.	—	—	
Pethidine, I.V. drip	"	—	—	2835	L.	L.	—	—	Anaemia.
I.V. drip & sedation	"	—	—	2665	L.	L.	—	—	
"	"	—	—	3232	L.	L.	—	—	
"	Low forceps	—	—	3062	L.	L.	—	Yes	
Pethidine, I.V. drip	"	—	—	2778	L.	L.	—	—	
"	"	—	—	3147	L.	L.	—	—	
"	"	—	—	3090	L.	L.	—	—	
"	"	—	—	3260	I.	L.	—	—	Pulmonary T.B.
"	Mid forceps	—	—	3260	L.	L.	—	—	
"	Low forceps	—	—	2892	L.	L.	—	—	
I.V. drip	L.S.C.S.	—	—	2892	L.	L.	—	—	
Pethidine, I.V. drip	"	—	—	3317	L.	L.	—	—	
"	Low forceps	—	—	3515	L.	L.	—	—	
"	"	700	1	2466	L.	L.	—	—	Manual removal of placenta.

PROLONGED LABOUR (Continued)

Reg. No.	Age	Maturity	Height (inches)	Previous Pregnancies		Position of foetus		Other obstetric abnormalities	Cause of delay as diagnosed	Time of rupture of mem- branes i.e. hours before delivery	Duration of labour (hrs.)		
				Before 28 wks.	After 28 wks.	At onset of labour	At delivery				1st stage	2nd stage	3rd stage
5122	28	40	58 $\frac{3}{4}$	—	—	R.O.T.	R.O.A.	—	Uterine inertia	10:00	26:00	0:20	0:10
5437	21	36	54	—	—	L.O.A.	L.O.A.	Contracted pelvis	„	24:00	26:00	0:25	1:05
5591	30	29	61	—	—	L.O.A.	L.O.A.	—	„	15:00	26:05	0:15	0:05
5622	21	41	59 $\frac{1}{4}$	—	—	L.O.T.	L.O.T.	—	„	24:00	31:48	—	—
5849	26	39	60	—	—	R.O.P.	R.O.A.	—	„	9:00	24:30	0:10	0:10
5916	26	40	58	—	—	L.O.P.	L.O.P.	—	„	22:00	33:15	0:10	0:05
6124	28	40	59 $\frac{1}{2}$	—	—	L.O.A.	L.O.A.	Unclassified toxaemia	„	28.45	28.45	—	—
MULTIPARA													
BOOKED													
752	34	40	59 $\frac{3}{4}$	—	8	L.O.A.	L.O.A.	—	Uterine inertia	6:45	18:40	0:05	0:05
851	26	43	59 $\frac{1}{4}$	1	1	R.O.T.	L.O.T.	—	C-P disproportion	4:55	18:30	—	—
1648	41	39	62 $\frac{1}{4}$	—	9	R.S.A.	R.S.A.	—	Breech	23:00	> 22:00	—	—
1818	30	44	60 $\frac{1}{4}$	—	3	L.O.A.	L.O.A.	—	Uterine inertia	36:00	36:00	—	—
2056	24	3	56 $\frac{1}{4}$	—	2	R.O.A.	R.O.A.	Contracted pelvis	„	20:50	19:50	0:15	0:05
2442	34	39	59	—	5	R.S.A.	L.S.A.	Contracted pelvis breech P.E.	„	22:00	22:00	—	—
NON-BOOKED													
3879	32	39	?	1	3	L.O.A.	L.O.A.	Severe P.F.	Uterine inertia	—	27:00	0:15	0:15
5152	26	40	?	—	2	L.S.A.	L.S.A.	Contracted pelvis.	„	30:00	30:00	—	—
MULTIPLE PREGNANCY													
BOOKED													
4130	35	34	62	1	—	L.O.P. L.O.P.	L.O.P. L.O.P.	Severe P.E. Twins	Uterine inertia	3 days	—	—	—

Treatment	Method of delivery	P.P.H.	Morbidity	Weight of child (grams)	Result		Child		Remarks
					M.	C.	Trauma	Resuscitation Yes/No	
I.V. drip	Mid forceps	—	—	3033	L.	L.	—	—	
I.V. drip & sedation	Low forceps	—	—	2098	L.	L.	—	—	Manual removal of placenta.
„	„	—	1	3062	L.	L.	—	—	
„	L.S.C.S.	—	—	3374	L.	L.	—	Yes	
„	Mid forceps	—	—	3856	L.	L.	—	—	
„	Low forceps	—	—	3175	L.	L.	—	—	
„	L.S.C.S.	—	—	2778	L.	L.	—	—	
—	Spont.	—	—	3510	L.	L.	—	—	P.P.S.
Sedation	L.S.C.S.	—	—	4320	L.	L.	—	—	
I.V. drip & sedation	„	—	—	3912	L.	L.	—	Yes	
„	„	—	—	3544	L.	L.	—	—	
„	Spont.	—	—	2466	L.	L.	—	—	
Sedation	L.S.C.S.	—	—	2662	L.	L.	—	—	Anaemia.
Sedation	Mid forceps	800	1	3342	L.	L.	—	Yes	
I.V. drip & sedation	L.S.C.S.	—	—	3517	L.	L.	—	„	
Sedation	L.S.C.S.	—	—	2211 2551	L.	L.	—	—	

TABLE 19

LABOUR FOLLOWING PREVIOUS CAESAREAN SECTION

Total No. of cases	55
Booked	54
Non-booked	1
Primiparae	—
Multiparae	55
Total No. of babies	55
Multiple pregnancies	—
Maternal death	—
S. B.	—
N. N. D.	2 (0.36%)
P. N. D.	2 (0.36%)

TYPE OF PREVIOUS C.S.:—

L. S. C. S.	51
Extraperitoneal C. S.	—
C. C. S.	1
Unknown	3

INDICATIONS FOR PREVIOUS C.S.:—

Uterine Inertia	6
Cervical Dystocia	1
Pre-eclampsia	5
Eclampsia	2
Eclampsia + Contracted Pelvis	1
Contracted Pelvis	5
Contracted Pelvis + Breech Presentation	2
Contracted Pelvis + Placenta Praevia	1
C. — P. Disproportion	2
Uterine Didelphys	1
Stab Wound of Uterus	1
Placenta Praevia	9
Accidental A. P. H.	1
A. P. H. of Extra-placental Cause	1
Breech Presentation + Big Baby	1
Breech Presentation in Primip + Uterine Inertia	2
Breech Presentation in Elderly Primip + Postmaturity	1
Face Presentation	2
Brow Presentation	1
Transverse Lie	3
Foetal Distress	3
Conjoined Twins	1
Postmaturity	2
Failed Forceps	1

METHOD OF PRESENT DELIVERY:—

Spontaneous	28
Low forceps	22
Mid forceps	3
Assisted breech	2

TABLE 20
INDUCTION OF LABOUR

Total No. of cases	169
Booked	156
Non-booked	13
Primiparæ	85
Multiparæ	84
Incidence	2.68%
Total No. of babies	174
Multiple pregnancies	5
S. B.	2 (1.15%)
N. N. D.	6 (3.49%)
P. N. D.	8 (4.6%)

INDICATIONS AND FOETAL LOSS:—

Indications	No. of cases			Medical			Surgical			Failed medical followed by surgical		
	S.B.	N.N.D.	P.N.D.	S.B.	N.N.D.	P.N.D.	S.B.	N.N.D.	P.N.D.	S.B.	N.N.D.	P.N.D.
Toxaemia	15	—	—	—	—	—	1	1	2	—	—	—
Post-maturity	28	—	—	—	—	—	—	1	1	—	—	—
A. P. H.	..	—	—	—	—	—	1	2	3	—	—	—
Leaking membranes	2	—	—	—	—	—	—	—	—	—	—	—
Large baby	—	—	—	—	—	—	—	—	—	—	—	—
Anencephaly	1	—	1	—	—	—	—	—	—	—	1	1
Diabetes	—	—	—	—	—	—	—	—	—	—	—	—
Ca. Breast	—	—	—	—	—	—	—	—	—	—	—	—
Hydramnios	—	—	—	—	—	—	—	—	—	—	—	—
Total	46	—	1	1	1	1	2	4	6	—	1	1

INDUCTION OF LABOUR (Continued)

MEDICAL INDUCTION (INCLUDING 46 CASES WHO SUBSEQUENTLY WERE FOLLOWED BY SURGICAL INDUCTION).

<i>Method of Induction</i>	<i>No. of Cases</i>	<i>No. Failed</i>
O. B. E.	82	53
Pitocin drip	4	3
O. B. E. + Pitocin drip	6	6
Total	92	62

SURGICAL INDUCTION (INCLUDING 46 CASES OF FAILED MEDICAL INDUCTION).

<i>Method of Induction</i>	<i>No. of Cases</i>	<i>No. Failed</i>
Stripping of membranes	58	11
A. R. M.	7	—
Stripping of membranes & A. R. M	56	7
A. R. M. + Pitocin drip	1	—
Stripping + Pitocin drip	1	—
Total	123	18

OF THE CASES OF SURGICAL INDUCTION FOLLOWING FAILED MEDICAL INDUCTION 14 FAILED.

TABLE 21

FORCEPS DELIVERIES

Total No. of cases	732
Booked	701
Non-booked	31
Primiparae	546
Multiparae	186
Total No. of babies delivered by forceps	743
Incidence	11.79%
Multiple pregnancies	22
Maternal death	—
S. B.	2 (0.27%)
N. N. D.	11 (1.48%)
P. N. D.	13 (1.75%)

TYPE OF FORCEPS:—

Type of Forceps	No. of cases	Primiparae			Multiparae			
		S.B.	N.N.D.	P.N.D.	No. of cases	S.B.	N.N.D.	P.N.D.
Low Forceps	488	2	3	5	140	—	7	7
Mid Forceps	40	—	—	—	39	—	—	—
Manual Rotation & Mid Forceps ..	23	—	1	1	13	—	—	—
Total	551	2	4	6	192	—	7	7

INDICATIONS:—

Prolonged 2nd Stage	265
Rigid Perineum	3
Uterine Inertia	3
Foetal Distress	214
Breech	84
A. P. H.	6
Prolapse of Cord	4
Prolonged 1st Stage	22
Cardiac Disease	24
Medical Complication	7
Toxaemia	45
Contracted Pelvis	25
Maternal Distress	11
Previous C. S.	25
Previous Myomectomy	1
Intra-uterine Death	1
Compound Presentation	1
Carcinoma Breast (IV)	1
Big Baby (4,763 gms.)	1
Total	<u>743</u>

TABLE 22

VERSION IN LABOUR

Total No. of cases	5
Booked	2
Non-booked	3
Primiparae	2
Multiparae	3
Incidence	0.08%
Total No. of babies	5
Multiple pregnancies	4
Maternal death	—
S. B.	—
N. N. D.	—
P. N. D.	—

TABLE 23

CAESAREAN SECTIONS

Total No. of cases	532
Booked	490
Non-booked	42
Primiparae	247
Multiparae	285
Incidence	8.45%
Total No. of babies	538*
Multiple pregnancies	7
Maternal death	—
S. B.	13 (2.42%)
N. N. D.	13 (2.48%)
P. N. D.	24 (4.46%)
Repeat C. S.	146
Type of C. S.:—	
L. S. C. S.	524
C. C. S.	2
Caesarean Hysterectomy	6

INDICATIONS AND FOETAL LOSS:—

<i>Indications</i>	<i>No. of cases</i>	<i>S.B.</i>	<i>N.N.D.</i>	<i>P.N.D.</i>
Contracted pelvis	121	1	—	1
C. P. with breech presentation	32	2	—	2
C. P. disproportion	21	—	—	—
Failed trial of forceps (and with contracted pelvis)	4	—	—	—
Previous C. S. with C. P.	105	—	2	2
Prolonged labour	28	1	1	2
Placenta praevia	36	1	3	3
Accidental haemorrhage	11	4	1	5
A. P. H. (? carcinoma of cervix)	1	—	—	—
Foetal distress	66	—	4	3
Previous C.S. (only mild C.P.)	40	1	1	2
Abnormal presentations	15	1	—	1
Prolapse of cord	6	—	—	—
Uterine inertia	10	—	—	—
Failed surgical induction	3	—	—	—
Severe pre-eclampsia	4	—	—	—
Breech and big baby	2	—	—	—
Breech and postmaturity	1	—	—	—
Obstructed labour (hydrops)	1	1	—	1
Ovarian cyst complicating pregnancy	1	—	—	—

* One section was done for a second twin.

CAESAREAN SECTIONS (Continued)

<i>Indications</i>	<i>No. of cases</i>	<i>S.B.</i>	<i>N.N.D.</i>	<i>P.N.D.</i>
Thick uterine septum with breech	1	—	—	—
Stenosis of vagina	2	—	—	—
Intrauterine infection	1	1	—	1
3 repeated manual removal of placenta . .	1	—	—	—
Elderly primipara & precious baby with minor indications, P.E.T. etc.	14	—	—	—
Diabetes mellitus	1	—	—	—
Acute abdomen (Carcinoma of liver, appendicitis)	2	—	—	—
Carcinoma of cervix	1	—	—	—
Carcinoma of breast (IV)	1	—	1	1
Total	532	13	13	24

CAUSES OF FOETAL MORTALITY:—

Prematurity	4
Bronchopneumonia	2
Atelectasis of lungs	8
Congenital abnormalities	6
Hydrops foetalis	4
Unknown	2

MORBIDITY AFTER C.S.:—

'Post-operative reaction'	18
Abdominal wound infection	3
Genital infection	7
Urinary tract infection	22
Breast engorgement	6
Chest infection	4
Typhoid fever	1
Bacillary dysentery	1
Unknown	3
Total	<u>65</u>

RUPTURE OF UTERUS (Continued)

Reg. No.	Age	Previous Pregnancies		Time of rupture	Type of rupture	Site of rupture	Cause of rupture	Type of labour	Duration of labour (hrs.)			Weight of child (grams)	Result		Remarks
		Before	After						28 wks.	28 wks.	1st stage		2nd stage	3rd stage	
MULTIPARA															
BOOKED															
953	35	—	5	38	Incomplete dehiscence of scar.	Dehiscence of previous scar (L.S.C.S.)	Previous scar	—	—	—	—	2835	L.	L.	Contracted pelvis. P.P.S.
5724	30	—	2	39	Complete	At previous inverted-T incision	2 previous L.S.C.S. Contracted pelvis	Admitted with strong pains 3 hrs. after onset of labour	—	—	—	3090	L.	S.B.	L.S.C.S. & subtotal hysterectomy
373	31	—	1	39	Incomplete	Dehiscent scar (L.S.C.S.)	C.S. scar	—	—	—	—	3600	L.	L.	L.S.C.S. for primip breech at 1st pregnancy.
NON-BOOKED															
2168	36	1	4	37	Complete	Laceration in left side lower segment down to vagina left fornix	Transverse lie	Obstructed	34:05	—	—	1701	L.	S.B.	Total hysterectomy

TABLE 25

CONTRACTED PELVIS

(O. C. less than 10 cm., T. D. less than 12 cm., A. P.O. less than 11 cm.)

Total No. of cases	568
Booked	549
Non-booked	19
Primiparae	290
Multiparae	278
Incidence	9.02%
Total No. of babies	574
Multiple pregnancies	6
Maternal death	—
S. B.	8 (1.39%)
N. N. D.	6 (1.06%)
P. N. D.	14 (2.44%)

TYPE OF PELVIS:—

Gynaecoid	367
Gynaecoid-Anthropoid	17
Gynaecoid-Platypelloid	4
Gynaecoid-Android	3
Platypelloid	118
Anthropoid	53
Android	2
Naegeles' pelvis	2
Deformed	2

METHOD OF DELIVERY AND FOETAL LOSS:—

<i>Method of delivery</i>	<i>No. of babies</i>	<i>S.B.</i>	<i>N.N.D.</i>	<i>P.N.D.</i>
Spontaneous	168	2	2	4
Low forceps	68	—	—	—
Mid forceps	23	—	1	1
Assisted breech	17	1	1	2
Breech extraction	1	1	—	1
C. S.	295*	3	2	5
Caesarean Hysterectomy	1	—	—	—
Laparotomy	1	1	—	1
Total	574	8	6	14

* 4 C. S. for twin pregnancies.

CONTRACTED PELVIS (Continued)

Reg. No.	Age	Height (inches)	Maturity	Type of pelvis	Radiological Results Pelvic Measurements					Position	Surgical induction
					O.C.	I.D.	I.S.	A.P.	Outlet T.O.		
PRIMIPARA											
BOOKED											
7822/59	38	59	39	Anthropoid	10.9	10.5	—	11.0	—	R.S.A.	—
8013/59	18	60	42	Gynaecoid	9.0	11.9	—	11.1	—	L.O.A.	—
8023/59	24	57	43	"	9.6	12.6	—	10.0	—	L.O.A.	—
8065/59	21	59½	42	Platypelloid	9.5	12.6	—	9.8	—	L.O.A.	—
8107/59	21	58	40	Gynaecoid	10.4	12.4	9.3	9.3	—	R.O.A.	—
8130/59	40	60	38	Gynaecoid-platypelloid	10.9	12.4	—	9.4	—	R.O.P.	—
14/60	23	59¼	39	Gynaecoid	11.0	11.3	—	11.8	—	L.O.A.	—
25	30	56½	39	"	10.8	11.2	—	10.0	—	R.O.P.	—
61	18	58	43	"	10.4	11.6	—	10.9	—	R.O.A.	—
62	30	60½	43	"	11.3	11.4	—	10.2	—	L.O.A.	—
96	29	57	40	"	9.5	12.6	—	11.0	—	R.O.A.	—
110	29	57¾	40	Platypelloid	9.9	12.1	—	9.5	—	R.O.T.	—
141	31	60	40	Gynaecoid	9.7	12.2	—	10.7	—	R.O.P.	—
143	24	57¼	45	"	9.8	12.0	9.9	11.4	—	L.O.A.	—
156	28	59¼	37	"	10.2	11.3	8.5	10.3	—	L.S.A.	—
168	27	57½	40	"	10.1	11.8	—	11.4	—	R.O.A.	—
198	25	57	41	"	10.4	11.4	—	11.5	—	L.O.A.	—
211	23	59½	43	Gynaecoid-Anthropoid	10.6	11.5	—	11.3	—	L.O.A.	—
308	33	58½	40	Gynaecoid	11.0	11.8	—	9.9	—	R.S.A.	—
330	24	59	41	Anthropoid	11.6	11.2	9.25	10.9	—	L.O.A.	—
338	20	57¼	38	Gynaecoid	9.9	11.4	—	10.5	—	L.O.A.	—
342	33	62¼	40	"	9.9	—	—	10.0	—	L.O.A.	—
343	25	60½	40	"	11.4	11.4	—	9.25	—	L.O.P.	—
353	23	60	42	"	11.3	11.9	—	12.2	—	R.O.A.	—
360	24	58	40	"	9.2	11.2	—	11.2	—	R.O.A.	—
367	30	59¼	39	"	9.65	11.7	—	10.4	—	L.O.T.	—
375	20	56	42	"	9.9	12.5	10.1	10.4	—	L.O.A.	—
389	22	59	42	"	10.6	11.8	—	11.7	—	R.O.P.	—
405	25	59	40	"	9.95	11.9	8.75	10.3	—	R.O.P.	—
443	23	59¾	41	Platypelloid	9.4	12.95	—	9.6	—	L.O.A.	—
488	30	61	38	"	9.6	12.5	—	10.6	—	L.O.A.	—
513	33	59	40	"	9.9	12.8	8.8	10.4	—	L.O.A.	—
518	28	58	731	Gynaecoid	9.6	12.25	—	10.9	—	L.O.A.	—
545	20	56	41	"	9.6	10.75	—	10.0	—	L.O.A.	—
555	36	58	40	Platypelloid	9.1	13.00	—	11.4	—	L.O.A.	—
564	29	58¼	36	Gynaecoid	9.4	11.85	9.6	11.0	—	L.O.A.	—
582	29	59	38	"	10.2	11.35	8.8	10.15	—	L.O.P.	—
615	28	59¾	40	"	10.5	11.7	—	11.0	—	L.O.A.	—
622	30	56¼	41	Gynaecoid-Android	9.0	11.85	—	11.4	—	L.O.A.	—
663	28	61¼	41	Gynaecoid-Anthropoid	9.8	11.85	—	10.8	—	R.O.A.	—
664	26	62¼	37	Anthropoid	12.9	11.95	—	11.6	—	L.S.A.	—
689	31	61¾	41	Deformed pelvis	8.35	13.2	—	11.2	—	R.O.A.	—
731	23	59	41	Gynaecoid	10.75	11.9	—	9.85	—	L.O.A.	—
780	34	57¾	41	"	10.3	11.55	—	10.35	—	L.O.P.	—
846	30	57¾	41	"	10.95	11.5	—	11.1	—	L.O.A.	—
853	24	57½	44	"	11.0	11.4	—	11.75	—	L.O.A.	—
860	22	61½	?	"	9.9	12.1	9.5	11.7	—	R.O.A.	—
876	34	61¼	42	"	10.0	11.9	—	11.5	—	R.O.A.	—
887	27	57½	42	Platypelloid	8.1	12.3	—	10.2	—	L.O.T.	—
895	27	58½	39	Gynaecoid	10.8	10.95	—	11.2	—	L.O.A.	—
896	21	58½	43	"	11.4	11.7	9.7	10.5	—	L.O.A.	—
907	27	57½	39	"	9.85	12.7	—	10.2	—	R.O.T.	—
931	29	58¼	42	"	9.7	12.0	—	10.45	—	L.S.A.	—
939	22	56¼	40	"	9.0	11.1	—	9.85	—	L.O.P.	—
950	31	61¼	39	Platypelloid	9.9	13.5	—	11.9	—	L.O.A.	—
984	30	62½	38	Gynaecoid-Anthropoid	12.1	11.95	8.8	10.4	—	R.O.P.	—
993	28	54	39	Gynaecoid	9.8	11.8	—	10.5	—	L.O.T.	—
1012	30	56¾	39	Android-Gynaecoid	9.6	10.8	—	9.85	—	R.O.P.	—
1018	24	59½	40	Gynaecoid	11.3	11.9	—	10.0	—	R.O.A.	—
1031	25	62	41	Platypelloid	9.5	13.4	—	11.1	—	R.O.A.	—
1033	28	57¾	38	"	9.0	12.8	—	11.6	—	L.O.P.	—
1079	24	56½	36	Gynaecoid	10.1	11.7	—	11.5	—	L.O.T.	—
1124	27	59½	39	Anthropoid	12.3	11.5	10.0	11.5	—	L.O.P.	—
1199	32	57¼	42	Gynaecoid	10.5	11.6	—	11.2	—	R.O.T.	—
1232	23	60½	40	Anthropoid	12.0	11.65	10.0	11.5	—	R.O.A.	—
1235	28	57	41	Gynaecoid-Anthropoid	11.2	11.7	9.25	10.6	—	L.O.A.	—
1249	26	55¼	40	Gynaecoid	11.3	11.75	—	11.0	—	L.O.A.	—
1259	25	58	41	"	10.2	11.8	—	10.8	—	L.O.A.	—
1261	28	56½	39	Platypelloid	9.7	13.4	—	12.0	—	R.O.A.	—
1338	27	58½	41	Gynaecoid	10.7	11.0	9.4	10.5	—	R.O.A.	—

Trial of labour (hours)	Management		Length of labour (hrs.)		Weight of child (grams)	Result		Morbidity (days)	Previous C.S.	Remarks
	Method	of delivery	1st stage	2nd stage		M	C			
—	Elective	L.S.C.S.	—	—	3572	L.	L.	—	—	
9:50	Spont.		9:20	0:30	2420	L.	L.	—	—	
—	"		3:00	0:15	2870	L.	L.	—	—	
9:15	"		9:10	0:05	2440	L.	L.	—	—	
—	"		1:35	0:40	2870	L.	L.	—	—	
—	Elective	L.S.C.S.	—	—	3232	L.	L.	—	—	
24:55	Mid forceps		30:00	1:55	2760	L.	L.	—	—	
—	L.S.C.S.		24:00	—	2665	L.	L.	—	—	
—	Spont.		6:20	0:55	3000	L.	L.	—	—	
—	"		5:03	0:45	3450	L.	L.	—	—	Anaemia.
—	L.S.C.S.		—	—	3232	L.	L.	—	—	
10:00	"		—	—	3203	L.	L.	—	—	
—	"		8:00	—	2466	L.	L.	2	—	
—	Spont.		7:40	0:05	2880	L.	L.	—	—	
—	Assisted breech		3:30	0:10	2360	L.	L.	—	—	Accidental Syphilis. haemorrhage.
2:05	Low forceps		4:55	0:20	3000	L.	L.	—	—	
7:55	"		7:30	0:25	3300	L.	L.	—	—	
—	Spont.		12:00	0:40	2750	L.	L.	—	—	
—	Elective	L.S.C.S.	—	—	2835	L.	L.	—	—	Anaemia.
4:40	Spont.		4:10	0:30	3300	L.	L.	—	—	P.E.
—	"		6:55	0:15	2410	L.	L.	—	—	Chronic asthma. P.E.
14:00	"		13:15	0:45	2600	L.	L.	—	—	Anaemia.
9:00	L.S.C.S.		9:00	—	3459	L.	L.	—	—	"
6:50	Spont.		6:05	0:45	3530	L.	L.	—	—	
7:20	"		7:10	0:10	2700	L.	L.	—	—	
6:58	L.S.C.S.		—	—	3400	L.	L.	—	—	Bronchial asthma.
6:00	Spont.		5:35	1:10	3250	L.	L.	—	—	Anaemia.
4:30	Low forceps		4:30	0:15	3060	L.	L.	—	—	
6:20	L.S.C.S.		6:20	—	2892	L.	L.	—	—	
20:00	"		20:00	—	3289	L.	L.	—	—	
4:20	Low forceps		3:25	0:55	3080	L.	L.	—	—	
—	"		8:30	0:05	3250	L.	L.	—	—	Anaemia.
10:35	Spont.		10:55	0:20	3230	L.	L.	—	—	
16:50	"		16:30	0:20	2720	L.	L.	—	—	P.E.
5:00	L.S.C.S.		5:00	—	3515	L.	L.	—	—	Anaemia.
12:00	"		12:00	—	3033	L.	L.	—	—	
17:30	L.S.C.S.		17:30	—	2863	L.	L.	—	—	P.E.
—	Spont.		9:25	0:20	3260	L.	L.	—	—	
—	Elective	L.S.C.S.	—	—	4082	L.	L.	—	—	
10:55	Low forceps		10:50	0:05	3270	L.	L.	—	—	Anaemia.
8:05	Assisted breech		7:50	0:15	2830	L.	L.	—	—	
—	L.S.C.S.		—	—	2920	L.	L.	—	—	
—	Spont.		10:30	0:25	3005	L.	L.	—	—	
—	L.S.C.S.		—	—	3750	L.	L.	—	—	Ovarian cyst — chance finding.
9:35	Manual rotation and mid forceps		9:25	0:10	3600	L.	L.	—	—	
—	Spont.		7:45	0:20	3230	L.	L.	—	—	
—	"		9:15	0:40	3515	L.	L.	—	—	
15:00	"		15:00	0:20	2381	L.	L.	—	—	
—	L.S.C.S.		—	—	2930	L.	L.	—	—	
9:40	Spont.		9:25	0:10	2700	L.	L.	—	—	
—	"		7:00	0:50	2563	L.	L.	—	—	
5:30	Low forceps		6:30	0:40	2700	L.	L.	—	—	
—	L.S.C.S.		—	—	3118	L.	L.	—	—	
6:00	"		—	—	2920	L.	L.	—	—	
4:00	Low forceps		4:05	0:05	3118	L.	L.	—	—	
—	L.S.C.S.		26:00	1:00	3800	L.	L.	—	—	Failed forceps.
10:05	Spont.		10:05	0:10	2722	L.	L.	—	—	P.E.
13:00	L.S.C.S.		13:00	—	2865	L.	L.	—	—	Anaemia.
5:55	Spont.		5:55	1:00	3232	L.	L.	—	—	
—	"		19:00	0:20	3742	L.	L.	—	—	
5:45	"		5:45	0:05	2608	L.	L.	—	—	Anaemia. Pulmonary T.B.
24:00	L.S.C.S.		24:00	—	3232	L.	L.	—	—	
24:00	"		24:00	—	2580	L.	L.	—	—	Pulmonary T.B.
24:00	"		24:00	—	3033	L.	S.B.	—	—	Intra-uterine death.
—	Spont.		11:30	0:05	2863	L.	L.	—	—	
6:10	Low forceps		5:30	1:30	2977	L.	L.	—	—	Anaemia.
—	"		17:30	0:20	2693	L.	L.	—	—	
3:15	Spont.		3:50	0:35	3252	L.	L.	—	—	
—	"		3:00	0:50	3175	L.	L.	—	—	
12:30	"		11:45	0:45	2637	L.	L.	—	—	

CONTRACTED PELVIS (Continued)

Reg. No.	Age	Height (inches)	Maturity	Type of pelvis	Radiological Results				Outlet	Position	Surgical induction	
					Pelvic Measurements							
					O.C.	T.D.	I.S.	A.P.	T.O.			
1341	26	57 $\frac{1}{4}$	40	Gynaecoid	11.3	11.3	9.7	9.5	—	—	R.O.P.	—
1350	32	59 $\frac{1}{4}$	41	Platypelloid	8.7	13.0	—	—	—	—	I.O.T.	—
1366	22	59	42	Gynaecoid	10.2	11.8	—	10.8	—	—	L.O.A.	—
1395	27	59	43	Platypelloid	9.6	12.8	10.1	12.0	—	—	R.O.A.	—
1404	24	57 $\frac{3}{4}$?	Gynaecoid	11.7	11.9	9.6	10.5	—	—	L.S.A.	—
1435	27	52	39	"	11.3	11.6	—	10.8	—	—	L.O.A.	—
1441	26	58	39	"	9.5	12.2	—	11.5	—	—	I.O.P.	—
1472	26	59 $\frac{1}{4}$	40	"	11.5	11.8	9.7	11.8	—	—	L.O.A.	—
1475	21	60	?	Android	10.5	11.2	—	10.5	—	—	L.O.A.	—
1537	21	56 $\frac{1}{2}$?	Gynaecoid	10.6	11.9	—	11.5	—	—	L.O.A.	—
1541	33	61 $\frac{1}{2}$	40	"	10.3	11.7	8.5	10.3	—	—	R.O.A.	—
1548	25	60 $\frac{1}{2}$	38	Anthropoid — Gynaecoid	11.2	11.4	8.6	12.0	—	—	L.O.A.	—
1556	34	62 $\frac{1}{4}$	41	Gynaecoid	11.7	11.9	9.9	11.0	—	—	L.O.A.	—
1561	21	58	41	Platypelloid	9.8	13.0	10.6	10.5	—	—	R.O.A.	—
1564	30	59	39	Anthropoid	11.3	11.0	8.5	11.1	—	—	L.O.A.	—
1577	20	58 $\frac{1}{2}$	40	Gynaecoid	11.5	11.6	8.5	10.7	—	—	L.O.A.	—
1578	37	60	38	Anthropoid	13.3	11.2	8.0	12.2	—	—	L.O.A.	—
1673	21	58 $\frac{1}{4}$	40	Gynaecoid	11.0	11.7	9.3	9.8	—	—	R.O.A.	—
1675	24	56	?	"	9.7	12.0	9.0	9.4	—	—	R.O.T.	—
1686	23	57	39	Platypelloid	9.3	12.5	9.6	9.7	—	—	R.O.P.	—
1694	18	51 $\frac{3}{4}$	39	Anthropoid	10.9	10.8	9.2	9.9	—	—	R.O.T.	—
1707	23	59 $\frac{3}{4}$?	Gynaecoid	10.3	11.7	8.8	10.4	—	—	L.O.T.	—
1709	31	59	39	"	9.9	11.9	9.1	12.7	—	—	L.O.A.	—
1740	24	58 $\frac{1}{2}$?	"	9.5	11.0	9.7	10.2	—	—	L.O.A.	—
1772	19	58	?	Platypelloid	9.0	12.8	—	10.4	—	—	R.O.A.	—
1856	28	61	39	Gynaecoid	9.8	12.0	8.7	9.8	—	—	L.O.A.	—
1879	20	58 $\frac{1}{2}$	40	"	10.5	11.9	9.9	10.4	—	—	I.O.T.	—
1920	26	56	42	"	10.8	11.3	9.2	11.0	—	—	L.O.A.	—
1926	42	59	39	"	10.0	11.7	9.4	10.7	—	—	L.O.A.	—
1967	19	63 $\frac{1}{4}$	42	"	11.2	11.8	9.6	12.5	—	—	L.O.A.	—
1992	26	57 $\frac{1}{2}$	39	"	11.2	11.2	9.6	10.1	—	—	L.O.A.	—
2011	35	59	41	Platypelloid	8.2	11.2	9.6	9.3	—	—	L.O.A.	—
2046	22	59 $\frac{1}{2}$	41	Gynaecoid	10.7	11.4	—	10.5	—	—	L.O.A.	—
2066	29	60 $\frac{1}{4}$	42	"	9.8	11.5	9.4	10.8	—	—	R.O.A.	—
2070	20	60	44	Platypelloid	9.5	12.1	11.4	—	—	—	L.O.P.	—
2079	39	58	36	Gynaecoid	10.5	11.7	9.0	9.3	—	—	R.O.A.	—
2083	35	59 $\frac{3}{4}$	40	"	11.9	11.7	9.9	10.3	—	—	L.O.A.	—
2084	22	60 $\frac{3}{4}$	40	"	11.6	11.6	9.5	10.8	—	—	L.O.A.	—
2089	22	58 $\frac{1}{2}$	40	"	10.4	11.5	—	10.4	—	—	L.O.A.	—
2093	26	60 $\frac{1}{2}$	40	"	10.5	11.5	9.6	10.7	—	—	R.O.A.	—
2222	25	56 $\frac{1}{4}$	40	"	10.7	11.4	8.1	11.0	—	—	L.O.A.	—
2240	27	57	40	Platypelloid	8.6	12.6	—	12.0	—	—	L.O.A.	—
2258	27	58	40	Gynaecoid	10.2	11.1	—	11.1	—	—	R.O.A.	—
2262	22	59 $\frac{1}{2}$	40	Platypelloid	9.7	12.7	—	12.3	—	—	L.O.A.	—
2264	32	54	41	"	8.7	12.0	9.3	10.5	—	—	L.O.A.	—
2288	30	59 $\frac{3}{4}$	39	Gynaecoid	10.6	11.4	—	10.7	—	—	L.O.T.	—
2300	26	59 $\frac{1}{2}$?	"	9.8	12.0	—	9.5	—	—	L.O.A.	—
2309	22	58	40	"	9.5	11.5	9.2	9.9	—	—	L.O.A.	—
2348	41	59	40	"	10.8	11.9	9.8	10.0	—	—	R.O.T.	—
2349	25	58	41	"	10.3	11.8	9.5	11.0	—	—	R.O.A.	—
2354	23	57 $\frac{1}{2}$?	"	11.1	11.7	9.1	10.8	—	—	L.O.A.	—
2408	37	59 $\frac{1}{2}$	36	"	9.9	11.7	8.9	10.7	—	—	L.O.T.	—
2436	21	58	36	"	8.7	11.6	8.5	11.5	—	—	L.O.A.	—
2467	24	59	40	"	9.5	12.1	9.8	9.5	—	—	R.O.A.	—
2519	20	58	40	"	11.0	11.8	9.1	10.8	—	—	L.O.A.	—
2529	33	51 $\frac{1}{2}$	40	"	9.7	11.8	9.6	10.6	—	—	L.O.A.	—
2567	29	59	38	"	9.7	11.8	9.4	10.5	—	—	L.S.P.	—
2582	35	59 $\frac{1}{2}$	41	Platypelloid	9.3	12.6	9.5	10.0	—	—	L.O.A.	—
2584	31	61	41	Gynaecoid	9.5	11.8	8.4	10.7	—	—	L.O.A.	—
2611	24	63	41	Anthropoid	12.4	11.4	—	12.5	—	—	R.O.P.	—
2627	29	61 $\frac{1}{4}$	41	Platypelloid	9.7	13.3	9.6	11.7	—	—	L.O.T.O.A.	—
2660	21	56 $\frac{1}{4}$	40	Gynaecoid	9.7	11.7	—	9.5	—	—	L.O.A.	—
2677	21	60	40	"	11.4	11.7	9.0	9.7	—	—	L.O.A.	—
2713	29	60 $\frac{1}{2}$	40	"	11.1	11.9	—	11.5	—	—	R.O.A.	—
2714	20	58	34	Anthropoid	11.9	11.5	10.4	10.7	—	—	R.O.A.	—
2789	33	63 $\frac{1}{4}$	39	Gynaecoid	11.2	11.4	9.4	11.9	—	—	L.O.A.	—
2812	36	59 $\frac{1}{4}$	39	"	12.2	11.9	—	11.3	—	—	R.S.P.	—
2823	30	59	36	Anthropoid	12.3	10.6	7.0	11.3	—	—	R.S.A.	—
2843	20	59 $\frac{1}{2}$	39	Gynaecoid	9.9	11.9	9.9	10.6	—	—	R.O.A.	—
2852	38	59 $\frac{1}{2}$	41	"	10.8	11.7	8.7	9.5	—	—	L.O.A.	—
2854	30	58	39	"	11.2	11.4	8.2	11.4	—	—	R.S.A.	—
2873	27	59	40	"	10.0	11.4	8.9	11.2	—	—	L.O.A.	—
2875	26	59	?	Platypelloid	9.5	12.4	10.4	—	—	—	R.O.P.	—
2931	27	62	38	"	9.4	12.4	—	11.0	—	—	L.O.A.	—
2981	15	58	38	Gynaecoid	10.6	11.9	9.9	11.9	—	—	L.O.A.	—
3022	25	60 $\frac{1}{4}$	39	Platypelloid	9.4	12.4	9.0	10.8	—	—	R.O.A.	—
3047	22	61	41	"	9.9	13.1	11.0	11.0	—	—	L.O.A.	—
3084	26	56 $\frac{3}{4}$	37	Gynaecoid- Anthropoid	10.3	11.35	10.15	11.2	—	—	L.O.A.	—

Trial of labour (hours)	Management		Length of labour (hrs.)		Weight of child (grams)	Result		Morbidity (days)	Previous C.S.	Remarks
	Method of delivery		1st stage	2nd stage		M	C			
7:00	L.S.C.S.		7:00	—	2664	L.	L.	—	—	
8:00	"		8:00	—	3260	L.	L.	—	—	
—	"		5:20	—	3629	L.	L.	—	—	
—	"		—	—	3570	L.	L.	—	—	
—	Assisted breech		6:40	0:10	2835	L.	L.	—	—	
17:20	Low forceps		17:20	1:30	3005	L.	L.	—	—	
8:00	L.S.C.S.		8:00	—	2750	L.	L.	—	—	
—	Spont.		11:30	0:15	2041	L.	L.	—	—	Mitral stenosis.
—	"		1:35	0:35	3402	L.	L.	—	—	P.F.
—	"		10:45	0:10	2325	L.	L.	1	—	
10:00	L.S.C.S.		10:00	—	2580	L.	L.	—	—	Anaemia.
10:35	Low forceps		10:35	0:05	2920	L.	L.	—	—	
12:00	L.S.C.S.		12:00	—	2864	L.	L.	—	—	
—	Low forceps		10:10	0:15	3232	L.	L.	—	—	
—	Spont.		14:45	0:20	2835	L.	L.	—	—	
9:00	"		9:20	0:55	2863	L.	L.	—	—	P.E.
—	Low forceps		6:30	0:10	3500	L.	L.	—	—	
20:00	L.S.C.S.		20:00	—	3370	L.	L.	—	—	Anaemia.
6:00	"		6:00	—	2920	L.	L.	—	—	"
2:30	"		—	—	2845	L.	L.	—	—	
13:35	Low forceps		13:35	0:30	2495	L.	L.	—	—	
18:00	L.S.C.S.		—	—	2863	L.	L.	—	—	
8:30	Spont.		8:10	0:20	3284	L.	L.	—	—	Anaemia.
—	Flective L.S.C.S.		—	—	3005	L.	L.	—	—	
—	"		—	—	3345	L.	L.	—	—	
9:00	L.S.C.S.		—	—	2863	L.	L.	—	—	
10:00	Spont.		10:20	1:20	2665	L.	L.	—	—	
—	Low forceps		11:05	0:15	2268	L.	L.	—	—	
—	L.S.C.S.		—	—	2863	L.	L.	—	—	
—	Spont.		22:50	0:40	3544	L.	L.	—	—	
18:30	Mid forceps		18:35	1:45	2948	L.	L.	—	—	
—	Flective L.S.C.S.		—	—	2948	L.	L.	—	—	
4:00	L.S.C.S.		4:00	—	3175	L.	L.	—	—	Moniliasis.
8:20	Mid forceps		8:20	0:15	3289	L.	L.	—	—	
—	L.S.C.S.		—	—	2948	L.	L.	—	—	
—	"		—	—	2495	L.	L.	—	—	Placenta praevia type II.
28:00	Mid forceps		28:00	0:15	2948	L.	L.	—	—	
—	Spont.		2:00	0:45	3260	L.	L.	—	—	Syphilis
20:30	Low forceps		20:30	0:10	2863	L.	L.	—	—	
10:40	"		10:40	0:35	2835	L.	L.	—	—	
18:00	L.S.C.S.		18:00	—	2948	L.	L.	—	—	
6:55	Spont.		15:55	0:10	2296	L.	L.	—	—	
4:00	L.S.C.S.		4:00	—	3062	L.	L.	—	—	
20:00	"		20:00	—	2551	L.	L.	—	—	
—	"		—	—	2892	L.	L.	—	—	
27:00	"		27:00	—	3062	L.	L.	—	—	
14:10	Spont.		14:10	0:10	2722	L.	L.	—	—	
11:00	L.S.C.S.		11:00	—	2835	L.	L.	—	—	
11:00	"		11:00	—	2835	L.	L.	—	—	
8:40	Spont.		8:40	0:20	2722	L.	L.	—	—	
5:25	"		5:25	0:35	3260	L.	L.	—	—	
7:30	L.S.C.S.		7:30	—	2892	L.	L.	—	—	Hookworm, anaemia.
18:00	"		18:00	—	2977	L.	L.	—	—	Anaemia.
—	Spont.		5:20	1:00	2948	L.	L.	—	—	
4:00	L.S.C.S.		4:00	—	2665	L.	L.	—	—	
—	"		—	—	2750	L.	L.	—	—	
—	"		—	—	2523	L.	L.	—	—	
10:35	Low forceps		10:35	1:00	3118	L.	L.	—	—	
—	L.S.C.S.		—	—	2948	L.	L.	—	—	Mitral stenosis.
11:00	"		11:00	—	3203	L.	L.	—	—	
14:25	Manual rotation and mid forceps		13:20	1:05	3289	L.	L.	—	—	
14:20	Low forceps		14:20	0:25	2268	L.	L.	—	—	
2:15	"		2:15	0:35	2551	L.	L.	—	—	
15:00	Spont.		14:45	0:35	2885	L.	L.	—	—	
—	"		6:55	0:30	2792	L.	L.	—	—	P.E. Anaemia.
8:00	Assisted breech		8:00	0:30	2705	L.	L.	—	—	
—	L.S.C.S.		—	—	2778	L.	L.	—	—	
13:00	Low forceps		12:40	0:15	2772	L.	L.	—	—	
10:00	"		9:55	0:35	2948	L.	L.	—	—	P.E.
—	Elective L.S.C.S.		1:30	0:15	3062	L.	L.	—	—	
—	L.S.C.S.		—	—	3147	L.	L.	—	—	
—	Spont.		8:10	0:50	2803	L.	L.	—	—	
15:10	L.S.C.S.		—	—	2300	L.	N.N.D.	—	—	Patent ductus. Baby died of disease of lung & bronchopneumonia.
8:00	Spont.		7:35	0:25	2778	L.	L.	—	—	Anaemia.
—	"		9:30	0:40	3203	L.	L.	—	—	
12:00	Low forceps		11:10	0:55	2892	L.	L.	—	—	P.P.H. Syphilis.
—	Spont.		6:00	0:55	3175	L.	L.	—	—	Anaemia.
—	Low forceps		2:15	0:20	3062	L.	L.	2	—	Severe P.E. Anaemia.

CONTRACTED PELVIS (Continued)

Reg. No.	Age	Height (inches)	Maturity	Type of pelvis	Radiological Results Pelvic Measurements				Outlet T.O.	Position	Surgical induction
					O.C.	T.D.	I.S.	A.P.			
3103	24	57	39	Gynaecoid	9.7	11.1	—	10.6	—	L.O.A.	—
3110	23	60	39	"	11.6	11.7	9.6	11.3	—	R.O.A.	—
3156	31	54½	40	"	9.7	12.1	8.5	9.0	—	R.O.P.-O.A.	—
3198	23	59	44	"	10.0	11.6	—	11.5	—	R.O.P.	—
3217	22	57¾	41	"	10.7	11.4	9.9	10.5	—	R.O.P.	—
3228	27	61	41	"	11.0	11.6	9.6	11.6	—	L.O.A.	—
3243	27	60	39	"	10.0	11.9	9.9	11.1	—	R.O.A.	—
3254	26	57	36	"	9.9	11.3	9.0	10.7	—	L.O.A.	—
3279	26	60	40	Anthropoid	12.3	11.6	10.1	11.2	—	L.O.A.	—
3309	31	58¼	37	Platypelloid	9.6	12.7	—	10.7	—	R.O.P.	—
3317	22	57	40	Gynaecoid	10.5	11.1	9.0	9.7	—	R.O.A.	—
3324	22	57	35	"	11.0	11.1	8.8	10.7	—	L.S.A.	—
3342	25	55½	41	"	9.8	11.4	9.7	10.2	—	R.O.P.-O.A.	—
3377	23	60½	35	Platypelloid	9.3	13.3	10.6	10.4	—	R.S.T.	—
3413	27	58	41	Gynaecoid	10.4	11.9	10.1	11.2	—	L.O.P.	—
3429	34	60¼	38	Platypelloid	9.0	12.3	8.5	11.0	—	L.O.A.	—
3451	32	57¼	39	Gynaecoid	9.8	12.5	10.3	10.3	—	L.O.P.	—
3452	29	59	40	"	10.2	10.9	8.6	9.5	—	L.O.A.	—
3459	24	59½	43	"	10.7	11.4	10.3	12.2	—	R.O.A.	—
3462	25	60½	43	Anthropoid	11.5	11.3	—	10.7	—	L.O.A.	—
3526	28	57¾	37	Platypelloid	9.7	12.6	9.0	9.7	—	R.O.P.	—
3532	23	57½	35	Gynaecoid	10.4	11.0	8.5	10.2	—	R.S.A.	—
3595	24	60	?	"	10.2	11.9	9.3	10.2	—	L.O.A.	—
3651	26	60	42	Anthropoid	11.4	11.3	8.2	10.9	—	L.O.A.	—
3672	31	59½	39	Gynaecoid	10.4	11.6	9.9	10.9	—	R.O.P.	—
3675	31	56½	39	"	9.7	11.8	9.0	11.4	—	L.O.A.	—
3685	20	59½	40	Anthropoid	12.2	11.3	9.4	11.0	—	L.O.A.	—
3728	34	59	41	Platypelloid	9.4	12.4	9.3	8.7	—	R.O.T.	—
3729	30	59	41	Gynaecoid	9.7	12.5	9.6	11.1	—	L.O.T.	—
3733	28	60½	41	Platypelloid	9.6	13.0	10.6	11.4	—	L.O.A.	—
3768	32	58	41	Gynaecoid	9.8	12.0	9.9	11.0	—	R.O.A.	—
3779	32	58¾	39	"	10.9	11.7	8.2	11.0	—	L.O.A.	—
3787	24	61½	41	"	10.5	11.7	—	10.5	—	L.S.P.	—
3802	18	60	41	Anthropoid	11.8	11.2	9.1	11.8	—	R.O.A.	—
3811	22	57½	40	Gynaecoid	10.7	11.9	9.8	10.8	—	L.O.A.	—
3846	22	61½	37	Anthropoid	12.4	11.8	9.8	13.0	—	R.O.A.	—
3970	30	59	40	Gynaecoid	9.3	11.9	9.6	10.3	—	R.O.A.	—
3984	25	60¼	40	Anthropoid	12.0	11.1	8.9	11.0	—	R.O.T.	—
3991	28	64	41	Gynaecoid	11.2	11.9	10.0	10.2	—	R.O.T.-O.A.	—
4004	27	60½	40	"	10.9	11.4	9.8	10.9	—	L.O.A.	—
4018	22	57	36	"	10.4	11.8	—	11.0	—	L.O.A.	—
4054	33	57	40	"	10.9	11.4	8.8	11.1	—	R.O.T.-O.A.	—
4090	25	60	40	"	10.6	11.8	9.4	11.0	—	R.O.P.	—
4101	26	55¼	40	"	8.9	11.2	9.0	9.9	—	L.O.P.	—
4171	28	58½	42	Platypelloid	8.0	11.8	8.5	11.4	—	R.S.A.	—
4175	20	59	40	"	10.0	11.3	—	11.1	—	L.O.A.	—
4181	30	60½	38	"	9.2	12.6	9.6	10.0	—	R.O.P.-O.A.	—
4189	21	59¾	39	Gynaecoid	10.4	11.0	9.35	10.4	—	L.O.A.	—
4206	29	62¼	37	"	10.9	11.8	10.2	12.2	—	L.O.T.	—
4262	25	60½	39	Anthropoid	12.0	11.5	9.25	11.0	—	L.O.T.	—
4269	23	60	41	Anthropoid- Gynaecoid	11.2	11.2	9.2	11.7	—	L.O.A.	—
4272	26	57	40	Platypelloid	8.6	12.9	10.3	10.8	—	R.S.A.	—
4295	19	60	?	Gynaecoid	11.0	11.9	—	12.0	—	L.O.A.	—
4321	23	60½	41	"	10.0	11.7	8.2	10.0	—	L.O.A.	—
4324	29	58¼	41	Anthropoid- Gynaecoid	11.1	11.55	8.5	11.1	—	R.O.A.	—
4332	23	60	39	Gynaecoid	11.2	11.1	8.8	10.0	—	L.O.A.	—
4384	21	59¾	41	"	11.5	11.7	9.4	10.8	—	L.S.A.	—
4416	31	60	42	"	11.0	11.9	8.5	11.0	—	L.O.T.	—
4423	28	55	36	"	9.5	11.2	8.3	11.0	—	R.O.P.	—
4426	34	60½	40	"	11.8	11.7	9.4	11.5	—	R.O.P.	—
4428	27	62	38	Anthropoid	11.2	11.2	9.2	12.3	—	R.O.A.	—
4429	21	60½	38	Gynaecoid	10.6	11.2	9.0	11.2	—	R.S.A.	—
4463	23	56	38	Platypelloid	8.3	11.6	9.35	10.8	—	R.O.T.	—
4472	27	61½	39	Gynaecoid	11.0	11.5	9.2	10.6	—	R.O.P.	—
4491	24	63	39	Anthropoid	12.6	11.5	9.4	10.8	—	L.S.A.	—
4554	27	53	37	"	10.0	10.2	8.7	10.0	—	R.O.T.	—
4598	23	57¼	35	Gynaecoid	10.4	11.2	8.7	11.1	—	L.O.P.-O.A.	—
4603	24	61	43	Platypelloid	9.4	13.3	10.2	11.6	—	L.O.A.	—
4621	29	61	40	"	9.9	13.3	10.9	10.1	—	L.S.A.	—
4622	22	63	40	Anthropoid- Gynaecoid	11.1	11.45	9.3	10.7	—	L.O.A.	—
4625	40	57	38	Gynaecoid- Platypelloid	9.6	12.3	9.9	9.4	—	R.O.T.	—

Trial of labour (hours)	Management		Length of labour (hrs.)		Weight of child (grams)	Result		Morbidity (days)	Previous C.S.	Remarks
	Method of delivery		1st stage	2nd stage		M	C			
—	Elective	L.S.C.S.	—	—	2840	L.	L.	—	—	Thrototoxicosis.
5:45	Spont.		5:45	0:40	3374	L.	L.	—	—	
6:05	Mid forceps		6:05	0:45	2807	L.	N.N.D.	—	—	Baby died of congenital heart, atelectasis.
4:00	L.S.C.S.		4:00	—	4423	L.	L.	—	—	
7:00	"		7:00	—	3572	L.	L.	4	—	
12:40	"		—	—	2700	L.	L.	—	—	
17:40	Spont.		17:40	0:40	3005	L.	L.	—	—	
—	L.S.C.S.		—	—	2580	L.	L.	—	—	
—	Spont.		9:40	0:20	2863	L.	L.	—	—	
13:30	L.S.C.S.		13:30	—	3203	L.	L.	—	—	
6:00	Spont.		6:00	0:10	2296	L.	L.	—	—	
10:00	Assisted breech		10:00	0:15	2126	L.	S.B.	—	—	Macerated foetus.
6:00	Mid forceps		6:00	0:05	2920	L.	L.	—	—	
—	L.S.C.S.		12:00	—	2387	L.	S.B.	—	—	Baby — Postmortem showed polycystic kidney.
5:15	"		5:15	—	3402	L.	L.	—	—	
8:00	"		8:00	—	2495	L.	L.	—	—	Severe P.E. Ovarian cyst—chance finding.
4:00	"		4:00	—	2551	L.	L.	—	—	
16:15	Low forceps		16:15	0:15	2608	L.	L.	—	—	
—	Spont.		1:30	0:15	2722	L.	L.	—	—	
—	Low forceps		4:15	1:25	3203	L.	L.	—	—	
25:00	L.S.C.S.		25:00	—	2863	L.	L.	5	—	P.E. Anaemia.
9:00	"		—	—	2381	L.	L.	—	—	
7:30	Spont.		7:00	0:30	2647	L.	L.	—	—	
12:00	Low forceps		11:50	1:00	3374	L.	L.	—	—	Anaemia. Moniliasis.
3:30	Mid forceps		3:30	6:15	2637	L.	L.	—	—	
9:00	L.S.C.S.		—	—	2495	L.	L.	—	—	
—	Low forceps		6:45	6:10	2863	L.	L.	—	—	P.E.
7:45	L.S.C.S.		7:45	—	3175	L.	L.	—	—	"
8:45	Manual rotation and mid forceps		8:45	0:10	3232	L.	L.	—	—	
15:00	L.S.C.S.		15:00	—	3489	L.	L.	—	—	
—	Low forceps		17:55	0:10	2693	L.	L.	—	—	P.E. Anaemia.
—	"		10:45	0:10	2835	L.	L.	—	—	
4:00	Assisted breech		4:00	0:35	2735	L.	L.	—	—	Anaemia.
—	Low forceps		8:35	1:15	2863	L.	L.	—	—	
6:10	Spont.		6:10	0:35	3402	L.	L.	—	—	
3:50	"		3:50	0:10	2977	L.	L.	—	—	Anaemia.
7:00	"		7:00	0:25	3175	L.	L.	—	—	"
6:40	Low forceps		6:40	0:40	2551	L.	L.	—	—	
11:40	"		11:40	0:05	3232	L.	L.	—	—	Mitral stenosis.
—	Spont.		6:20	0:25	2665	L.	L.	—	—	Anaemia.
—	"		4:45	0:05	2381	L.	L.	—	—	"
5:00	Low forceps		12:30	1:00	3175	L.	L.	—	—	
7:20	L.S.C.S.		7:20	—	3317	L.	L.	—	—	Anaemia. Syphilis.
6:00	"		6:00	—	2835	L.	L.	1	—	
—	"		9:00	—	2977	L.	L.	—	—	P.E. Syphilis.
15:00	"		15:00	—	2693	L.	L.	2	—	
10:15	Mid forceps		9:45	0:30	2618	L.	L.	2	—	Anaemia.
13:45	Low forceps		13:30	0:05	3005	L.	L.	—	—	
14:00	L.S.C.S.		—	—	3884	L.	L.	—	—	
29:00	"		—	—	3232	L.	L.	—	—	
8:30	Spont.		8:30	0:35	3090	L.	L.	—	—	
—	L.S.C.S.		—	—	2523	L.	L.	—	—	
7:10	Spont.		7:10	0:10	2920	L.	L.	—	—	
8:45	Low forceps		7:15	1:25	2523	L.	L.	—	—	P.E. Anaemia.
5:30	Spont.		5:30	0:25	3118	L.	L.	—	—	
12:00	"		12:00	0:15	2353	L.	L.	—	—	
—	L.S.C.S.		6:00	—	2551	L.	L.	—	—	
8:30	"		8:30	—	3147	L.	L.	—	—	
8:00	"		8:00	2:50	2580	L.	L.	—	—	Failed forceps.
7:00	Low forceps		7:00	1:00	2807	L.	L.	—	—	
1:55	Spont.		1:50	0:05	2353	L.	L.	—	—	P.E.
—	Assisted breech		3:05	0:10	2693	L.	L.	—	—	
7:30	L.S.C.S.		7:30	—	2495	L.	S.B.	—	—	
20:00	"		20:00	—	3317	L.	L.	—	—	P.E.
—	Assisted breech		8:45	0:10	2608	L.	L.	—	—	
10:00	L.S.C.S.		10:00	—	2353	L.	L.	—	—	Anaemia.
7:00	Manual rotation and mid forceps		7:00	1:20	2722	L.	L.	—	—	
—	Low forceps		10:20	0:10	3629	L.	L.	—	—	
6:05	Assisted breech		5:40	0:15	3118	L.	L.	—	—	Anaemia.
4:20	Spont.		4:00	0:20	2523	L.	L.	—	—	"
11:00	L.S.C.S.		11:00	—	2608	L.	L.	—	—	

CONTRACTED PELVIS (Continued)

Reg. No.	Age	Height (inches)	Maturity	Type of pelvis	Radiological Results Pelvic Measurements					Position	Surgical induction
					O.C.	T.D.	I.S.	Outlet A.P.	T.O.		
4648	43	56½	35	Platypelloid	8.7	12.0	9.4	9.4	—	R.O.T.	—
4666	24	57	39	Gynaecoid	11.1	11.5	9.3	11.1	—	R.O.A.	—
4696	27	59	41	"	11.3	11.9	10.3	11.6	—	R.O.A.	—
4734	22	58	42	"	10.3	11.8	10.2	11.0	—	L.O.A.	—
4755	31	58½	40	"	10.8	11.6	9.0	11.7	—	L.O.A.	—
4783	23	62	40	"	11.2	11.8	8.5	11.1	—	L.O.A.	—
4785	16	57½	39	"	11.2	11.4	—	11.2	—	R.S.A.	—
4795	25	58	38	Platypelloid	9.4	13.1	11.1	11.4	—	L.O.A.	—
4804	28	55	41	Gynaecoid	10.8	11.3	8.9	10.2	—	R.O.P.	—
4817	29	59½	38	"	11.3	11.3	7.9	11.0	—	L.O.A.	—
4818	33	57½	42	"	9.9	12.4	9.2	10.7	—	R.O.A.	—
4844	25	59½	42	"	9.9	11.6	9.3	10.9	—	L.O.P.	—
4854	25	55¾	39	"	10.5	11.4	8.8	10.4	—	R.O.T.	—
4873	20	62	38	Platypelloid	9.7	12.8	—	11.2	—	L.S.A.	—
4885	22	59	44	Gynaecoid	10.0	11.8	9.9	10.0	—	L.O.A.	—
4890	28	57½	39	"	10.3	11.5	8.7	10.6	—	R.O.T.-O.A.	—
4908	22	61	40	"	11.1	11.1	—	9.5	—	L.O.A.	—
4935	25	61½	38	"	10.5	12.0	9.7	9.5	—	R.O.T.-O.A.	—
4963	26	59½	38	"	11.5	11.7	9.5	10.6	—	L.O.A.	—
4973	32	58¾	41	Anthropoid	11.1	11.8	9.6	11.1	—	R.O.A.	—
4982	28	56	44	Gynaecoid	10.6	11.65	10.2	11.2	—	L.O.A.	—
5027	24	59	40	"	10.2	11.9	8.4	10.6	—	L.O.P.	—
5029	34	58	40	Platypelloid	9.8	12.2	9.3	10.2	—	R.O.T.	—
5066	36	61	39	Anthropoid	12.0	11.9	11.1	11.0	—	R.S.A.	—
5067	24	58	42	Platypelloid	9.0	13.0	—	—	—	R.O.P.-O.P.	—
5096	22	59½	40	Gynaecoid	10.6	11.8	9.3	10.5	—	L.O.A.	—
5125	23	59	39	"	8.8	11.3	10.7	10.8	—	R.O.T.	—
5141	21	56	40	Anthropoid	12.3	11.6	8.8	11.1	—	L.O.A.	—
5179	20	55	39	Gynaecoid	10.3	11.3	8.5	10.6	—	L.S.A.	—
5195	29	59	44	Anthropoid	12.2	11.6	9.0	11.3	—	R.O.T.-O.A.	—
5200	31	NB	41	"	12.5	11.8	9.9	11.4	—	L.S.A.	—
5203	30	61¼	41	"	11.3	11.0	8.1	10.2	—	L.O.T.	—
5208	28	60½	41	Gynaecoid	9.9	11.8	9.6	10.8	—	L.O.T.	—
5257	33	60½	42	Platypelloid	9.5	12.5	10.2	9.7	—	R.O.P.	—
5278	26	62½	41	Anthropoid	11.7	11.6	10.0	10.7	—	L.O.A.	—
5301	19	57	37	Gynaecoid	9.4	11.9	—	10.3	—	L.O.A.	—
5326	23	53	39	Platypelloid	9.1	12.4	9.6	10.5	—	L.O.A.	—
5341	20	58	39	Gynaecoid	9.5	12.0	9.7	9.5	—	L.O.A.	—
5355	21	59	41	Platypelloid	9.7	12.8	10.6	11.7	—	L.O.A.	—
5365	28	57½	?	Gynaecoid	10.5	11.2	8.3	10.5	—	R.S.A.	—
5375	25	62	40	Anthropoid	12.5	11.6	9.5	10.7	—	L.O.A.	—
5414	24	59½	40	Gynaecoid	11.9	11.9	8.8	11.5	—	L.O.A.	—
5434	26	54½	36	"	9.5	11.8	9.2	10.8	—	L.G.A.	—
5437	21	54	36	Anthropoid	10.6	10.5	8.5	10.9	—	L.O.A.	—
5453	28	59½	40	Platypelloid	8.5	12.4	—	10.0	—	L.O.T.	—
5480	26	58½	41	Gynaecoid	9.8	12.0	10.0	10.9	—	L.S.A.	—
5507	27	55	37	Anthropoid	11.5	10.5	9.0	10.7	—	R.O.A.	—
5525	23	56½	38	Gynaecoid	11.4	11.5	—	10.8	—	R.O.A.	—
5580	25	57	38	"	9.8	10.9	7.9	10.7	—	L.O.A.	—
5607	18	62	42	"	11.7	11.9	9.9	10.0	—	R.O.A.	Stripping. A.R.M.
5666	33	57	40	"	10.8	11.0	9.2	9.6	—	L.O.T.	—
5757	32	58	40	"	10.1	11.5	9.7	9.1	—	R.O.A.	—
5877	25	57	40	"	10.3	11.2	—	10.3	—	L.O.T.	—
5904	27	60	40	"	11.0	11.8	—	11.0	—	R.O.A.	—
5910	22	58½	34	Platypelloid	9.7	12.6	9.6	11.1	—	L.O.A.	—
5935	20	61¾	41	Gynaecoid	11.4	11.6	9.8	10.5	—	L.O.A.	—
5962	21	61	40	"	10.6	11.4	10.0	8.9	—	L.O.T.-O.A.	—
5987	22	60	42	"	9.7	12.4	9.6	11.0	—	L.O.T.	—
6028	37	58½	42	Platypelloid	9.5	13.0	—	11.5	—	R.O.T.	—
6112	21	57½	38	Gynaecoid	10.6	11.0	9.5	10.0	—	R.O.T.-O.A.	—
6173	27	56½	40	"	10.9	11.2	8.4	10.0	—	R.O.A.	—
6192	23	56½	40	"	9.9	11.2	—	10.6	—	L.O.A.	—
6213	29	60	42	Platypelloid	9.7	12.7	—	12.2	—	L.O.T.	—
6232	30	61¾	40	Android	9.8	12.2	9.1	11.7	—	L.O.A.	—
6338	25	61	40	Gynaecoid	9.8	12.0	9.4	10.9	—	R.O.A.	—
NON-BOOKED											
8190/59	26	60½	44	Platypelloid	9.5	12.9	—	9.9	—	R.S.A.	—
654	27	58¼	40	Gynaecoid- Platypelloid	9.7	12.7	—	9.4	11.5	R.O.A.	—
1098	23	57	42	Platypelloid	9.2	11.2	—	10.8	—	R.O.A.	—
1473	23	52	?	Gynaecoid	11.0	11.8	8.4	10.8	—	L.O.A.	—

Trial of labour (hours)	Management		Length of labour (hrs.)		Weight of child (grams)	Result		Morbidity (days)	Previous C.S.	Remarks
	Method of delivery		1st stage	2nd stage		M	C			
—	L.S.C.S		6:00	—	2778	L.	L.	—	—	
8:00	Spont.		8:00	0:05	3481	L.	L.	—	—	
—	"		7:40	0:30	2495	L.	L.	—	—	
—	"		11:35	1:00	3941	L.	L.	—	—	
4:45	Low forceps		4:45	0:10	2665	L.	L.	—	—	Mitral stenosis.
—	"		10:45	0:25	3232	L.	L.	—	—	Anaemia.
—	"		—	—	3629	L.	L.	—	—	Eclampsia.
1:30	Spont.		?	?	2892	L.	L.	—	—	
8:00	L.S.C.S.		—	—	3912	L.	L.	—	—	
11:00	"		11:00	—	2183	L.	L.	—	—	
—	Spont.		10:55	0:05	2693	L.	L.	—	—	
7:00	L.S.C.S.		7:00	—	2637	L.	L.	—	—	P.E. Anaemia
11:00	"		11:00	—	3033	L.	L.	—	—	P.F.
—	Assisted breech		1:50	0:40	3402	L.	L.	—	—	
11:10	Spont.		16:25	0:35	2831	L.	L.	—	—	
5:15	Low forceps		4:15	0:55	2268	L.	L.	—	—	
9:15	"		9:15	1:25	2863	L.	L.	—	—	
6:50	Manual rotation and mid forceps		6:50	1:00	2807	L.	L.	—	—	
15:45	Spont.		15:45	0:05	3175	L.	L.	—	—	
10:00	Low forceps		10:00	1:15	3062	L.	L.	—	—	
18:00	Spont.		18:00	0:05	2948	L.	L.	—	—	
36:00	L.S.C.S.		36:00	—	2892	L.	L.	—	—	
25:00	"		25:00	—	3147	L.	L.	—	—	
—	"		—	—	3600	L.	L.	—	—	P.E.
8:00	Manual rotation and mid forceps		7:55	0:15	2948	L.	L.	—	—	
19:50	Low forceps		19:50	0:55	3260	L.	L.	—	—	
5:00	L.S.C.S.		5:00	—	2580	L.	L.	—	—	
3:00	Spont.		2:30	0:30	3033	L.	L.	—	—	
—	L.S.C.S.		—	—	3118	L.	L.	—	—	Anaemia.
15:00	Low forceps		14:30	1:10	2920	L.	L.	—	—	
—	L.S.C.S.		—	—	3260	L.	L.	1	—	
6:30	"		6:30	—	3572	L.	L.	—	—	
17:00	"		17:00	—	3175	L.	L.	—	—	
13:00	"		13:00	—	3175	L.	L.	—	—	
10:35	Low forceps		10:20	0:15	3175	L.	L.	—	—	Anaemia.
—	"		15:10	1:35	3515	L.	L.	—	—	"
6:20	Spont.		5:30	0:50	2750	L.	L.	—	—	
12:15	"		11:55	0:15	3260	L.	L.	—	—	
20:30	"		20:00	0:35	2750	L.	L.	—	—	
—	L.S.C.S.		—	—	2778	L.	L.	1	—	
19:00	Low forceps		18:50	0:10	2948	L.	L.	—	—	
—	"		7:30	1:05	3147	L.	L.	—	—	P.E.
6:20	Spont.		6:00	0:20	2240	L.	L.	—	—	
26:25	Low forceps		26:00	0:25	2098	L.	L.	—	—	Manual removal of placenta.
10:00	L.S.C.S.		10:00	—	2892	L.	L.	—	—	
—	"		—	—	3629	L.	L.	3	—	
6:30	Spont.		6:10	0:20	2013	L.	N.N.D.	—	—	
—	"		1:05	0:10	2637	L.	L.	1	—	
6:35	Low forceps		6:15	0:20	2381	L.	L.	—	—	
4:45	Spont.		4:30	0:15	2863	L.	L.	—	—	P.E. Anaemia.
16:00	L.S.C.S.		—	—	3289	L.	L.	—	—	
—	Low forceps		4:25	0:10	2296	L.	L.	—	—	Trichomonas vaginitis.
9:00	L.S.C.S.		—	—	3459	L.	L.	—	—	
—	Spont.		10:55	0:25	4281	L.	L.	—	—	Anaemia
8:00	"		8:10	1:05	2608	L.	L.	—	—	
—	L.S.C.S.		—	—	3147	L.	L.	—	—	
—	Low forceps		8:10	0:40	3062	L.	L.	—	—	
5:00	L.S.C.S.		5:00	—	3090	L.	L.	1	—	
10:00	"		10:00	—	3175	L.	L.	—	—	
11:20	"		10:35	0:45	2977	L.	L.	—	—	Anaemia.
5:00	"		5:00	—	2948	L.	L.	—	—	
—	Mid forceps		15:45	1:15	3175	L.	L.	—	—	Anaemia.
13:00	L.S.C.S.		13:00	—	3906	L.	L.	—	—	
13:00	"		13:00	—	3289	L.	L.	—	—	
5:10	Low forceps		5:00	0:10	3175	L.	L.	—	—	
—	L.S.C.S		—	—	2438	L.	L.	—	—	Anaemia.
10:00	"		10:00	—	2970	L.	L.	—	—	Hepato-splenomegaly.
—	Elective L.S.C.S.		—	—	2835	L.	L.	—	—	
22:15	Low forceps		22:15	0:10	2722	L.	L.	—	—	

CONTRACTED PELVIS (Continued)

Reg. No.	Age	Height (inches)	Previous Pregnancies			Maturity	Type of Pelvis	Radiological Results Pelvic Measurements				Outlet T.O.	Position
			Before	After	28 wks.			28 wks.	O.C.	I.D.	I.S.		
MULTIPARA													
BOOKED													
6651/59	33	57¼	1	2	38	Gynaecoid	10.0	10.2	—	10.8	—	—	R.O.A.
7391/59	27	56	—	2	41	"	9.3	11.2	8.9	10.7	11.3	—	L.O.A.
7728/59	32	62½	—	4	40	Platypelloid	9.7	12.7	9.3	12.5	10.0	—	L.O.A.
7941/59	33	61¾	1	1	41	Gynaecoid	11.2	11.2	—	10.7	10.0	—	R.O.A.
7975/59	27	59¾	—	1	36	"	10.0	11.5	—	11.0	—	—	R.O.A.
8042/59	27	57½	1	1	42	"	9.5	11.9	9.0	10.3	—	—	L.O.T.
8047/59	36	62½	—	3	41	Platypelloid	9.2	12.0	—	—	—	—	R.O.A.
8055/59	25	57¼	—	1	40	"	9.6	12.5	—	10.8	—	—	R.O.T.
8067/59	31	57¾	—	3	39	Anthropoid	11.4	10.8	—	—	—	—	R.O.T.
8075/59	38	61½	—	2	41	Gynaecoid	11.6	11.6	8.5	11.2	—	—	L.O.A.
8133/59	39	57	—	1	40	"	9.0	11.8	—	10.8	—	—	R.O.A.
47/60	29	57	1	2	37	Platypelloid	9.8	12.5	—	—	—	—	L.O.T.
55	29	57¾	—	1	40	Gynaecoid	9.6	11.7	—	—	—	—	R.O.A.
71	37	58	1	4	39	"	10.6	11.8	9.6	11.0	—	9.0	R.O.A.
90	30	56	—	1	39	"	10.5	11.2	—	—	—	—	L.O.A.
102	31	56¾	—	2	39	"	8.4	11.7	—	8.1	9.8	—	R.O.A.
106	27	62	—	2	44	Platypelloid	8.1	12.5	—	11.2	10.3	—	L.O.A.
151	28	58¼	—	2	40	"	9.6	12.4	—	10.5	—	—	R.O.A.
218	36	60½	—	2	39	Gynaecoid	10.0	11.8	—	10.9	10.1	—	L.O.T.
251	42	NS	—	3	37	Platypelloid	9.6	12.1	—	12.2	—	—	L.O.A.
257	34	59¾	—	3	41	Gynaecoid	10.5	11.8	9.3	11.9	—	—	R.O.T.
258	22	58	—	1	44	Gynaecoid- Anthropoid	12.2	11.65	—	12.9	—	—	L.O.A.
266	39	60¾	—	2	38	Gynaecoid	10.4	11.8	—	11.1	8.6	—	L.O.A.
270	36	56¾	1	5	40	"	9.7	11.3	—	11.5	—	—	L.O.A.
305	30	60½	—	4	40	"	9.5	12.1	—	12.0	—	—	R.O.P.
328	35	57¾	—	1	40	Platypelloid	9.2	12.6	—	10.4	—	—	L.O.A.
336	28	56	—	1	42	Gynaecoid	9.3	12.5	—	9.5	—	—	R.O.P.
345	27	55	—	1	38	"	9.2	11.4	11.5	9.8	—	—	L.O.A.
388	35	61	—	3	42	"	9.9	11.6	—	12.6	—	—	L.O.A.
446	35	58½	3	1	40	"	11.4	11.8	—	10.7	—	—	L.O.A.
449	27	59	—	1	39	Platypelloid	9.4	13.2	10.0	11.7	—	—	L.O.A.
452	30	58¼	—	1	40	Gynaecoid	10.5	11.4	—	9.3	—	—	L.O.T.
520	37	58¾	—	6	43	Platypelloid	9.6	12.6	—	—	—	—	L.O.A.
550	31	56¾	1	2	39	"	9.6	12.9	—	11.3	—	—	L.O.P.
557	29	61	—	1	39	Gynaecoid	9.7	12.4	—	11.4	—	—	R.O.A.
586	31	62½	—	2	40	"	11.6	11.8	—	9.7	—	—	L.O.A.
627	32	58	—	5	38	"	9.5	12.0	—	10.2	—	—	R.O.T.
632	32	62¼	—	3	40	Anthropoid	12.5	11.8	—	11.0	—	—	L.O.A.
655	31	58¼	—	1	39	Platypelloid	8.2	12.7	—	10.8	—	—	R.O.A.
678	24	58	—	2	40	Gynaecoid	10.6	11.3	—	11.0	—	—	L.O.A.
712	30	58½	—	3	39	"	11.0	11.9	—	9.5	—	—	L.O.A.
756	34	59	—	3	38	"	10.0	11.4	8.4	10.1	—	—	R.O.A.
779	27	59	—	1	40	"	9.0	12.4	8.9	10.7	—	—	L.O.A.
786	26	60½	—	1	38	Anthropoid	12.6	11.4	—	12.3	—	—	L.O.A.
815	33	57½	—	2	39	Gynaecoid	10.9	11.85	9.15	10.0	—	—	L.O.A.
852	43	59½	2	5	43	Gynaecoid- Platypelloid	9.95	12.8	—	11.1	—	—	L.S.A. L.O.P.
885	38	63½	—	4	41	Anthropoid	2.3	11.9	—	13.4	—	—	L.O.A.
921	33	59	—	1	40	Gynaecoid	10.98	12.45	9.53	9.6	—	—	R.O.P.-O.A.
924	31	57	—	4	41	"	11.0	11.95	—	10.5	—	—	R.S.A.
926	24	59	—	1	40	Gynaecoid- Anthropoid	11.65	11.4	—	10.8	—	—	L.O.A.
929	34	58¼	—	1	41	Gynaecoid	11.0	11.0	—	10.6	—	—	R.O.P.
940	26	58½	—	1	40	"	11.0	11.4	—	10.5	—	—	R.O.T.-O.A.
953	35	58	—	5	38	Platypelloid	9.9	13.1	—	12.5	11.5	—	R.O.A.
995	40	59½	5	2	38	Gynaecoid- Anthropoid	11.6	11.6	—	10.9	—	—	L.O.A.
1001	23	57	—	1	42	Gynaecoid	10.2	11.65	9.15	11.0	—	—	L.O.A.
1092	25	55	—	1	38	"	10.4	11.9	8.8	9.6	—	—	R.S.A.
1238	28	56	—	1	44	"	9.8	12.0	—	10.6	—	—	L.O.A.
1163	43	59¼	1	7	37	Platypelloid	9.9	12.8	9.8	9.9	—	—	R.O.A.
1164	34	58¾	1	4	39	Gynaecoid	9.7	12.4	9.9	10.2	—	—	R.O.T.
1176	25	56	—	1	44	"	10.2	11.4	9.1	10.5	—	—	R.O.A.
1212	35	57	—	5	39	"	9.1	12.0	—	10.5	10.4	—	R.O.A.
1277	28	58¾	—	3	?	Platypelloid	8.2	11.8	—	—	9.9	—	R.O.A.
1309	23	58¾	1	2	37	Gynaecoid	9.5	11.9	—	10.6	—	—	L.O.T.
1335	34	55	1	2	37	Naegele's pelvis	8.3	9.5	7.5	10.5	—	—	R.S.A.
1372	29	59½	—	5	36	Platypelloid	9.7	12.8	—	10.6	—	—	L.O.A.
1384	29	60½	—	1	42	Gynaecoid	9.9	12.4	—	11.7	—	—	R.O.A.
1387	27	63	—	1	40	"	11.0	12.3	—	9.8	—	—	L.O.A.
1416	36	59	—	1	40	Platypelloid	9.4	12.5	10.5	10.3	—	—	L.O.A.
1431	30	52½	—	4	37	Gynaecoid	10.2	11.7	9.5	10.7	—	—	R.O.T.

Surgical Induction	Management of labour (hours)		Length of labour (hrs.)		Weight of child (grams)	Result		Morbidity (days)	Prenatal C.S.	Remarks
	Method of delivery		1st stage	2nd stage		M	C			
—	—	Caesarean hysterectomy	—	—	2640	L.	L.	—	1	Cardiac disease.
—	—	Elective L.S.C.S.	—	—	2770	L.	L.	—	2	" "
—	—	" "	—	—	3500	L.	L.	—	—	" "
—	—	" "	—	—	2997	L.	L.	—	1	P.P.S.
—	—	" "	—	—	2580	L.	L.	N.N.D.	1	Hydrop foetalis.
—	3:10	L.S.C.S.	3:10	—	3827	L.	L.	—	—	—
—	—	Elective L.S.C.S.	—	—	2778	L.	L.	—	2	P.P.S.
—	4:00	L.S.C.S.	4:00	—	3289	L.	L.	—	—	—
—	—	" "	—	—	2685	L.	L.	—	1	P.P.S.
—	—	Elective L.S.C.S.	—	—	2800	L.	L.	—	2	" "
—	7:00	L.S.C.S.	—	—	3550	L.	L.	—	—	—
—	—	Elective L.S.C.S.	—	—	2430	L.	L.	—	2	—
—	—	" "	—	—	2920	L.	L.	—	1	—
—	—	L.S.C.S.	6:30	1:15	3400	L.	L.	—	—	—
—	—	Elective L.S.C.S.	—	—	3147	L.	L.	—	1	—
—	—	" "	—	—	3345	L.	L.	—	2	P.P.S.
—	—	" "	—	—	2693	L.	L.	—	1	—
—	—	" "	—	—	3402	L.	L.	—	2	—
—	—	Spont.	13:35	0:20	3450	L.	L.	—	—	—
—	4:10	" "	4:10	0:10	3062	L.	L.	—	—	—
—	—	" "	3:20	0:10	4200	L.	L.	—	—	Anaemia.
—	—	" "	2:30	0:10	2790	L.	L.	—	—	—
—	—	Elective L.S.C.S.	—	—	2722	L.	L.	—	—	—
—	—	Spont.	3:10	0:15	2980	L.	L.	—	—	—
A.R.M.	8:35	" "	8:25	0:10	3100	L.	L.	—	1	—
—	—	Elective L.S.C.S.	—	—	2778	L.	L.	—	1	Pulmonary T.B. Anaemia. Syphilis.
—	14:00	L.S.C.S.	14:00	—	3370	L.	L.	—	—	—
—	—	" "	7:00	—	3118	L.	L.	—	1	—
—	6:30	Spont.	6:20	0:10	3140	L.	L.	—	—	—
—	2:45	" "	2:35	0:10	3800	L.	L.	—	—	—
—	—	Elective L.S.C.S.	—	—	3402	L.	L.	—	—	—
—	13:00	L.S.C.S.	13:00	—	3280	L.	L.	—	—	Syphilis.
—	—	Spont.	22:50	0:05	3150	L.	L.	—	—	External version at onset of labour. P.P.S.
—	—	" "	11:15	0:15	2950	L.	L.	—	—	—
—	—	" "	9:15	0:25	2880	L.	L.	—	—	—
—	—	" "	11:05	0:05	3050	L.	L.	—	—	Anaemia.
—	6:00	L.S.C.S.	5:30	1:00	4100	L.	L.	—	—	—
—	—	Spont.	4:40	0:25	2630	L.	L.	—	—	—
—	—	Elective L.S.C.S.	—	—	2400	L.	L.	—	—	—
—	—	" "	—	—	3260	L.	L.	—	—	—
—	—	Spont.	4:15	0:05	3000	L.	L.	—	—	—
—	—	L.S.C.S.	—	—	2722	L.	L.	—	1	Syphilis.
—	1:00	Spont.	1:00	0:10	2722	L.	L.	—	—	—
—	—	" "	6:30	0:20	3570	L.	L.	—	—	—
—	—	Assisted breech	3:20	0:05	3100	L.	L.	—	—	—
—	—	L.S.C.S.	—	—	3970	L.	L.	—	—	Prolapse of cord.
—	—	Spont.	4:30	0:05	3530	L.	L.	—	—	—
—	—	Mid forceps	10:55	0:55	2917	L.	L.	—	—	—
—	—	Assisted breech	3:15	0:05	3400	L.	L.	—	—	—
—	—	Low forceps	11:05	0:20	2740	L.	L.	1	—	P.E.
—	15:30	L.S.C.S.	15:30	—	3033	L.	L.	—	—	—
—	2:15	Manual rotation and mid forceps	2:15	0:20	3402	L.	L.	—	—	—
—	—	L.S.C.S.	—	—	2835	L.	L.	—	1	Anaemia.
—	—	Spont.	0:07	0:03	2098	L.	L.	—	—	Pulmonary T.B. Anaemia.
—	4:50	" "	4:50	0:10	2495	L.	L.	—	—	—
—	—	L.S.C.S.	2:30	—	2863	L.	L.	—	—	—
—	—	" "	—	—	2948	L.	L.	—	—	Mitral stenosis.
—	—	Mid forceps	3:05	0:10	3062	L.	L.	—	—	—
—	3:10	Spont.	3:10	0:05	2977	L.	L.	—	—	—
—	—	L.S.C.S.	—	—	2948	L.	L.	—	1	P.P.S.
—	—	" "	—	—	3402	L.	L.	—	1	—
—	—	Elective L.S.C.S.	—	—	3515	L.	L.	—	3	—
—	—	L.S.C.S.	3:00	—	1984	L.	L.	—	—	Accidental haemorrhage. Anaemia.
—	—	" "	—	—	2722	L.	L.	—	1	P.P.S.
—	—	Spont.	2:10	0:05	3175	L.	L.	—	—	Anaemia.
—	12:40	" "	12:40	0:10	2920	L.	L.	—	—	Congenital elongation of cervix.
—	—	L.S.C.S.	—	—	3856	L.	L.	—	1	—
—	—	" "	—	—	3175	L.	L.	—	—	—
—	10:00	" "	10:00	—	3685	L.	L.	—	—	Subtotal hysterectomy for secondary P.P.H. Anaemia.

CONTRACTED PELVIS (Continued)

Reg. No.	Age	Height (inches)	Previous Pregnancies		Maturity	Type of Pelvis	Radiological Results				Outlet	Position
			Before	After			O.C.	T.D.	I.S.	A.P.		
1454	27	57½	—	4	36	Platypelloid	9.4	12.3	—	11.8	—	R.O.A.
1464	32	58½	1	3	40	"	9.7	12.5	—	10.8	—	L.O.A.
1469	24	58¾	—	3	40	Gynaecoid	10.1	11.5	—	10.0	—	R.O.A.
1496	26	57¾	—	1	40	Anthropoid	11.7	11.0	—	11.0	—	R.O.A.
1610	39	59½	—	1	38	Gynaecoid	9.6	12.1	8.3	10.6	—	R.O.A.
1623	28	54	—	1	40	"	9.4	11.7	—	9.5	—	R.O.A.
1642	27	57½	—	2	41	"	11.0	11.4	—	11.5	9.6	L.O.A.
1672	29	56½	—	1	40	"	9.8	11.2	—	11.2	—	L.O.A.
1705	29	55½	—	2	44	Anthropoid	11.4	10.8	—	10.7	—	R.O.A.
1715	39	57	—	5	38	Gynaecoid	9.8	12.3	—	10.4	10.1	R.O.P.
1721	29	57½	—	2	36	"	11.1	11.0	—	—	—	R.O.P.
1747	30	60	—	2	40	"	11.2	11.3	—	10.6	—	L.O.A.
1790	21	57	—	1	36	"	10.6	10.9	—	11.8	—	L.O.A.
1839	37	59½	—	7	39	"	10.7	11.8	9.6	13.0	—	L.O.T.
1846	35	57	—	2	40	"	10.8	11.6	—	11.0	—	R.S.A.
1858	33	58½	—	1	39	"	9.2	10.0	—	10.3	—	R.O.A.
1870	27	57¾	—	1	38	"	9.1	11.9	—	11.2	—	L.O.A.
1892	33	58¾	—	2	39	"	8.7	11.6	9.3	10.9	9.1	R.O.P.
1902	26	60¼	—	1	41	"	9.7	12.0	—	9.4	—	L.O.P.
1921	28	57	—	1	40	Platypelloid	9.2	12.7	—	10.7	—	L.O.A.
1952	29	56	—	1	39	Gynaecoid	11.2	11.4	—	10.7	—	L.O.A.
1954	30	58¼	—	4	42	Platypelloid	9.1	12.1	—	10.4	—	R.O.A.
1993	35	59	—	2	39	Gynaecoid	11.7	11.4	9.3	11.2	—	L.O.A.
2028	28	57½	—	1	39	Anthropoid	12.0	11.7	9.8	10.1	—	R.S.A.
2031	27	58½	—	1	37	Gynaecoid	9.7	12.3	—	10.0	9.0	R.O.A.
2037	42	57	1	12	39	"	10.0	11.8	9.0	11.3	—	L.O.A.
2044	24	56½	—	1	40	"	10.6	11.4	9.6	10.8	—	L.O.A.
2056	24	56¼	—	2	?	Platypelloid	8.3	13.5	—	10.1	11.4	R.O.A.
2073	29	58	2	4	41	Anthropoid	11.7	11.7	9.9	11.1	—	L.O.A.
2112	26	56	—	1	40	Gynaecoid	9.3	11.4	—	—	—	L.O.A.
2128	27	57½	—	4	38	"	9.7	11.5	9.6	9.9	—	R.O.P.
2136	27	54¼	—	2	39	Gynaecoid	9.3	11.0	9.0	9.5	—	L.S.A.
2137	29	56	—	1	40	Platypelloid	9.7	12.9	10.2	10.0	—	R.O.A.
2163	26	57½	—	1	40	Anthropoid	11.4	11.2	9.1	11.7	—	L.O.A.
2192	38	57½	1	4	40	Gynaecoid	10.3	11.5	—	—	9.7	L.O.A.
2229	24	57¾	—	1	39	Platypelloid	9.0	12.2	—	9.8	—	L.S.A.
2279	29	58¾	—	1	40	Gynaecoid	11.4	11.8	—	11.0	—	R.O.A.
2367	23	56¾	—	2	38	"	11.0	11.6	9.7	11.5	—	L.O.A.
2381	27	58½	—	3	33	Platypelloid	9.8	13.3	—	11.6	—	R.O.P.
2390	26	58	1	2	35	Gynaecoid	9.5	11.1	9.0	10.7	—	L.O.A.
2414	32	60¼	—	1	39	"	11.0	11.5	8.8	12.0	—	L.O.A.
2417	32	56¼	1	2	41	"	9.0	11.8	—	—	—	R.S.A.
2439	28	56¾	—	1	39	"	9.2	12.3	9.8	10.3	—	R.O.A.
2442	34	59	—	5	39	"	10.2	11.9	9.6	11.0	—	L.S.A.
2471	26	57	—	2	31	"	10.2	10.25	—	9.7	—	R.O.A.
2499	32	57¼	—	1	43	"	10.7	11.6	—	9.9	10.0	R.O.A.
2530	41	60	3	4	41	"	9.3	12.1	—	10.5	—	L.O.A.
2531	31	59½	1	6	36	"	11.0	11.3	—	9.0	—	L.O.A.
2534	21	57	—	2	38	"	9.0	11.9	—	9.2	—	R.O.A.
2570	27	55	1	2	39	"	9.3	10.8	8.2	10.3	—	R.O.P.
2588	31	55¼	—	7	44	"	8.8	11.8	—	11.3	—	R.O.A.
2603	32	53¾	—	2	41	"	8.9	12.1	—	10.5	—	L.O.A.
2604	30	56½	—	1	40	"	9.9	10.6	—	10.4	—	L.O.A.
2608	34	55½	1	7	41	"	9.8	11.5	—	11.8	—	R.O.T.
2640	38	57¾	—	2	42	"	10.5	12.2	8.2	12.0	—	L.O.A.
2644	32	58½	—	1	41	"	11.5	11.7	—	10.3	—	R.O.A.
2687	40	56½	—	4	41	Platypelloid	9.6	12.5	10.2	10.9	—	R.O.A.
2702	24	56½	—	2	41	"	9.1	12.1	—	9.9	—	L.S.A.
2820	35	59	—	1	39	Gynaecoid	9.8	12.1	—	10.3	9.5	L.O.A.
2836	31	59¾	—	1	40	"	9.7	11.9	8.4	11.7	—	R.O.P.
2861	34	58	—	2	39	"	10.1	11.85	—	—	—	R.O.A.
2885	29	56¼	—	5	37	Platypelloid	8.6	12.8	—	11.3	—	L.O.A.
2917	34	57½	—	1	41	Gynaecoid	8.9	11.8	8.3	10.0	—	L.O.A.
2936	31	58¾	—	5	42	"	10.9	11.8	10.0	11.8	—	L.O.A.
2955	39	57½	—	7	44	Platypelloid	9.4	12.5	—	11.1	—	L.O.A.
2957	24	58½	—	1	40	Gynaecoid	9.65	11.7	—	10.4	—	R.O.A.
2966	30	58	—	2	38	Platypelloid	9.5	12.3	—	10.5	—	R.O.A.
2970	29	57	—	2	42	Gynaecoid	10.6	11.7	—	11.7	—	L.O.A.
2980	26	NS	—	2	34	"	10.3	11.5	8.8	—	—	R.O.A.
3083	24	58½	—	5	37	"	9.9	11.9	9.6	11.4	—	L.O.A.
3085	28	59	—	2	38	"	9.2	11.9	—	10.3	—	L.O.A.
3112	44	54¾	—	11	37	"	10.1	10.9	9.1	11.3	—	L.O.A.
3125	31	57	—	1	39	"	10.7	11.8	—	10.9	—	R.O.A.
3145	27	57	—	2	39	"	9.2	12.2	—	—	—	L.O.T.
3176	24	56	—	2	40	Anthropoid	10.1	10.5	—	—	9.0	R.O.A.
3195	25	57	—	2	39	Gynaecoid	10.7	11.0	—	10.4	—	L.O.A.
3203	28	58½	—	5	40	Naegle's pelvis	10.0	9.9	6.2	12.2	9.0	L.O.A.

Surgical Induction	Management Trial of labour (hours)		Length of labour (hrs)		Weight of child (grams)	Result		Morbidity (days)	Pretensions (days)	Remarks
	Method of delivery	1st stage	2nd stage	M		C				
—	6:00	Spont.	5:45	0:25	3544	L.	L.	—	—	
—	—	L.S.C.S.	—	—	3945	L.	L.	—	1	P.P.S.
—	—	"	—	—	2920	L.	L.	—	2	"
—	—	Spont.	5:30	0:10	3459	L.	L.	—	—	
—	—	"	3:55	0:10	3459	L.	L.	—	—	Anaemia.
—	—	L.S.C.S.	—	—	2972	L.	L.	—	1	
—	—	Elective L.S.C.S.	—	—	3005	L.	L.	—	—	Moniliais.
—	—	L.S.C.S.	—	—	3495	L.	L.	—	1	
—	—	Spont.	4:30	0:10	3515	L.	L.	—	—	
—	—	L.S.C.S.	—	—	3175	L.	L.	—	—	
—	15:00	"	15:00	—	3459	L.	L.	—	—	
—	—	"	—	—	2722	L.	L.	—	1	
—	4:15	Spont.	4:10	0:05	3260	L.	L.	—	—	Anaemia.
—	6:00	L.S.C.S.	6:00	—	3856	L.	L.	—	—	"
—	—	Elective L.S.C.S.	—	—	3289	L.	L.	—	2	P.P.S.
—	—	L.S.C.S.	—	—	2892	L.	L.	—	1	
—	—	Spont.	11:30	0:10	1700	L.	N.N.D.	—	—	Anaemia. Hydramnios. Anencephaly.
—	—	L.S.C.S.	—	—	2948	L.	L.	—	2	
—	6:00	"	6:00	—	3289	L.	L.	—	—	
—	—	Elective L.S.C.S.	—	—	3090	L.	L.	—	1	Pulmonary T.B. (active).
—	—	"	—	—	3203	L.	L.	—	1	
—	4:00	Low forceps	4:15	0:20	3742	L.	L.	—	—	P.P.S.
—	1:50	"	1:50	—	2892	L.	L.	—	—	
—	—	Assisted breech	5:10	0:20	2863	L.	L.	—	—	
—	—	L.S.C.S.	—	—	2637	L.	L.	1	—	Accidental haemorrhage. Placenta praevia type I.
—	10:00	"	—	—	3500	L.	L.	—	—	P.P.S.
—	2:00	Spont.	1:55	0:05	2722	L.	L.	—	—	Anaemia.
—	—	"	19:50	0:15	2466	L.	L.	—	—	
—	1:30	"	1:30	0:50	3042	L.	L.	—	—	
—	—	L.S.C.S.	—	—	2600	L.	L.	—	1	
—	8:00	"	—	—	2693	L.	L.	—	—	Pulmonary T.B. Prolapse of uterus.
—	7:35	L.S.C.S.	—	—	2102	L.	L.	—	—	
—	2:35	Spont.	2:20	0:05	3037	L.	L.	—	—	
—	3:40	"	7:25	1:15	2948	L.	L.	—	—	
—	—	L.S.C.S.	—	—	2807	L.	L.	—	1	
—	6:50	"	—	—	2835	L.	L.	—	—	Bronchopneumonia.
—	16:15	Spont.	16:15	0:15	2835	L.	L.	—	—	
—	—	L.S.C.S.	—	—	3232	L.	L.	—	2	
—	—	"	—	—	3345	L.	L.	—	3	
—	—	C.C.S.	—	—	3062	L.	L.	—	2	Previous rupture of uterus. P.P.S.
—	—	Low forceps	3:15	0:15	3090	L.	L.	—	—	
—	—	L.S.C.S.	—	—	2977	L.	L.	—	2	
—	—	"	—	—	3345	L.	L.	—	1	
Stripping & A.R.M.	22:00	"	22:00	—	2862	L.	L.	—	—	Anaemia.
—	—	"	—	—	1786	L.	L.	—	1	
—	—	"	—	—	3402	L.	L.	—	—	
—	0:40	Mid forceps	0:40	0:05	3317	L.	L.	—	—	
—	6:30	"	6:30	0:15	2892	L.	L.	—	—	
—	—	L.S.C.S.	—	—	2835	L.	L.	—	2	P.P.S.
—	—	"	—	—	3289	L.	L.	—	—	
—	15:50	Spont.	15:50	0:05	3060	L.	L.	—	—	
—	—	L.S.C.S.	—	—	3140	L.	L.	—	—	
—	20:10	Spont.	20:10	0:30	2750	L.	L.	—	—	
—	12:00	L.S.C.S.	—	—	3827	L.	L.	—	—	
—	3:00	"	3:00	—	2665	L.	L.	—	—	
—	—	Elective L.S.C.S.	—	—	3515	L.	L.	—	—	
—	—	Spont.	4:12	0:08	2863	L.	L.	—	—	
—	—	L.S.C.S.	—	—	2890	L.	L.	—	—	
—	—	Low forceps	1:35	0:25	2948	L.	L.	—	1	Anaemia.
—	17:00	L.S.C.S.	17:00	—	3175	L.	L.	—	—	P.E.
—	—	"	—	—	2897	L.	L.	—	2	
—	2:00	Spont.	2:00	—	3208	L.	L.	—	—	
—	—	L.S.C.S.	—	—	2438	L.	L.	—	1	
—	12:00	Spont.	12:00	0:05	2835	L.	L.	—	—	
—	8:57	L.S.C.S.	7:15	1:42	3575	L.	L.	1	—	
—	—	Spont.	4:00	0:20	3742	L.	L.	—	—	
—	—	Elective L.S.C.S.	—	—	3175	L.	L.	—	1	
—	—	Spont.	4:15	0:10	3232	L.	L.	—	—	Anaemia.
—	—	"	22:25	0:10	2608	L.	L.	—	—	
—	—	L.S.C.S.	—	—	2693	L.	L.	—	1	
—	—	Elective L.S.C.S.	4:00	—	2835	L.	L.	1	1	
—	3:30	L.S.C.S.	—	—	3912	L.	L.	—	—	
—	—	"	—	—	2948	L.	L.	—	1	
—	—	"	—	—	3118	L.	L.	—	2	
—	—	Elective L.S.C.S.	—	—	2557	L.	L.	3	2	P.P.S.
—	—	L.S.C.S.	—	—	2900	L.	L.	—	2	"
—	—	"	—	—	3317	L.	L.	—	2	T.B. Hip. P.P.S.

CONTRACTED PELVIS (Continued)

Reg. No.	Age	Height (inches)	Previous Pregnancies		Maturity	Type of Pelvis	Radiological Results Pelvic Measurements					Position
			Before 28 wks.	After 28 wks.			O.C.	T.D.	I.S.	A.P.	Outlet T.O.	
3225	27	57½	—	2	38	Gynaecoid-Anthropoid	10.8	11.1	9.2	12.0	—	L.O.A.
3231	30	59½	—	4	36	Platypelloid	9.4	13.0	10.3	12.0	—	R.O.A.
3272	27	59½	—	5	?	Gynaecoid	10.95	11.68	—	10.9	—	R.O.A.
3311	39	58	1	5	40	"	10.7	11.9	—	12.0	—	L.O.A.
3366	28	58	—	1	39	"	9.6	11.1	—	10.0	—	R.O.A.
3389	30	NS	—	2	42	"	9.8	12.3	10.0	11.3	—	R.O.A.
3406	23	54¾	—	1	38	"	10.1	11.5	8.7	10.0	—	R.O.T.
3408	39	58¾	—	1	39	Platypelloid	9.5	12.4	—	11.3	—	R.O.A.
4320	29	59¾	—	3	42	"	9.0	12.3	9.7	10.0	—	L.O.T.
3422	40	60	1	7	42	"	9.0	13.1	9.2	11.2	—	R.O.A.
3449	36	59	—	1	40	"	9.6	12.2	—	10.4	9.5	L.O.T.
3480	30	60¼	—	2	39	"	8.8	12.2	—	10.7	10.8	R.O.A.
3485	29	57½	—	1	40	Gynaecoid	9.9	11.2	—	9.2	—	L.O.A.
3492	24	55	1	2	38	"	10.1	11.1	—	11.5	10.3	L.O.P.
3508	33	59	—	6	?	"	11.5	11.5	8.8	10.7	—	L.O.P.
3552	21	60	—	1	41	"	10.4	11.5	—	10.5	—	L.O.A.
3613	31	56¼	—	1	39	"	9.2	11.7	—	10.2	—	L.O.A.
3620	29	55	—	1	40	Platypelloid	9.6	12.5	10.2	11.2	—	L.O.A.
3626	27	60	—	1	42	Gynaecoid	11.4	11.8	9.3	11.4	—	L.O.A.
3631	27	58	—	2	38	"	9.75	12.6	8.6	10.5	—	R.O.A.
3643	39	60	—	1	40	Platypelloid	8.8	12.3	—	9.8	—	R.O.T.
3652	30	55	—	5	40	Gynaecoid	10.0	11.9	—	10.5	—	L.O.A.
3705	24	56¼	—	1	37	"	9.9	10.55	—	9.3	—	L.O.A.
3715	28	58	—	1	41	"	10.3	11.3	—	10.05	—	L.O.A.
3726	39	59	—	8	34	"	10.3	11.8	9.9	10.3	—	L.O.A.
3736	31	56½	1	4	39	"	10.0	11.8	8.6	12.3	—	L.O.P.
3745	31	61¼	1	3	43	"	10.7	11.8	9.6	10.9	—	R.O.T.
3775	33	58½	—	1	40	Anthropoid-Gynaecoid	11.0	10.6	7.5	11.0	—	R.O.P.
3783	38	60¾	—	8	41	Platypelloid	9.4	12.4	10.1	10.0	—	L.O.T.
3863	29	60¼	—	4	43	"	9.3	12.6	11.0	11.0	—	R.O.A.
3875	25	53½	1	1	38	Gynaecoid	9.5	11.0	—	10.2	—	R.O.T.
3939	32	58½	—	4	37	Platypelloid	8.4	11.5	9.5	11.7	—	R.O.T.
9949	21	59	—	1	42	Gynaecoid	9.5	11.3	—	10.6	—	L.O.A.
3960	38	N.S.	—	3	42	Anthropoid	11.4	10.7	8.5	10.5	—	R.O.A.
3980	32	60	—	1	39	"	12.0	11.6	9.6	12.5	—	L.S.A.
4014	24	58¾	—	1	38	Gynaecoid	10.0	11.7	—	10.2	10.1	L.O.A.
4017	30	59	1	3	40	Platypelloid	9.5	12.4	—	11.4	—	L.O.A.
4056	34	57	—	1	38	Gynaecoid	9.7	11.6	—	11.2	—	R.O.A.
4057	30	59	—	4	38	"	10.7	11.9	10.1	11.0	—	L.O.A.
4058	34	60	—	1	39	"	10.6	11.6	8.6	9.8	—	L.O.A.
4132	32	62½	—	3	40	"	9.7	12.0	8.7	10.1	—	L.O.P.
4194	27	58	1	1	38	Anthropoid	11.9	11.4	9.0	11.1	8.7	R.O.A.
4258	34	56¼	—	4	38	Gynaecoid	8.9	11.8	—	—	—	R.O.A.
4260	38	59	—	5	37	"	9.0	11.9	9.25	9.5	—	R.O.A.
4286	27	58¼	—	1	40	"	10.4	11.3	9.6	10.1	—	R.O.A.
4132	30	55½	—	3	33	"	8.6	11.1	—	—	—	L.O.A.
4386	32	59½	—	1	37	"	11.3	11.4	9.2	11.2	—	L.O.A.
4389	34	61½	—	1	33	"	11.0	11.6	8.6	11.0	—	L.O.A.
4419	23	57¼	—	1	41	Anthropoid	12.2	10.9	9.0	11.3	—	L.O.A.
4471	32	N.S.	—	1	39	Gynaecoid	10.6	11.6	9.5	11.0	—	R.O.P.
4492	24	56¼	—	2	36	"	8.9	12.4	—	10.4	—	L.O.A.
4505	21	58	—	1	40	"	10.4	12.3	9.3	9.6	—	R.O.A.
4510	29	60	—	2	37	Platypelloid	9.0	13.2	—	10.5	—	R.O.P.
4539	25	57	—	4	41	Gynaecoid	10.8	11.8	9.6	10.7	—	R.M.P.
4590	31	58	1	1	39	"	9.8	12.9	—	10.5	—	R.O.A.
4626	24	58½	—	2	41	"	10.1	11.8	9.2	10.9	—	L.O.A.
4690	25	58	—	1	39	Platypelloid	8.6	11.5	9.7	9.2	—	L.O.P.
4694	30	60½	2	1	45	Gynaecoid	11.4	11.4	9.6	11.4	—	R.M.P.
4707	25	58¾	—	2	?	"	10.0	11.6	9.6	9.5	—	R.O.A.
4708	33	57	—	1	38	"	9.2	10.8	—	9.5	—	R.O.P.
4710	32	58¾	—	3	42	"	10.5	11.4	—	12.0	10.8	R.O.T.
4786	41	48½	1	1	39	Deformed	12.0	10.5	—	10.8	—	L.O.A.
4789	34	53¼	—	1	40	Gynaecoid	9.5	11.1	—	10.0	—	Transverse
4828	26	58½	—	1	39	"	9.7	11.9	8.7	11.6	—	L.O.T.
4853	25	57½	—	1	40	Anthropoid	11.6	11.75	—	10.8	—	R.O.A.
4878	30	59½	1	5	38	Gynaecoid	11.3	11.4	8.8	12.0	—	L.O.A.
4906	36	57½	—	8	37	Platypelloid	8.9	13.3	10.1	12.0	—	L.O.A.
4943	32	55	—	4	42	Gynaecoid	9.7	10.8	8.8	9.5	—	BROW
4966	30	53½	—	2	39	"	10.2	10.8	7.6	11.5	—	R.O.T.
4989	30	60	—	2	39	"	10.0	11.7	—	10.8	—	R.O.P.
5072	22	56	—	1	38	"	10.3	11.5	—	9.1	—	R.O.P.-O.A.
5114	32	60	—	5	39	"	9.7	—	—	10.7	—	R.O.P.
5117	27	59½	—	1	41	"	9.5	12.3	—	10.5	—	R.O.A.

Surgical Induction	Management Trial of labour (hours)		Length of labour (hrs.)		Weight of child (grams)	Result		Morbidity (days)	Previous C.S.	Remarks
	Method of delivery	1st stage	2nd stage	M		C				
—	—	Spont.	7:00	0:05	3289	L.	L.	—	—	
—	5:35	"	5:35	0:10	2580	L.	L.	—	—	
—	4:10	Low forceps	4:10	0:10	3515	L.	L.	—	—	
—	11:30	L.S.C.S.	11:30	—	4802	L.	L.	—	—	
—	—	"	—	—	2920	L.	L.	—	—	Prolapse of cord when cervix 3 cm. dilated.
—	—	Spont.	2:15	0:20	2835	L.	L.	—	1	
—	—	Elective L.S.C.S.	—	—	2608	L.	L.	1	1	
—	—	"	—	—	3005	L.	L.	—	1	
—	—	L.S.C.S.	—	—	3030	L.	L.	1	1	P.P.S.
—	7:30	Spont.	7:30	0:15	3289	L.	L.	—	—	
—	—	Elective L.S.C.S.	—	—	2863	L.	L.	1	—	
—	—	L.S.C.S.	—	—	3600	L.	L.	—	2	P.P.S.
—	—	"	—	—	3203	L.	L.	—	—	
—	—	"	8:00	—	2835	L.	L.	—	1	
—	—	"	9:50	3:00	3827	L.	L.	2	—	
—	6:25	Spont.	6:20	0:05	3062	L.	L.	—	—	
—	7:00	"	6:55	0:05	2693	L.	L.	—	—	
—	—	L.S.C.S.	—	—	2551	L.	L.	—	1	
—	7:00	Spont.	7:00	0:05	3317	L.	L.	—	—	
—	—	L.S.C.S.	—	—	3005	L.	L.	1	1	Anaemia. Syphilis.
—	—	"	—	—	3062	L.	L.	—	1	
—	3:30	Spont.	3:30	0:10	3203	L.	L.	—	—	
—	—	L.S.C.S.	—	—	3175	L.	L.	—	1	
—	—	"	—	—	2722	L.	L.	—	1	
—	—	L.S.C.S.	15:25	1:45	3439	L.	L.	—	—	Anaemia.
—	5:50	Low forceps	6:52	0:58	2637	L.	L.	—	—	Uterine didelphys.
—	—	Manual rotation and mid forceps	3:10	0:50	3544	L.	L.	—	—	
—	5:00	L.S.C.S.	5:00	—	2977	L.	L.	—	—	Pulmonary T.B. Anaemia.
—	6:00	"	6:00	—	3856	L.	L.	—	—	Anaemia
—	—	Spont.	5:15	0:05	2381	L.	L.	—	—	"
—	—	L.S.C.S.	—	—	3118	L.	L.	—	—	
—	6:00	"	6:00	—	3459	L.	L.	—	1	
—	—	"	—	—	2296	L.	L.	1	—	
—	4:00	Spont.	3:50	0:10	3402	L.	L.	—	—	
—	—	Assisted breech	13:20	0:10	3374	L.	L.	—	—	P.E.
—	—	Spont.	1:50	0:10	2778	L.	L.	—	—	
—	—	L.S.C.S.	—	—	3600	L.	L.	—	1	
—	—	"	—	—	3203	L.	L.	—	1	
—	3:00	Spont.	3:00	0:25	3125	L.	L.	—	1	Previous C.S. for face presentation.
—	—	Low forceps	3:55	0:35	2892	L.	L.	—	—	
—	4:10	L.S.C.S.	4:10	—	3402	L.	L.	1	—	
—	1:10	Spont.	4:30	0:10	2977	L.	L.	—	—	
—	5:55	"	5:50	0:05	2268	L.	L.	—	2	
—	—	L.S.C.S.	—	—	3060	L.	L.	—	—	Prolapse of cord. Anaemia.
—	—	"	—	—	2722	L.	L.	—	1	
—	—	"	—	—	2353	L.	L.	—	2	Anaemia. P.P.S.
—	7:30	Low forceps	7:30	0:30	2495	L.	L.	—	1	
—	3:00	Spont.	3:00	0:10	1785	L.	L.	—	—	
—	5:10	"	5:00	0:10	3515	L.	L.	—	—	
—	—	L.S.C.S.	—	—	3090	L.	L.	—	1	
—	—	"	—	—	3033	L.	L.	—	—	Anaemia.
—	—	Spont.	2:15	0:10	3402	L.	L.	—	—	"
—	—	Elective L.S.C.S.	—	—	2722	L.	L.	—	1	
—	—	Spont.	10:00	0:20	2835	L.	L.	—	—	Anencephaly.
—	—	L.S.C.S.	—	—	3260	L.	L.	—	1	
—	—	"	—	—	3572	L.	L.	—	2	Anaemia. P.P.S.
—	—	Elective L.S.C.S.	—	—	3203	L.	L.	—	—	
—	4:00	Manual rotation and mid forceps	4:45	0:15	2693	L.	L.	—	—	
—	9:00	L.S.C.S.	9:00	—	3402	L.	L.	—	—	
—	—	"	—	—	2268	L.	L.	—	1	
—	—	Elective L.S.C.S.	—	—	3487	L.	L.	—	2	
—	—	C.C.S.	—	—	3402	L.	L.	—	1	
—	—	L.S.C.S.	—	—	2863	L.	L.	—	1	
—	6:55	Spont.	6:45	0:10	3629	L.	L.	—	—	
—	4:00	Low forceps	3:50	0:10	3629	L.	L.	—	1	
—	3:05	Spont.	3:00	0:05	3175	L.	L.	—	—	P.P.S.
—	—	L.S.C.S.	—	—	3600	L.	L.	—	—	
—	—	"	—	—	3374	L.	L.	—	—	
—	—	"	—	—	2948	L.	L.	—	1	
—	8:00	Spont.	8:50	0:30	2892	L.	L.	—	—	
—	—	L.S.C.S.	—	—	2381	L.	L.	—	1	
—	8:15	Spont.	8:10	0:05	3600	L.	L.	—	—	Anaemia. P.P.S. Ligation of left internal iliac artery.
—	—	L.S.C.S.	—	—	3544	L.	L.	—	1	P.P.H. Anaemia.

CONTRACTED PELVIS (Continued)

Reg. No.	Age	Height (inches)	Previous Pregnancies		Maturity	Type of Pelvis	Radiological Results Pelvic Measurements				Outlet T.O.	Position
			Before 28 wks.	After 28 wks.			O.C.	T.D.	I.S.	A.P.		
5154	26	57¼	—	1	42	Platypelloid	8.6	12.1	—	10.5	—	R.S.A.
5156	28	58	—	1	36	Gynaecoid	10.3	11.1	—	9.1	—	L.O.A.
5187	35	59½	—	7	41	Platypelloid	9.4	12.5	—	9.6	—	L.O.T.
5264	31	59	1	1	40	"	9.9	13.2	—	10.9	—	R.O.A.
5302	22	57½	—	1	40	Gynaecoid	9.0	11.8	—	9.4	—	R.S.A.
5308	26	59½	—	1	37	Platypelloid	8.8	14.8	—	11.2	—	L.O.A.
5312	27	61½	—	1	40	Android-Gynaecoid	10.5	10.8	—	—	10.3	L.O.T.
5367	29	61	—	4	39	Gynaecoid	9.5	12.3	—	11.0	—	R.O.P.
5396	27	60	—	1	40	Platypelloid	9.6	12.2	—	9.5	—	L.O.A.
5427	37	57	1	3	38	"	8.7	12.3	—	—	10.6	L.O.A.
5429	23	55¼	1	2	36	Gynaecoid	10.0	11.1	8.5	10.2	—	R.O.A.
5454	31	61	4	4	39	"	11.5	11.7	—	—	11.0	L.O.A.
5470	26	60	—	1	38	"	11.0	11.9	—	10.4	—	L.O.A.
5556	30	59¼	—	1	39	"	11.0	11.2	—	11.4	—	R.O.T.
5557	35	60	—	5	40	"	10.1	11.9	9.4	11.5	—	R.O.T.
5569	21	60	—	1	40	"	9.4	11.8	—	8.9	—	L.O.T.
5657	24	61	—	1	40	Anthropoid	11.0	11.35	—	10.3	—	R.O.A.
5659	21	57½	—	2	41	Gynaecoid	10.7	11.3	—	10.6	—	L.O.A.
5686	38	N.S.	—	9	41	Platypelloid	9.4	12.6	—	10.6	—	L.O.A.
5724	30	59	—	2	39	Gynaecoid	9.3	11.2	8.3	—	—	Transverse lie
5427	24	51¼	—	2	39	"	9.1	11.7	—	10.3	—	R.O.A.
5740	25	61	—	1	39	Gynaecoid	10.0	11.8	—	11.4	—	R.O.A.
5826	31	54¼	—	1	39	Android-Gynaecoid	11.5	11.1	—	10.0	—	R.O.A.
5832	23	57½	—	1	40	Gynaecoid	9.0	12.25	—	10.9	—	L.O.A.
5845	30	59	—	5	32	Platypelloid	9.0	12.0	9.25	11.4	—	Transverse
5890	31	56½	2	3	39	Gynaecoid	10.4	11.2	9.0	11.0	—	R.O.A.
5892	30	N.S.	—	1	41	"	11.9	11.9	—	11.3	—	R.O.P.-O.A.
5909	31	56	—	1	40	"	9.6	11.2	—	10.8	—	L.O.P.
5914	25	57½	—	1	40	"	10.5	11.8	10.0	11.4	—	L.O.A.
5931	34	N.S.	—	4	40	"	9.8	11.6	9.2	10.4	—	R.O.A.
5945	32	62½	1	4	37	Anthropoid-Gynaecoid	9.8	13.0	10.6	10.3	—	L.S.A.
5951	34	56¼	—	6	43	Gynaecoid	9.7	12.0	11.1	11.0	—	L.S.A.
6006	30	56	—	1	41	Platypelloid	8.3	11.3	—	9.3	—	L.O.A.
6047	36	58	—	6	35	Gynaecoid	8.9	12.1	—	11.1	—	R.O.P.
6115	29	55½	—	1	?	"	9.3	11.3	—	—	—	L.O.A.
6141	31	60½	1	4	41	Gynaecoid-Anthropoid	11.7	11.8	10.9	12.2	—	R.S.A.
6153	34	59¼	—	2	38	Platypelloid	9.8	12.8	9.3	10.0	—	L.O.A.
6170	34	55½	—	1	42	Gynaecoid	9.6	11.1	—	11.1	—	L.O.A.
6189	22	59	—	1	40	"	11.2	11.8	—	9.8	—	L.O.A.
6283	40	57	—	5	39	Platypelloid	8.5	12.2	9.7	10.0	—	R.O.A.
6287	30	57	—	1	39	Gynaecoid	9.4	11.8	—	11.3	—	L.O.A.
NON-BOOKED												
6567/59	25	61	—	3	?	Gynaecoid	11.7	11.7	—	10.9	—	L.O.A.
1508/60	29	58	—	2	44	Anthropoid	11.5	11.5	—	12.5	—	R.O.A.
1768	25	58½	—	1	?	"	11.7	11.2	9.1	10.9	—	R.S.A.
2117	30	N.S.	—	2	40	"	11.3	11.3	10.0	11.6	—	R.O.A.
2156	37	61½	—	4	39	Gynaecoid	11.7	11.4	8.7	10.8	—	L.O.A.
2489	23	59	—	1	46	Platypelloid	8.7	12.9	—	10.0	—	L.O.A.
3232	38	N.S.	—	3	40	"	11.3	11.4	9.1	11.0	—	L.O.A.
3263	30	N.S.	—	1	39	"	9.3	12.2	10.0	12.3	—	R.O.T.
5231	29	N.S.	—	3	41	"	8.4	12.1	9.1	10.4	—	L.O.T.
5490	31	58½	1	6	34	"	8.0	12.7	11.3	11.2	—	L.S.A.
5565	31	N.S.	—	2	40	"	8.4	12.3	10.3	10.5	—	L.O.T.
5751	36	N.S.	—	2	40	"	9.5	13.0	9.2	11.0	—	R.O.T.
5805	25	N.S.	—	3	42	"	9.6	13.5	—	11.4	—	L.O.P.
MULTIPLE PREGNANCY BOOKED												
628	28	58¼	—	1	40	Gynaecoid	9.1	12.15	—	10.4	—	{ R.O.A. L.O.A.
3381	37	59	—	7	41	"	9.4	12.1	9.4	10.8	—	{ L.O.A. L.O.A.
3702	25	61	—	2	36	Anthropoid	12.3	11.8	—	10.1	—	{ L.O.A. ?
4341	23	60	—	—	39	Gynaecoid	10.0	11.5	8.6	9.9	—	{ L.O.A. R.O.A.
4945	35	58¼	—	3	34	"	9.9	12.4	9.2	9.8	—	{ L.O.A. R.S.A.
NON-BOOKED												
4914	25	N.S.	—	—	37	Gynaecoid	9.5	11.8	8.5	11.5	—	{ L.S.A. L.S.A.

Surgical Induction	Management of labour (hours)		Length of labour (hrs.)		Weight of child (grams)	Result		Morbidity (days)	Previous C.S.	Remarks
	Method of delivery	1st stage	2nd stage	M		C				
—	4:00	L.S.C.S.	4:00	—	2835	L.	L.	—	—	
—	—	—	—	—	3118	L.	L.	—	1	P.P.S.
—	7:00	"	7:00	—	3069	L.	L.	—	—	
—	—	—	—	—	2778	L.	L.	—	1	P.P.S.
—	—	—	2:00	—	2977	L.	L.	—	—	
—	14:35	Low forceps	14:20	0:15	2920	L.	L.	—	—	
—	—	L.S.C.S.	—	—	3629	L.	L.	—	—	
—	3:30	"	—	—	3317	L.	L.	—	1	
—	—	—	—	—	3260	L.	L.	—	1	
—	4:25	Spont.	4:15	0:05	3179	L.	L.	—	—	
—	—	L.S.C.S.	—	—	2453	L.	L.	—	—	
—	4:00	Spont.	3:55	0:05	3147	L.	L.	—	1	
—	—	"	3:40	0:15	2948	L.	L.	—	—	Anaemia.
—	—	L.S.C.S.	—	—	3175	L.	L.	—	1	
—	—	"	—	—	3090	L.	L.	—	—	
—	—	"	12:00	—	2801	L.	L.	—	—	Syphilis.
—	—	"	—	—	2523	L.	L.	—	1	
—	—	"	—	—	2353	L.	L.	—	1	
—	—	Spont.	2:35	0:05	3345	L.	S.B.	—	—	Prolapse of cord. Anaemia.
—	—	Laparotomy	3:00	—	3090	L.	S.B.	—	2	Subtotal hysterectomy for rupture of uterus.
—	—	L.S.C.S.	—	—	2948	L.	L.	—	2	P.P.S.
—	—	"	—	—	3175	L.	L.	—	1	
—	—	"	—	—	2948	L.	L.	—	1	
—	—	"	—	—	3118	L.	L.	—	1	
—	—	"	3:30	—	2438	L.	L.	—	—	
—	—	Spont.	2:45	0:10	3289	L.	L.	—	—	P.P.S.
—	7:00	"	7:35	0:10	3289	L.	L.	—	—	
—	—	L.S.C.S.	—	—	3232	L.	L.	1	—	Anaemia.
—	—	Spont.	8:45	0:10	2665	L.	L.	—	—	
—	—	"	13:20	0:05	2835	L.	L.	—	—	
—	—	Assisted breech	12:55	0:10	3430	L.	L.	—	—	
—	—	L.S.C.S.	—	—	3742	L.	L.	—	—	
—	—	"	—	—	3430	L.	L.	—	—	
—	9:00	"	9:00	—	3459	L.	L.	—	1	
—	—	"	—	—	2863	L.	L.	—	1	
—	—	Assisted breech	12:30	0:05	3062	L.	L.	—	—	
—	6:00	Mid forceps	6:00	0:10	3118	L.	L.	—	—	
—	—	L.S.C.S.	—	—	3685	L.	L.	—	1	
—	3:00	Spont.	3:10	0:10	3130	L.	L.	—	—	
—	—	"	1:10	0:05	2722	L.	L.	—	—	
—	—	"	0:40	0:05	2948	L.	L.	—	—	
—	—	Elective L.S.C.S.	—	—	3005	L.	L.	—	—	Habitual abortion — Shirodkar operation done in 1959.
—	—	Spont.	7:55	0:05	3345	L.	L.	—	—	
—	—	L.S.C.S.	—	—	2722	L.	L.	—	—	Unclassified toxæmia.
—	—	Spont.	2:30	0:10	2268	L.	L.	—	—	Diabetes mellitus. Schizophrenia.
—	12:00	Low forceps	12:00	0:40	2572	L.	L.	—	—	
—	—	L.S.C.S.	—	—	3515	L.	L.	1	—	
—	5:35	Spont.	5:35	0:10	3289	L.	L.	—	—	
—	—	L.S.C.S.	—	—	2268	L.	L.	—	—	
—	—	"	8:00	4:20	2900	L.	L.	—	—	
—	—	Breech extraction	5:00	0:50	2381	L.	S.B.	—	—	
—	—	L.S.C.S.	15:00	—	3175	L.	L.	—	—	
—	4:00	"	4:00	—	3884	L.	L.	—	—	
—	—	"	—	—	3515	L.	L.	—	2	P.P.S.
—	—	L.S.C.S.	—	—	{ 2300	{ L.	{ L.	—	—	Previous hydrops foetalis.
—	—	{ Spont.	—	—	{ 2550	{ L.	{ L.	—	—	
—	—	{ Spont.	0:20	0:05	{ 2693	{ L.	{ L.	—	—	Anaemia. P.P.S.
—	—	{ Spont.	—	—	{ 2778	{ L.	{ L.	—	—	
—	—	L.S.C.S.	—	—	{ 1871	{ L.	{ L.	—	2	
—	—	{ Spont.	—	—	{ 1921	{ L.	{ L.	—	—	
Stripping	11:00	"	11:00	—	{ 2523	{ L.	{ L.	—	2	P.E.
—	—	{ Breech assisted	4:30	0:20	{ 2240	{ L.	{ L.	—	—	
—	—	{ Spont.	—	—	{ 2070	{ L.	{ L.	—	—	
—	—	{ Breech assisted	—	—	{ 2410	{ L.	{ N.N.D.	—	—	
—	—	L.S.C.S.	—	—	{ 2438	{ L.	{ L.	—	—	
—	—	{ Spont.	—	—	{ 1070	{ L.	{ L.	—	—	

TABLE 26

ANTEPARTUM HAEMORRHAGE

Total No. of cases	189
Booked	137
Non-booked	52
Primiparae	49
Multiparae	140
Incidence	3%
Total No. of babies	193
Multiple pregnancies	4
Maternal death	—
S. B.	16 (8.29%)
N. N. D.	18 (10.17%)
P. N. D.	32 (16.58%)

TYPE OF A.P.H.:—

Accidental A. P. H.	60 (31.75%)
Placenta Praevia	48 (25.40%)
A. P. H. of unknown origin	45 (23.81%)
A. P. H. from extra-placental cause	36 (19.05%)

TABLE 27

ACCIDENTAL ANTEPARTUM HAEMORRHAGE

Total No. of cases	60
Booked	39
Non-booked	21
Primiparae	8
Multiparae	52
Incidence:—	
Over all	9.53%
Among A.P.H.	31.75%
Total No of babies	64
Multiple pregnancies	4
Maternal death	—
S. B.	13 (20.31%)
N. N. D.	11 (21.57%)
P. N. D.	24 (37.5%)
CAUSES WHERE PRESENT:—	
Toxaemia	8
Premature separation of placenta in 2nd twin	2
Unknown	50

AGE GROUP:—

<i>Age Group</i>	<i>Primiparae</i>	<i>Multiparae</i>
20 & under	—	1
21 — 25	4	6
26 — 30	4	13
31 — 35	—	13
36 — 40	—	12
41 & over	—	7
	8	52

MANAGEMENT OF ACCIDENTAL HAEMORRHAGE IN SINGLE PREGNANCY:—

	<i>No. of cases</i>
Non-interference	23
A.R.M.	18
L.S.C.S.	9
A.R.M. + L.S.C.S.	2
A.R.M. + Hysterectomy	2
A.R.M. + L.S.C.S. + Hysterotomy	2
Total	56
Number of cases requiring blood transfusion	16
Number of cases requiring fibrinogen	9

METHOD OF DELIVERY & FOETAL LOSS IN SINGLE PREGNANCY:—

<i>Method of delivery</i>	<i>No. of cases</i>	<i>Foetal Loss</i>		
		<i>S.B.</i>	<i>N.N.D.</i>	<i>P.N.D.</i>
Spontaneous	37	8	7	15
Forceps	6	—	1	1
C.S.	11	2	1	3
C.S. + Hysterectomy	2	2	—	2
Total	56	12	9	21

ACCIDENTAL ANTEPARTUM HAEMORRHAGE (Continued)

Reg. No.	Age	Previous Pregnancies		Maturity (wks.)	At admission	At delivery	Condition on admission	If in labour	Cause of A.P.H. if known	Treatment	Blood transfusion	Amount of bleeding		Result	Remarks
		Before 28 wks	After 28 wks									Concealed (ml.)	Revealed (ml.)		
PRIMIPARA															
BOOKED															
156	28	—	—	?	?	?	Good	Yes	—	—	—	150	—	L.	Syphilis.
880	28	—	—	34	35	35	"	"	A.R.M. & Low forceps	—	—	250	250	L.	Pulmonary T.B.
1658	27	—	—	35	35	35	Fair	"	L.S.C.S.	1 pint	—	360	—	L.	Fibrinogen.
1706	23	—	—	P41	P41	P41	Good	"	—	—	—	100	100	L.	S.B.
3161	22	—	—	39	39	39	"	"	—	—	—	50	10	L.	Low forceps.
6271	28	—	—	36	36	37	"	No	A.R.M.	—	—	5	35	L.	
NON-BOOKED															
4831	24	—	—	30	30	31	Good	No	—	—	—	68	300	L.	
MULTIPARA															
BOOKED															
7778/59	42	—	9	About 16	27	30	Good	No	—	—	—	20	115	L.	Anaemia.
7340	31	—	1	31	31	31	"	Yes	—	—	—	30	50	L.	Diagnosed after delivery
317	33	1	10	30	39	39	"	No	A.R.M.	—	—	100	305	L.	Low forceps. Anaemia.
631	37	1	8	38	38	38	"	"	—	—	—	600	—	L.	Anaemia.
849	35	—	6	37	37	37	Poor	Yes	A.R.M. + subtotal hysterectomy	12 pints	—	1,000	50	L.	Hydrocephrosis.
1069	22	—	1	39	38	37	Good	Yes	—	—	—	50	150	L.	Fibrinogen. P.E. P.P.H
1160	30	—	1	38	38	38	"	No	—	—	—	100	60	L.	
1284	33	—	5	37	37	37	Poor	"	L.S.C.S.	2 pints	—	480	800	L.	Anaemia. P.P.S.
1309	23	1	2	37	37	37	Good	Yes	"	1 pint	—	100	300	L.	Fibrinogen
1417	26	—	1	32	32	32	"	"	A.R.M.	—	—	60	300	L.	Contracted pelvis.
1663	35	—	12	41	41	41	"	"	—	—	—	200	—	L.	
1678	31	2	4	36	36	36	"	No	L.S.C.S.	—	—	50	120	L.	
2031	27	1	1	37	37	37	"	"	"	—	—	30	100	L.	Contracted pelvis.
2733	21	1	1	30	30	30	"	"	A.R.M. & pitocin drip	—	—	300	50	L.	
2830	42	1	7	38	38	38	"	Yes	—	—	—	10	200	L.	Essential hypertension.

ACCIDENTAL ANTEPARTUM HAEMORRHAGE (Continued)

Reg. No.	Age	Previous Pregnancies		Maturity (wks.)			Condition on admission	If in labour	Cause of A.P.H. if known	Treatment	Blood transfusion	Amount of bleeding		Result	Remarks
		Before 28 wks	After 28 wks	At Ist haem.	On admission	At delivery						Concealed (ml.)	Revealed (ml.)		
3080	29	—	6	35	35	37	Good	No	—	A.R.M.	—	150	75	L.	P.P.S.
3157	44	2	10	38	38	38	Fair	"	—	A.R.M. + L.S.C.S. + total hysterectomy	4 pints	630	1,500	L.	S.B. Fibrinogen.
3193	37	—	6	40	40	40	Good	Yes	—	—	—	5	300	L.	N.N.D.
3218	25	—	1	36	36	36	"	"	—	—	—	70	30	L.	Low forceps. Brecch.
3411	24	—	1	34	34	34	"	"	—	—	—	500	150	L.	S.B. Proclapse of cord. Syphilis.
3819	33	—	7	33	33	33	"	"	—	A.R.M.	—	660	40	L.	S.B. Anaemia.
3828	35	—	6	36	36	36	Fair	"	—	A.R.M.	2 pints	650	650	L.	S.B. Anaemia. Pulmonary T.B.
3886	42	—	5	41	41	41	Good	No	—	"	—	40	200	L.	N.N.D. Anaemia.
4086	37	—	9	30	30	30	Good	No	—	—	—	50	—	L.	N.N.D. B.O.A.
4197	35	—	1	32.6	32.6	32.6	"	Yes	—	—	2 pints	1,500	100	L.	L.
4634	37	—	5	38	38	38	Poor	No	—	L.S.C.S.	—	20	120	L.	L.
4958	31	—	4	34	34	34	Good	Yes	—	A.R.M.	—	25	70	L.	L.
5244	34	—	9	?	?	?	"	No	—	—	—	10	20	L.	N.N.D. Anaemia.
5853	29	—	1	Spotting since pregnancy	28	28	"	"	—	—	—	450	—	L.	L.
5990	39	1	4	—	38	38	"	Yes	—	A.R.M.	—	100	350	L.	L.
6144	36	—	6	39	39	40	"	No	—	A.R.M. + L.S.C.S.	—	—	—	L.	L.
NON-BOOKED															
7844/59	40	—	3	38	38	38	Good	Yes	Severe P.E.	A.R.M. + Mid forceps	—	—	50	L.	S.B. Fibrinogen.
190/60	30	—	3	38	38	38	Fair	"	—	A.R.M. + Pitocin drip + L.S.C.S.	8 pints	1,800	180	L.	S.B.
653	43	—	6	38	38	38	Good	No	Severe P.E.	A.R.M.	2 pints	300	50	L.	L.
1118	39	—	8	33	33	33	Poor	Yes	—	A.R.M. + subtotal hysterectomy	4 pints	1,650	350	L.	S.B. Fibrinogen.
1119	38	—	5	41	41	41	"	Yes	—	L.S.C.S.	8 pints	500	400	L.	S.B. " P.P.H.
1260	28	1	5	38	38	38	Good	No	—	A.R.M. + Low forceps	1 pint	300	300	L.	L.
1294	35	—	6	35	35	35	"	"	—	L.S.C.S.	—	70	50	L.	L.
1356	38	1	7	32	32	32	"	"	—	A.R.M. + L.S.C.S. + total hysterectomy	—	10	546	L.	L.
1553	41	1	9	35	35	35	Fair	Yes	—	—	7 pints	1,200	700	L.	S.B. Fibrinogen. Anaemia. P.P.H.

ACCIDENTAL ANTEPARTUM HAEMORRHAGE (Continued)

Reg. No.	Previous Pregnancies		Maturity (wks.)		Condition on admission	If in labour	Cause of A.P.H.	Treatment	Blood transfusion	Amount of bleeding		M.	Result	Remarks
	Before 28 wks	After 28 wks	At 1st adm.	On admission						Concealed (ml.)	Revealed (ml.)			
1863	20	—	30	30	Good	Yes	—	—	—	20	—	L.	L.	Anaemia.
2755	30	4	33	33	Fair	No	—	1 pint	—	200	400	L.	N N D.	Fibrinogen.
2769	28	1	28	28	Good	—	—	—	—	20	400	L.	S.B.	—
2839	33	2	38	38	—	—	—	—	—	10	150	L.	L.	—
2841	28	5	31	31	—	—	—	1 pint	—	20	810	L.	N.N.D.	—
2898	42	2	31	31	—	—	—	—	—	5	310	L.	L.	P.P.S.
3180	31	8	29	29	—	—	—	—	—	10	100	L.	L.	—
3769	27	5	38	38	—	—	—	3 pints	—	500	800	L.	L.	Pulmonary T.B.
5133	27	2	?	?	—	—	—	—	—	30	50	L.	L.	—
5133	37	10	31	31	—	No	—	—	—	—	—	L.	L.	—
MULTIPLE PREGNANCY														
BOOKED														
1327	27	—	244	244	Good	Yes	—	A.R.M.	1 pint	370	2500	L.	L.	Retroplacental clots in placenta of 1st twin.
3637	25	—	After delivery of 1st twin.	39	—	—	Separation of placenta	Breech extraction of 2nd twin	2 pints	—	1,000	L.	L.	P.E. Bleeding after delivery of 1st twin.
NON-BOOKED														
260	26	—	1	38	38	Fair	Yes	(1) Spont (2) L.S.C.S.	5 pints	100	—	L.	L.	Bleeding after delivery of 1st twin.
1329	38	2	6	32	32	Good	—	—	—	180	200	L.	L.	Transverse lie—2nd twin, external version & breech extraction. Retroplacental clots in placenta of 2nd twin.

TABLE 28

PLACENTA PRAEVIA

Total No. of cases	48
Booked	30
Non-booked	18
Primiparae	8
Multiparae	40
Incidence:—	
Over all	7.62%
Among A. P. H.	25.4%
Total No. of babies	48
Multiple pregnancies	—
Maternal death	—
S. B.	1 (2.08%)
N. N. D.	4 (8.51%)
P. N. D.	4 (8.33%)

TYPE AND FOETAL LOSS:—

Type	No. of cases	S.B.	N.N.D.	P.N.D.
I	13	—	1	1
II	13	—	1	—
III	10	1	1	2
IV	12	—	1	1
TOTAL	48	1	4	4

MANAGEMENT AND FOETAL LOSS:—

Management	No. of cases	S.B.	N.N.D.	P.N.D.
E.U.A. & A.R.M.	8	—	1	1
E.U.A. & L.S.C.S.	39	1	2	2
L.S.C.S. + Hysterectomy	1	—	1	1
TOTAL	48	1	4	4

Number of cases requiring blood transfusion 17

PLACENTA PRAEVIA (Continued)

Reg. No.	Age	Pregnancies Before 28 wks.	Pregnancies After 28 wks.	At 1st haem.	Maturity On adm.	At del.	Condition on admission	If in labour	Type 1, 2, 3, 4,	1 treatment	Blood transfu- sion (ml.)	Amount of bleeding (ml.)	Result M.	Remarks
PRIMIPARA														
BOOKED														
113	25	—	—	38	38	38	Good	No	4	L.S.C.S.	1500	1000	L.	Anaemia. P.E.
146	27	—	—	44	44	44	"	"	1	"	—	—	L.	Chance finding.
2079	39	—	—	35	35	36	"	"	2	"	—	400	L.	Contracted pelvis.
2085	37	—	—	39	39	40	"	"	1	"	—	150	L.	Anaemia.
3548	30	—	—	38	38	39	"	"	1	"	—	15	L.	P.E.
3853	30	—	—	16	36	41	"	"	3	"	—	—	L.	"
4301	23	—	—	38	38	38	"	"	2	"	2000	1500	L.	"
NON-BOOKED														
2757	30	—	—	30	30	30	Good	No	2	L.S.C.S.	500	520	L.	N.N.D.
MULTIPARA														
BOOKED														
332	41	7	7	31	31	34	Good	No	3	L.S.C.S.	500	225	L.	Anaemia.
425	34	—	—	41	38	41	"	"	1	A.R.M.	—	108	L.	"
798	38	1	1	39	39	39	"	"	1	L.S.C.S.	—	430	L.	Anaemia.
882	41	1	1	38	38	38	Fair	"	4	L.S.C.S.	1000	200	L.	"
1355	41	—	—	32	32	33	Good	No	3	L.S.C.S.	—	450	L.	"
1406	32	—	—	37	39	40	"	"	3	"	—	—	L.	"
1933	31	—	—	39	39	39	"	"	1	A.R.M.	—	100	L.	Anaemia.
1984	38	2	7	31	31	35	"	"	2	L.S.C.S.	—	90	L.	"
2023	28	—	—	35	35	40	"	"	2	"	—	100	L.	"
2037	42	1	12	39	39	39	"	"	1	"	—	5	L.	Contracted pelvis.
2266	40	—	—	34	34	34	"	"	3	"	—	355	L.	toaemia.
2835	33	—	—	37	37	37	"	"	2	"	—	100	L.	"
3659	31	—	—	39	39	39	"	"	1	"	—	100	L.	P.E.
3750	37	—	—	36	36	39	"	"	4	"	1500	140	L.	"
4636	28	1	3	43	43	43	"	"	1	E.U.A. & strapping	—	45	L.	"
4754	37	—	—	37	37	37	"	Yes	2	A.R.M.	—	200	L.	Anaemia.
5080	37	—	—	32	32	32	"	No	3	L.S.C.S.	500	420	L.	N.N.D.
5294	34	—	—	34	36	37	"	"	4—5	"	—	330	L.	P.P.S.
5518	23	—	—	39	39	39	"	Yes	1	E.U.A. & low forceps	—	200	L.	"

PLACENTA PRAEVI (Continued)

Reg. No.	Age	Previous Pregnancies Before 28 wks.	At 1st haem.	Maturity On adm.	At del.	Condition on admission	If in labour	Type 1,2,3,4	Treatment	Blood transfusion (ml.)	Amount of bleeding (ml.)	Result M.	C.	Remarks
5823	40	—	36	36	38	Good	No	3	L.S.C.S.	—	10	L.	L.	
5911	29	2	31	40	40	"	"	2	A.R.M.	—	1920	L.	L.	
6046	32	—	39	39	39	"	"	4	L.S.C.S.	—	100	L.	L.	
6098	38	—	36	36	36	"	"	3	"	—	500	L.	L.	
NON-BOOKED														
1099	41	2	10	40	40	Good	No	2	L.S.C.S.	—	950	L.	L.	
1761	27	—	27	33	34	"	"	4	"	1000	572	L.	L.	
1849	26	1	38	38	39	"	"	4	"	1500	520	L.	L.	P.E.
1929	26	—	34	34	35	"	"	4	"	500	430	L.	L.	"
2049	20	—	29	29	29	"	Yes	1	"	—	390	L.	N.N.D.	Breech assisted.
2157	27	—	40	40	40	"	"	1	E.U.A.	500	600	L.	L.	Breech extraction.
2402	37	—	35	38	38	"	No	2	L.S.C.S.	—	360	L.	L.	
3105	38	1	30	30	30	"	"	4	L.S.C.S. & hysterectomy	500	725	L.	N.N.D.	
3263	30	—	37	37	39	"	"	4	L.S.C.S.	—	380	L.	L.	Contracted pelvis.
3320	38	—	32	32	35	"	"	4	"	1000	1110	L.	L.	Syphilis.
3978	27	—	28	33	34	"	"	2	"	—	635	L.	L.	
4365	28	—	30	31	32	"	"	4	"	500	795	L.	L.	
4623	32	—	—	37	37	"	"	1	"	—	—	L.	L.	Chance finding.
4740	39	—	38	38	38	"	"	2	"	1500	980	L.	L.	Pulmonary T.B., Prolapse of cord.
5236	31	—	16	35	35	Good	No	2	"	1000	450	L.	L.	Suprarenal failure.
5730	31	—	37	37	37	"	Yes	2	"	—	400	L.	L.	P.F.S.
6095	30	2	37	37	37	"	No	3	"	1000	1000	L.	S.B.	

TABLE 29

A.P.H. OF UNKNOWN ORIGIN

Total No. of cases	45
Booked	33
Non-booked	12
Primiparae	14
Multiparae	31
Incidence	{ Over all 7.15%
	{ Among A.P.H. 23.81%
Total No. of babies	45
Multiple pregnancies	—
Maternal death	—
S. B.	2 (4.44%)
N. N. D.	3 (6.98%)
P. N. D.	4 (8.89%)

MANAGEMENT:—

No interference	38
E. U. A.	4
E. U. A. & A. R. M.	1
L. S. C. S.	2

METHOD OF DELIVERY AND FOETAL LOSS:—

<i>Method of Delivery</i>	<i>No of cases</i>	<i>S.B.</i>	<i>N.N.D.</i>	<i>P.N.D</i>
Spontaneous	33	1	3	3
Forceps	3	—	—	—
L.S.C.S.	6	—	—	—
Breech	3	1	—	1
Total	45	2	3	4

TABLE 30

A.P.H.—EXTRA-PLACENTAL CAUSES

Total No. of cases	36
Booked	35
Non-booked	1
Primiparae	19
Multiparae	17
Incidence	{ Over all 5.72%
	{ Among A.P.H. 19.05%
Total No. of babies	36
Multiple pregnancies	—
Maternal death	—
S. B.	—
N. N. D.	—
P. N. D.	—
 CAUSES:—	
Cervical erosion	30
Cervical Polyp	3
Varicose veins	2
Prolapse of uterus	—
Carcinoma of cervix	—
Vulval warts	1

TABLE 32

MANUAL REMOVAL OF PLACENTA

Total No. of cases	46
Booked	42
Non-Booked	4
Primiparae	21
Multiparae	25
Incidence	0.73%
Total No. of babies	47
Multiple pregnancies	1
Maternal death	—
Previous manual removal	1

CAUSES:—

	<i>Primiparae</i>	<i>Multiparae</i>	<i>Total</i>
Adherent placenta	16	24	40 (86.96%)
Retained placenta	3	1	4 (8.70%)
Bleeding	1	—	1 (2.17%)
Broken cord	1	—	1 (2.17%)
Total	21	25	46

AMOUNT OF BLEEDING:—

	<i>Primiparae</i>	<i>Multiparae</i>	<i>Total</i>
100 c.c. & less	10	1	11 (23.91%)
101 — 200	5	10	15 (32.60%)
201 — 300	1	5	6 (13.04%)
301 — 400	1	3	4 (8.70%)
401 — 500	1	—	1 (2.17%)
501 — 1000	3	2	5 (10.87%)
1001 & over	—	4	4 (8.70%)
Total	21	25	46

METHOD OF DELIVERY:—

	<i>Primiparae</i>	<i>Multiparae</i>	<i>Total</i>
Spontaneous	12	22	34 (73.91%)
Forceps	8	2	10 (21.74%)
Breech	1	1	2 (4.35%)

DURATION OF LABOUR

Prolonged labour	2	—	2 (4.35%)
2nd stage over 1 hr.	3	1	4 (8.7%)
3rd stage over 1 hr.	12	16	28 (60.9%)

TABLE 33

**TUMOURS ASSOCIATED WITH PREGNANCY
LABOUR OR PUERPERIUM**

Total No. of cases	30
Booked	29
Non-booked	1
Primiparae	11
Multiparae	19
Incidence	0.48%
Total No. of babies	30
Multiple pregnancies	—
Maternal death	—
S. B.	—
N. N. D.	—
P. N. D.	—

TYPE OF TUMOUR:—

Ovarian — Simple serous cystadenoma	8
Benign teratoma	5
Fibroma	1
Pseudomucinous cystadenoma	1
Uterine — Uterine fibroids	8
Carcinoma of cervix	2
Others — Lipoma of Posterior vaginal wall	1
Tubo-ovarian cyst	1
Parovarian cyst	3
Caesarean section	20
Chance finding at Abdominal operation	13

TABLE 34

MULTIPLE PREGNANCY

Total No. of pregnancies	67
Booked	55
Non-booked	12
Primiparae	21
Multiparae	46
Twin pregnancy	66
Triplets	1
Incidence	1.06%
Total No. of babies	135
Maternal death	1

MATERNAL COMPLICATIONS:—

	<i>No. of cases</i>	<i>Percentage of total</i>
Toxaemia of pregnancy	26	38.81
Pre-eclampsia	25	37.31
Unclassified	1	1.49
Anaemia	20	29.85
Hydramnios	3	4.48
Accidental haemorrhage	5	7.46

TWIN PREGNANCY

Total No. of cases	66
Binovular	34 (51.52%)
Uniovular	31 (46.97%)
Unknown	1 (1.52%)
Total No. of babies	132

FOETAL MORTALITY:—

	<i>First Twin</i>	<i>Second Twin</i>	<i>Total</i>
S. B.	2	1	3
N. N. D.	3	5	8
P. N. D.	5	6	11

PRESENTATION:—

	<i>No. of cases</i>	<i>Percentage</i>
Vertex-Vertex	35	53.03
Vertex-Breech	16	24.24
Breech-Vertex	5	7.58
Breech-Breech	1	1.52
Vertex-Transverse	2	3.03
Vertex-Compound	1	1.52
Vertex-Unknown	2	3.03
Unknown-Compound	1	1.52
Breech-Unknown	1	1.52
Unknown-Unknown	2	3.03

PRESENTATION AND FOETAL LOSS:—

<i>Presentation</i>	<i>First Twin</i>		<i>Second Twin</i>			<i>Total</i>	
	<i>No. of cases</i>	<i>Foetal loss</i>	<i>Percentage</i>	<i>No. of cases</i>	<i>Foetal loss</i>		<i>Percentage</i>
Vertex	56	5	8.93	40	2	5.0	96
Breech	7	—	—	17	2	11.76	24
Transverse	—	—	—	2	—	—	2
Compound	—	—	—	2	—	—	2
Unknown	3	—	—	5	2	40.0	8
Total	66	5	7.58	66	6	9.09	132

MULTIPLE PREGNANCY (Continued)

FOETAL BODY WEIGHT AND FOETAL LOSS:—

Weight	Foetal Loss						Total
	First Twin			Second Twin			
	No. of cases	Foetal loss	Percentage	No. of cases	Foetal loss	Percentage	
1,500 & under ..	2	2	100.00	3	1	33.33	5
1,501 — 2,000 ..	4	1	25.00	7	2	28.57	11
2,001 — 2,500 ..	29	1	3.45	18	3	16.67	47
2,501 — 3,000 ..	21	1	4.76	30	—	—	51
3,001 — over ..	9	—	—	8	—	—	17
Unknown				—	—	—	1
Total	66	5	7.58	66	6	9.09	132

MATURITY AND FOETAL LOSS:—

Maturity	No. of cases	Foetal Loss	
		1st Twin	2nd Twin
30 — 31 6/7 wks.	2	1	—
32 — 33 6/7	4	1	1
34 — 35 6/7	8	1	2
36 — 37 6/7	12	1	—
38 — 39 6/7	17	1	3
40 wks. & over	19	—	—
Unknown	4	—	—
Total	66	5	6

METHOD OF DELIVERY AND FOETAL LOSS:—

Method of delivery	First Twin		Second Twin	
	No. of cases	Foetal loss	No. of cases	Foetal loss
Spontaneous	45	4	26	1
C.S.	6	—	7*	1
Forceps—Low	12	1	11	1
Mid	—	—	3	—
Assisted breech	3	—	14	2
Breech extraction	—	—	2	1
Internal version & breech extraction ..	—	—	3	—
Total	66	5	66	6

INDICATIONS FOR C.S.:—

	No. of cases
Cervical dystocia	1
Prolonged 1st stage, precious baby	1
Foetal distress, acute hydramnios	1
Contracted pelvis	1
Previous C.S., contracted pelvis	1
2nd twin, transverse lie with prolapse of arm	1*
Severe contracted pelvis, both breech presentation	1
Total	7

* A second twin was admitted as an emergency.

MULTIPLE PREGNANCY (Continued)

Reg. No.	Age (inches)	Height (inches)	Pregnancies Before	After	Maturity	Presentation		Method of delivery		Sex	Weight (grams)		Type, if known	Result Child		Remarks
						1st twin	2nd twin	1st twin	2nd twin		1st	2nd		M.	2nd	
PRIMIPARA																
BOOKED																
7292/59	23	N.S.	—	—	?	L.O.A.	R.O.P.	Low forceps	Low forceps	F.	2250	2600	Uniovular	L.	L.	P.E. A.R.M. Anaemia.
68/60	28	59	—	—	38	R.O.A.	L.O.P.	L.S.C.S.	L.S.C.S.	M.	2290	2550	Binovular	L.	L.	P.E.
80	28	61½	—	—	39	R.O.A.	R.O.A.	Low forceps	Low forceps	M.	2430	2420	Uniovular	L.	L.	Anaemia.
195	28	59¼	—	—	38	L.O.A.	R.O.A.	Spont.	Spont.	F.	2150	2350	"	L.	L.	P.E. Mild concealed
694	24	62	—	—	41	R.O.T. →	R.O.T. →	Bipolar version and breech extraction	Bipolar version and breech extraction	F.	2683	3147	Binovular	L.	L.	accidental haemorrhage —2nd twin.
951	29	61	—	—	35	L.S.A.	R.O.A.	Spont.	Spont.	M.	2375	1814	"	L.	L.	R.O.A.
1044	23	60½	—	—	40	L.O.A.	L.O.A.	"	"	M.	3033	2637	Uniovular	L.	L.	Anaemia.
1383	23	60¼	—	—	36	L.O.A.	L.S.A.	"	"	M.	2466	2580	"	L.	L.	"
1953	25	64	—	—	38	L.O.A.	R.O.A.	Low forceps	Low forceps	M.	2523	2665	Binovular	L.	L.	"
2377	32	60½	—	—	30	L.O.A.	R.O.A.	Spont.	Spont.	M.	880	1240	Uniovular	L.	N.D.	"
2629	25	61	—	—	36	L.O.A.	L.O.A.	"	"	F.	2211	1673	"	L.	L.	P.E. Anaemia
3067	28	62	—	—	35	L.O.A.	L.S.A.	Low forceps	Assisted breech— extraction	M.	2381	2381	"	L.	L.	Hydranmios, Anaemia.
3347	32	58½	—	—	37	R.O.A.	R.O.A.	Spont.	Spont.	M.	2098	2296	"	L.	L.	Anaemia.
3637	25	60	—	—	39	L.O.A.	L.O.A.	"	Breech extraction	F.	2211	2353	"	L.	N.D.	P.E. Cervical erosion.
3674	21	58	—	—	41	R.S.P.	R.O.A.	Assisted breech— forceps for A.C.H.	Spont.	F.	2268	2750	Binovular	L.	L.	Accidental haemorrhage.
4130	35	62	1	—	34	?	?	L.S.C.S.	L.S.C.S.	M.	2211	2251	Uniovular	L.	L.	P.E. Prolonged labour.
4341	23	60	—	—	39	L.O.A.	R.O.A.	"	"	M.	2523	2240	"	L.	L.	P.E. Hydranmios, Contracted pelvis.
5347	23	57¼	—	—	38	L.O.A.	L.O.A.	Spont.	Spont.	F.	2183	1928	Binovular	L.	L.	"
NON-BOOKED																
1240	23	62	—	—	33	L.O.A.	L.O.P. →	Spont.	Spont.	M.	2296	2637	Uniovular	L.	L.	Anaemia.
2997	25	N.S.	—	—	39	L.O.P.	R.O.A.	Low forceps	Internal version and breech extraction	M.	2551	2410	Binovular	L.	S.B. N.N.D.	P.E. Prolapse of cord—2nd twin.
4872	23	N.S.	—	—	34	Compound presentation	Compound presentation	Spont.	Spont.	M.	2211	2665	Uniovular	L.	L.	"

MULTIPLE PREGNANCY (Continued)

Reg. No.	Age (months)	Height (inches)	Previous Pregnancies Before After 28 wks. 28 wks.	Maturity	Presentation		Method of delivery		Sex		Weight (grams)		Type, if known	Result Child		Remarks		
					1st twin	2nd twin	1st twin	2nd twin	1st twin	2nd twin	1st twin	2nd twin		M.	F.		1st	2nd
4529	29	62	—	4	41	L.S.A.	L.O.A.	Spont.	M.	M.	3289	3345	Binovular	L.	L.	"		
4559	31	58½	—	5	31	L.S.A.	R.O.P. → O.A.	Assisted breech	F.	F.	3402	3232	Uniovular	L.	L.	"		
4741	35	59	—	2	40	R.O.A.	L.O.T.	Spont.	F.	F.	2639	3090	Binovular	L.	L.	P.E., Anaemia.		
4833	41	N.S.	—	9	?	L.O.A.	R.S.P.	"	M.	M.	2722	2750	"	L.	L.	"		
4880	32	60	—	4	37	R.O.A.	L.O.A.	"	M.	M.	3090	2438	Uniovular	L.	L.	"		
4945	35	58½	—	3	34	L.O.A.	R.S.A.	"	M.	M.	2070	2410	Binovular	L.	L.	N.N.D. Contracted pelvis.		
5205	35	62½	—	5	33	R.O.A.	R.S.A.	"	F.	F.	1817	2268	"	D.	L.	L.		
5334	36	64½	—	4	42	R.S.A.	?	Assisted breech	F.	F.	2693	2948	"	L.	L.	L.		
5419	38	64	1	6	37	R.O.A.	L.O.T.	"	F.	F.	2268	2835	"	L.	L.	L.		
5585	32	N.S.	—	4	40	R.O.A.	R.O.T.	Mid forceps	F.	M.	2948	2920	"	L.	L.	L.		
5820	29	N.S.	1	4	37	R.O.A.	Compound-L.O.A.-atm	Low forceps	F.	F.	2240	2187	Uniovular	L.	L.	L.		
NON-BOOKED																		
260	26	N.S.	—	1	38	?	?	Spont.	F.	F.	?	1750	?	L.	L.	S.B.		
626	37	61	—	4	?	L.O.A.	R.O.A.	Low forceps	M.	F.	2350	2630	Binovular	L.	L.	L.		
1329	38	60¼	2	6	37	R.O.A.	L.A.A.	Spont.	F.	M.	2013	2381	"	L.	N.N.D.	L.		
1397	36	N.S.	—	3	38	R.O.A.	R.O.P.	Low forceps	M.	M.	2698	2977	Uniovular	L.	L.	L.		
2474	40	N.S.	2	9	33	L.O.A.	L.S.A.	Assisted breech	M.	F.	2920	2551	Binovular	L.	L.	L.		
2732	25	N.S.	—	2	41	L.O.A.	R.S.P.	Assisted breech—forceps	F.	F.	2665	2722	"	L.	L.	L.		
2882	31	N.S.	—	3	42	L.O.A.	L.A.A.	for A.C.H. Internal version	F.	F.	3260	3033	Uniovular	L.	L.	L.		
3381	37	59	—	7	41	L.O.A.	R.O.A.	Spont.	M.	M.	2693	2778	"	L.	L.	L.		
4914	25	N.S.	—	1	37	L.S.A.	L.S.A.	L.S.C.S.	F.	F.	2438	1070	Binovular	L.	L.	L.		
Triplets																		
No.	Age	Height (inches)	Previous Pregnancies Before After 28 wks. 28 wks.	Maturity	Presentation		Method of delivery		Sex		Weight (grams)		Type if known		Result Child		Remarks	
1244	38	N.S.	—	6	38	R.S.A.	L.O.A.	L.S.P.	Low forceps	F.	F.	1200	1150	1100	Uniovular	L.	L.	N.N.D. N.N.D. P.P.S.

MULTIPARA BOOKED

TABLE 35

PUERPERAL PYREXIA

Total No. of cases	410
Incidence	6.51%
Genital in Origin	108 (26.34%)
Extragenital in Origin	254 (61.95%)
Unknown in Origin	48 (11.71%)

PUERPERAL PYREXIA (GENITAL)

Total No. of cases	108
Booked	99
Non-booked	9
Primiparae	70
Multiparae	38
Incidence:—	
Overall	1.71%
Among puerperal pyrexia	26.34%
Maternal death	—

CAUSES:—

Uterine infection	83
Urinary tract infection + uterine	2
Perineal wound infection	23

METHOD OF DELIVERY:—

Spontaneous	67
Forceps	26
C. S. (1st operation)	15

RESPONSIBLE ORGANISMS:—

B. coli	28
Gram negative bacilli	2
B. fusiformis	2
Haemolytic streptococcus	6
Coagulase positive staphylococcus	12
Streptococci faecalis	5
B. proteus	2
Pseudomonas	1
Staphylococci pyogenes	9
Streptococci pneumonia	1
Streptococci pyocyanea	2
Streptococci pyogenes	14
Unknown	18
Mixed	6
Total	<u>108</u>

PUERPERAL PYREXIA (Continued)

PUERPERAL PYREXIA (EXTRA-GENITAL)

Total No. of cases	254
Booked	244
Non-booked	10
Primiparae	98
Multiparae	156
Incidence:—	
Overall	4.03%
Among puerperal pyrexia	61.95%
Maternal death	1
CAUSES:—	
Respiratory tract infection	21
Urinary tract infection	76
Breast engorgement	63
Post operative fever	56
Dehydration	11
Skin infection	2
Bacillary dysentery	5
Amoebic dysentery	1
Carious tooth	2
P.P.S. wound infection	3
C. S. wound infection	4
Haematoma of broad lig.	1
Parotitis	1
Paratyphoid	1
Mesenteric Thrombosis	1
Carcinoma of liver	1
Staph. septicaemia	1
Conjunctivitis	1
Typhoid	1
Blood transfusion reaction	2
Total	<u>254</u>

DETAILS OF MATERNAL DEATHS

1. Booked Case. Serial No. 5205/60 Age: 35 Gravida: 6 Para: 5.

First seen in antenatal clinic on 13/10/60 when 29½ weeks pregnant. Twin pregnancy diagnosed. No complaints. B.P. 120/80, pulse 76/m. No oedema. Urine clear.

Admitted for rest on 28/10/60 but went into premature labour on 7/11/60 (33 weeks pregnant). First twin (1817 gm.) delivered spontaneously but 2nd (2268 gm.) by assisted breech extraction.

Post partum sterilization was done on 9/11/60 under local anaesthesia. She complained of severe abdominal pain but no vomiting 33 hours afterwards. Temperature was 103°F and pulse 160/m. There was generalised tenderness of the abdomen and bowel sounds were absent. She was treated as a case of general peritonitis with Antibiotics and intravenous fluid. The pain disappeared and there was no vomiting 2 days later. But temperature and pulse were 99°F and 110/m respectively.

Seven days after the operation she again complained of abdominal pain. She was transferred to the Surgical Unit for laparotomy. At operation a loop of ileum, 47 inches in length was found to be gangrenous due to mesenteric thrombosis and was resected. An end-to-end anastomosis done. Patient stood the operation well and there was marked improvement for 2 days. On the 3rd day her condition suddenly deteriorated and she died.

Post-mortem: Another 5 feet of ileum found to be gangrenous as a result of extension of mesenteric thrombosis.

No other abnormalities.

2. Non-booked case. Serial No. 4469/60 Age: 28 Gravida: 1 Para: 0.

Admitted on 22/9/60 clinically 23½ weeks pregnant. She gave a history of productive cough and fever for 2 weeks, dyspnoea on exertion for 8 days and orthopnoea and oedema of ankles for 3 days. On further questioning, she said she had palpitation on exertion since 12 years of age.

Her general condition was very poor—she was orthopnoeic and cyanotic and the neck veins were engorged. There was oedema of the legs up to the knees and the liver was palpable 2 inches below the costal margin. B.P. 140/110, Pulse 110/m, Respiration 32/m (laborous). Heart greatly enlarged with apex beat at the 6th intercostal space. A rough systolic murmur was heard all over the precordium. Chest full of rales.

In spite of treatment her condition deteriorated rapidly and she died 8 hours after admission.

Post-mortem: Tight mitral stenosis.

Marked hypertrophy of right ventricle.

Bilateral pleural effusion, pulmonary oedema and patchy consolidation.

Chronic passive congestion in the liver and spleen.

TABLE 36

SUMMARY OF FOETAL LOSS

Total No Live-births and stillbirths 6,365

Total No. of Live-births 6,304

	S.B.		N.N.D.		P.N.D.	
	No. of cases	Incidence	No. of cases	Incidence	No. of cases	Incidence
Total No.	61	9.58	78	12.37	135	21.21
Booked	47	(per 1,000	68	(per 1,000	112	(per 1,000
Non-booked	14	L.B. & S.B.)	10	L.B.)	23	L.B. & S.B.)
Primiparae	22		25		45	
Multiparae	39		53		90	

CAUSES:—

1. Unknown	23	37.70%	24	30.77%	46	34.07%
Mature	10		3		13	
Premature	13		21*		33	
2. Birth Trauma	10	16.39%	7	8.97%	17	12.59%
Contracted pelvis	2		—		2	
Breech delivery	—		3		3	
Cord accident	4		1		5	
Ruptured uterus	1		—		1	
Acute asphyxia	—		1		1	
Obstructed labour	1		—		1	
Forceps delivery	2		2		4	
3. Toxaemia	—	—	6	7.69%	6	4.44%
Pre-eclampsia	—		5		5	
Eclampsia	—		—		—	
Unclassified	—		—		—	
Essential hypertension	—		—		—	
Chronic Nephritis	—		1		1	
4. A. P. H.	14	22.95%	16	20.51%	30	22.22%
Placenta praevia	—		2		2	
Accidental haemorrhage	13		11		24	
Unknown cause	1		3		4	
5. Foetal abnormalities	8	13.11%	15	19.23%	21	15.56%
Congenital heart	—		4		3	
Anencephaly	3		5		8	
Multiple anomalies	3		3		5	
Cyclops	1		—		1	
Exomphalos	—		1		1	
Achondroplasia	1		1		2	
Imperforate anus	—		1		1	

SUMMARY OF FOETAL LOSS (Continued)

6. Miscellaneous	6	9.84%	10	12.82%	15	11.11%
Hydrops	4		1		5	
Intrauterine pneumonia	2		1		3	
Gastro-enteritis	—		1		1	
Bronchopneumonia	—		5		5	
Kernicterus	—		1		—	
Carcinoma of breast with early induction	—		1		1	

* Including 14 with atelectasis, some probably with hyaline membrane.

BIRTH WEIGHT:—

<i>Birth Weight (grams)</i>	<i>Foetal Loss</i>		
	<i>S.B. No. of cases</i>	<i>N.N.D. No. of cases</i>	<i>P.N.D. No. of cases</i>
Under — 1,000	2	3	5
1,001 — 1,500	7	16	22
1,501 — 2,000	13	20	33
2,001 — 2,500	14	17	29
2,501 — 3,000	11	13	23
3,001 — 3,500	12	7	19
3,501 — 4,000	1	2	3
4,001 — 4,500	—	—	—
4,501 & over	—	—	—
Not recorded	1	—	1

METHOD OF DELIVERY:—

<i>Method of Delivery</i>	<i>Foetal Loss</i>		
	<i>S.B. No. of cases</i>	<i>N.N.D. No. of cases</i>	<i>P.N.D. No. of cases</i>
Spontaneous	32	49	80
Low Forceps	2	5	7
Mid Forceps	—	1	1
Breech Delivery	14†	10*	23
Caesarean Section	11	13	22
Caesarean Hysterectomy	2	—	2
Total	61	78	135

* Including 5 forceps of A.C.H. and including 2 cases delivered by Caesarean Section.

† Including 2 cases delivered by Caesarean Section.

TABLE 37

PREMATURE LIVE-BORN INFANTS

Total No. of cases	483
Booked	434
Non-booked	49
Primiparae	249
Multiparae	234
Incidence	76.62 per 1,000 L.B.
Neo-natal deaths	53
Neo-natal mortality rate (per 1,000)	109.73

MATURITY (WEEKS):—

	<u>No. of cases</u>	<u>Neo-natal death (loss per 1,000)</u>
Under 32 weeks	41	17 = 414.63
32 — 33 „	56	13 = 232.14
34 — 35 „	62	4 = 64.52
36 — 37 „	97	5 = 51.55
38 — 39 „	118	7 = 59.32
40 — 41 „	48	3 = 62.50
42 weeks & over	4	—
Unknown	57	4 = 70.18

BIRTH WEIGHT (GRAMS):—

	<u>No. of cases</u>	<u>Neo-natal death (loss per 1,000)</u>
Under — 1,000	4	3 = 750.00
1,001 — 1,500	22	15 = 681.82
1,501 — 2,000	72	19 = 263.89
2,001 — 2,500	385	16 = 41.56

CAUSES OF DEATH IN PREMATURE INFANTS:—

	<u>No. of cases</u>
Birth trauma	2
Bronchopneumonia	2
Acute asphyxia	3
Atelectasis	34
Congenital malformation	9
Unknown	3

TABLE 38

CONGENITAL MALFORMATIONS

	<u>Booked</u>	<u>Non-booked</u>	<u>Total</u>
Total No. of cases	54	2	56
Incidence	8.88 per 1,000 live and stillbirths.		
Among Live Births	45	1	46 (0.73%)
Among Still-births	9	1	10*(16.39%)
Among Premature Live Births	17	1	18*(3.73%)
Among neo-natal deaths	13	1	14*(17.95%)

TYPE OF MALFORMATION:—

	<u>No. of cases</u>
Cyclops	1
Anencephaly	8
Mongolism	2
Deformity of ear	2
Hypospadias	2
Deformity of limbs	8
Exomphalos	1
Cystic hygroma	1
Achondroplasia	2
Epidermolysis bullosa	1
Cavernous haemangioma	1
Gartner's cyst	1
Harelip and cleft palate	12
Oesophageal atresia	1
Malformation of anus	2
Congenital heart disease	5
Multiple anomalies	6

* Some were not the cause of death.

TABLE 40

NEO-NATAL INFECTIONS

Total No. of cases	72
Incidence	11.42 per 1,000 live births.
Neo-natal deaths	8 (all bronchopneumonia)
Neo-natal mortality rate	11.11%

TYPE OF INFECTIONS:—

Respiratory tract infection	37
Skin infection	8
Umbilical infection	11
Conjunctivitis	14
Bacillary dysentery	1
Mastitis	1

TABLE 41

HAEMOLYTIC DISEASES

Total No. of cases	3
Incidence	0.48 per 1,000 live births.


Type of haemolytic disease:—

Rh incompatibility	—
ABO incompatibility	3
Treatment given	Nil
Result of infants	All living

TABLE 42
X-RAY DEPARTMENT STATISTICS

Month	Chest	Bones	Abdomen	Pelvimetry	Placento- graphy	Cystography	Hysterosal- pingography	Intravenous Urography	Cervico- graphy	Barium Sivallou	Screening	Placenta Specimen	Cholangio- graphy
Jan.	366	1	17	124	7	1	4	1	—	—	1	—	—
Feb.	627	10	14	131	4	5	10	2	—	—	—	—	—
Mar.	703	12	18	124	3	7	3	—	—	—	—	—	—
Apr.	630	9	10	116	8	5	7	1	—	1	—	—	—
May	813	12	12	103	12	8	6	2	—	—	—	—	—
June	727	5	25	103	12	7	5	1	—	1	—	9	—
July	733	6	18	117	13	5	5	—	—	—	1	—	1
Aug.	712	—	29	135	11	7	5	1	—	—	—	—	—
Sept.	742	3	18	118	6	9	4	1	—	—	—	—	—
Oct.	624	5	14	100	3	4	3	1	1	—	—	—	—
Nov.	602	8	10	95	3	4	9	2	—	—	—	—	—
Dec.	577	10	8	101	4	6	6	—	—	—	2	—	—
TOTAL	7856	81	193	1367	86	68	67	12	1	2	4	9	1

X17171540



S 610

Tsan Yuk Hospital.
Clinical report of ...
for the year ...

